
Global Emergence of Trimethoprim/Sulfamethoxazole Resistance in *Stenotrophomonas maltophilia* Mediated by Acquisition of *sul* Genes

Mark A. Toleman,* Peter M. Bennett,† David M.C. Bennett,† Ronald N. Jones,‡ and Timothy R. Walsh*

Trimethoprim/sulfamethoxazole (TMP/SMX) resistance remains a serious threat in the treatment of *Stenotrophomonas maltophilia* infections. We analyzed an international collection of 55 *S. maltophilia* TMP/SMX-sensitive (S) (n = 30) and -resistant (R) (n = 25) strains for integrons; *sul1*, *sul2* and *dhfr* genes; and insertion element common region (ISCR) elements. *sul1*, as part of a class 1 integron, was detected in 17 of 25 TMP/SMX-R. Nine TMP/SMX-R strains carried *sul2*; 7 were on large plasmids. Five TMP/SMX-R isolates were positive for ISCR2, and 4 were linked to *sul2*; 2 others possessed ISCR3. Two ISCR2s were adjacent to *floR*. Six TMP/SMX-S isolates harbored novel ISCR elements, ISCR9 and ISCR10. Linkage of ISCR3, ISCR9, and ISCR10 to *sul2* and *dhfr* genes was not demonstrated. The data from this study indicate that class 1 integrons and ISCR elements linked to *sul2* genes can mediate TMP/SMX resistance in *S. maltophilia* and are geographically widespread, findings that reinforce the need for ongoing resistance surveillance.

Nosocomial *Stenotrophomonas maltophilia* are intrinsically resistant to a plethora of antimicrobial agents that severely limit commonly used empiric standard antimicrobial therapies. *S. maltophilia* is resistant to many β -lactams, β -lactamase inhibitors, and aminoglycosides (1,2). A recent survey of SENTRY (www.jmilabs.com) Antimicrobial Surveillance Program isolates indicated that the newer fluoroquinolones demonstrated good efficacy; the most active were levofloxacin (6.5% resistance) and

gatifloxacin (14.1%) (3). Furthermore, the resistance to the polymyxins (20%–32%) is higher than observed in *Pseudomonas aeruginosa* (3,4). Because of low resistance levels (\approx 5%), trimethoprim/sulfamethoxazole (TMP/SMX) remains the therapy of choice worldwide. A recent study encompassing data from Europe, Latin America, and North America indicates that the level of resistance to TMP/SMX is 3.8%; however, previous studies indicate that the level is higher in Latin America than North America (5,6). Although surveillance studies are few, resistance to TMP/SMX appears to be emerging, and recent in vitro modeling studies have shown that combination therapies of TMP/SMX plus ciprofloxacin and TMP/SMX plus tobramycin exhibit a greater killing capacity than TMP/SMX alone (7,8).

S. maltophilia exhibits an array of mechanisms that singularly or collectively contribute to its multidrug resistance status. Intrinsic resistance includes inducible efflux pumps (2) and multiple β -lactamase expression (1) but not mutations in the quinolone resistance-determining region (9). In addition, *S. maltophilia* can acquire resistance through integrons, transposons, and plasmids (10). Recently, class 1 integrons have been characterized from *S. maltophilia* strains isolated in Argentina and Taiwan, which indicates that they contribute to TMP/SMX resistance through the *sul1* gene carried as part of the 3' end of the class 1 integron (10).

In addition to class 1 integrons, other mobile elements are associated with *sul* genes. For example, *Vibrio cholerae* serogroup O139 is resistant to several antimicrobial agents, including SMX, and it has been recently shown that the *sul2* gene was part of a cluster located on a novel genetic

*School of Medicine at Cardiff University, Cardiff, Wales, U.K.;

†School of Medical Sciences at University of Bristol, England, UK;

‡The JONES Group/JMI Laboratories, North Liberty, Iowa, USA

element of the integrative conjugative element group named SXT. The resistance genes harbored by SXT are embedded in a composite transposon-like structure and were probably acquired recently (11). Within this antimicrobial drug resistance region, an insertion element common region (ISCR) sequence, ISCR2, is adjacent to a *sul2* gene that moves by 1-ended transposition. Thus, the possibility exists that *sul2* genes can transfer intra- and intergenerally, including into *S. maltophilia*. Herein, we describe the molecular characterization of an international collection of *S. maltophilia* isolates and determine their mechanism of resistance to TMP/SMX, including the first report of *sul2* genes and the first description of insertion element common region (ISCR) elements carried in *S. maltophilia*.

Methods and Materials

Bacterial Strains

During 1998–2003, a total of 1,744 *S. maltophilia* isolates collected worldwide were forwarded to the SENTRY Program (Europe, USA, and Australia) and tested for antimicrobial drug susceptibility. A TMP/SMX resistance phenotype was demonstrated for 71. From these isolates, 25 nonclonal strains from patients in North America, Latin America, and Europe were analyzed by using molecular methods together with 30 representative isolates that were TMP/SMX-susceptible. Isolates were identified by using the Vitek System and confirmed by using API20NE (bioMérieux, Hazelwood, MO, USA).

Susceptibility Methods

Isolates were tested for susceptibility to TMP/SMX according to procedures of the Clinical and Laboratory Standards Institute (CLSI, formerly the National Committee for Clinical Laboratory Standards [NCCLS]) (12,13) by using broth microdilution methods (TREK Diagnostics, Cleveland, OH, USA). MIC results were confirmed with TMP/SMX. Etests were performed according to the manufacturer's directions (AB Biodisk, Solna, Sweden).

Molecular Materials

PCR primers were purchased from Sigma-Genosys Ltd. (Pampisford, UK) and are listed in the Table. General reagents for DNA manipulation were obtained from Invitrogen (Groningen, the Netherlands). All other reagents were obtained from Sigma Chemical Co. or BDH (both of Poole, England, UK).

Strain Typing

Clonality among the *S. maltophilia* isolates was assessed by pulsed-field gel electrophoresis (PFGE) fol-

lowed by *Xba*I digestion of genomic DNA. This assessment was conducted according to the standard 1-day protocol (16).

Plasmid Isolation

Bacterial plasmids were isolated by the alkaline lysis method described by Grinsted and Bennett (17). Essentially, an overnight 10-mL culture was centrifuged (12,000× *g*) and suspended in water (250 μL) before 200 μL of lysis solution (0.2 mol/L NaOH, 1% sodium dodecyl sulfate [SDS]) was added. After lysis, 125 μL of neutralizing solution (0.3 mol/L potassium acetate, 1 mmol/L EDTA) was added. After precipitation, the suspension was centrifuged (12,000× *g*) and washed twice with 500 μL of a 50/50 (v/v) phenol/chloroform solution. The DNA was precipitated from the solution with the addition of 0.7 volumes of iso-amyl alcohol. The DNA/RNA pellet was washed twice in 1 mL 70% ethanol before being dried. The DNA was dissolved in 30 μL with 0.1 U RNase.

Southern Hybridization

ISCR and *sul2* PCR amplicons generated with primers CRF/CRFF-r were labeled with P³²-CTP by random primer extension by using a commercially available kit (Stratagene, Amsterdam, the Netherlands) according to the manufacturer's instructions. Unincorporated nucleotides were removed by passing the labeled DNA through a Sephadex column (Nick column, Pharmacia Bio-tech, Uppsala, Sweden).

Agarose gels used for Southern transfer were denatured for 45 min in denaturing solution (0.5 mol/L NaOH, 1.5 mol/L NaCl) before being neutralized in 0.5 mol/L Tris-HCl, pH 7.5, 1.5 mol/L NaCl for 30 min. DNA was then transferred to Hybond (Amersham, Buckinghamshire, UK) nylon membrane by vacuum by using a custom-made Southern blotting apparatus. The nylon filter was prehybridized for at least 2 h with a blocking solution (6× SSC [1× SSC is 0.15 mol/L NaCl plus 0.014 mol/L sodium citrate], 0.1% [w/v] polyvinylpyrrolidone 400, 0.1% Ficoll [v/v], 0.1% bovine serum albumin, 0.5% SDS, 150 μg/mL denatured calf thymus DNA) at 65°C. The labeled denatured probe was then added to the solution and incubated overnight at 65°C. Finally, the filter was washed (300 mL 2× SSC, 0.1% [w/v] SDS followed by 0.1× SSC 0.1% SDS) at 65°C. Autoradiographic images were recorded on Hyperfilm-MP (Pharmacia Bio-tech), which was exposed overnight with intensifying screens.

PCR Analysis

The presence of class 1 integrons in each strain was assessed by using class 1 specific primers. Gene cassettes embedded within the class 1 integrons were determined by using primers listed in the Table. Isolates were also

screened for *sul1*, *sul2*, and *sul3* by using *sul1*-F and -R, *sul2*-F and -R, and *sul3*-F and -R, respectively. Seven positive class 1 integron PCR products were chosen randomly, extracted from agarose gels after size separation, and sequenced with IntF, IntR, and custom-made oligonucleotide primers (Table).

The presence of ISCR elements in each strain was also determined by using primers CRF/CRFF-r designed to amplify the same 700-bp fragment internal to the open reading frames (ORFs) of *ISCR1–5* (Table). Full-length *ISCR2* elements were amplified with primers designed to target the ends of *ISCR2*. Primers used to amplify genes often associated with *ISCR2* or *ISCR3* are also given (Table). Because *dhfr* genes are associated with ISCR elements, we also performed molecular analysis of them.

PCRs were conducted in a final volume of 20 μ L by using 10 μ L ABgene Expand Hi-fidelity Master Mix (ABgene House, Surrey, UK). Primers were used at final concentrations of 10 μ mol/L, and 1 μ L of an overnight bacterial culture (optical density 1.0 at 600 nm) was added as source of DNA template. The cycling parameters were as follows: 95°C for 5 min, followed by 30 cycles of 95°C

for 1 min, 55°C for 1 min, and 68°C for 1–4 min, depending on the sequence to be amplified, and ending with a 5-min incubation at 68°C.

DNA Sequencing and Analysis

Sequencing was conducted on both strands by the dideoxyl-chain termination method with a Perkin-Elmer Biosystems 377 DNA sequencer (Perkin-Elmer, Waltham, MA, USA). Sequence analysis was performed with the Lasergene DNASTAR software package (SelectScience Ltd., Bath, UK). Sequence alignments were conducted with the ClustalW program (www.ebi.ac.uk/clustalw) and the PAM 250 matrix.

The sequence of *ISCR2*, together with the adjacent *sul2* region and the novel *ISCR9* and *ISCR10*, has been deposited in GenBank. The genetic locus *ISCR2-glmM/sul2* from isolates 5232, 4647, 3800, and 2107 has been attributed the accession nos. AM182031, 182030, 182029, and 181666, respectively. *ISCR9* and *ISCR10* have been given the numbers AM182033 and AM182032, respectively.

Table. Oligonucleotide primers used in this study, Cardiff, 2007

Primer	Sequence (5'→3')	GenBank accession no.	Target	Reference
Ina-F	GCCTGTTTCGGTTCGTAAGCT		<i>intl</i>	(14)
Int-R	CGGATGTTGCGATTACTIONTCG		<i>intl</i>	(14)
<i>sul1</i> -F	ATGGTGACGGTGTTCGGCATTCTGA		<i>sul1</i>	(15)
<i>sul1</i> -R	CTAGGCATGATCTAACCCCTCGGTCT		<i>sul1</i>	(15)
<i>sul2</i> -F	GAATAAATCGCTCATCATTTTCGG	AJ289135	<i>sul2</i>	(15)
<i>sul2</i> -R	CGAATTCTTTCGGTTCCTTTCAGC	AJ289135	<i>sul2</i>	(15)
<i>aacA4</i> -F	AACCTGCGAGCGATCCGATG		<i>aacA4</i>	(14)
<i>aacA4</i> -R	ATGTACACGGCTGGACCATC		<i>aacA4</i>	(14)
<i>aacA7</i> -F	AATGGATAGTTTCGCCGCTCG		<i>aacA7</i>	This study
<i>aacA7</i> -R	TTCCGGAAGCAGCGCTACTTG		<i>aacA7</i>	This study
CRF	CACTWCCACATGCTGTTKCC	AF231986	All ISCR	This study
CRF-r	GMMACAGCATGTGGWAGTG	AF231986	All ISCR	This study
CRFF	GGRYGCAACGSCCTCAAGCG	AF231986	All ISCR	This study
CRFF-r	CGCTTGAGSCGTTGCRYCC	AF231986	All ISCR	This study
LECR2	CACTGGCTGGCAATGTCTAG	AF231986	<i>ISCR2</i>	This study
RECR2	CTTTGGACCGCAGTTGACTC	AF231986	<i>ISCR2</i>	This study
FloF	TGCACATCCTGGCTTCACTG	AF231986	<i>floR</i>	This study
FloR	ATTACAAGCGCGACAGTGGC	AF231986	<i>floR</i>	This study
<i>dfrA20f</i>	GGGAAACACCGAGAATGGG	AJ605332	<i>dfrA20</i>	This study
<i>dfrA20R</i>	TTCTTCTTCCCATTCTCCCC	AJ605332	<i>dfrA20</i>	This study
<i>dfrA9F</i>	CAGATTCGGTGGCATGAACC	X57730	<i>dfrA9</i>	This study
<i>dfrA9R</i>	GACCTCAGATACGAGTTTCC	X57730	<i>dfrA9</i>	This study
<i>dfrA10F</i>	TGTAGCGCGTGGTGTAAACG	AY055428	<i>dhfr10</i>	This study
<i>dfrA10R</i>	ACGTCTACGTGAGTATCCCG	AY055428	<i>dhfr10</i>	This study
<i>strA</i> F	TCTGTGCGACCTGCTTGATC	AY055428	<i>strA</i>	This study
<i>strA</i> R	CATTGCTGATGAACTGCGCG	AY055428	<i>strA</i>	This study
<i>tetA</i> F	CGCTGTTTGTGATTACACCC	AJ250203	<i>tetA</i>	This study
<i>tetA</i> R	CAGCGAGATGCGATATATCC	AJ250203	<i>tetA</i>	This study
<i>glmM</i> R	GAGTCAACTGCGGTCCAAC	AJ289135	<i>glmM</i>	This study
<i>glmM</i> F	ACGGTATTCTGGCAAAGCC	AJ289135	<i>glmM</i>	This study

Results

TMP/SMX MICs

TMP/SMX MICs separated the isolates into an obvious bimodal distribution. The TMP/SMX-resistant isolates possessed MICs >32 mg/L, whereas the sensitive controls used as molecular comparators possessed TMP/SMX MICs ranging from 0.5 to 2 mg/L (Online Appendix Table, available from <http://www.cdc.gov/EID/content/13/4/559-appT.htm>).

Detection and Determination of Class 1 Integrons

Of the 25 TMP/SMX-resistant *S. maltophilia* isolates that we analyzed, 17 possessed the *sul1* gene as part of the 3' end of a class 1 integron. None of the TMP/SMX-susceptible *S. maltophilia* isolates yielded positive *sul1* PCR products. PFGE analysis (data not shown) showed that only 2 isolates (9189 and 12221 from Chile) are clonally related (online Appendix Table). To our knowledge, this is the first report of *sul1*-positive *S. maltophilia* isolates from North America and Europe. The *sul1*-positive isolates are widespread, being from Europe, North America, and South America. Most (5) were isolated from Brazil. The integrons associated with the *sul1* gene vary in size; however, when 2 strains were isolated from the same country (e.g., 3438 and 3444, 9189 and 12221, and 98 and 14469), they possessed integrons of the same size, despite not being clonally related (Online Appendix Table). Seven of these integrons were randomly selected to examine their gene cassettes. The genetic context of the class 1 integrons and procured gene cassettes are shown in Figure 1. Strains 1893 (Germany) and 9431 (Brazil) possessed only the *int* and *sul/qac* genes. The class 1 integrons from strains 4891 (USA), 9189 (Chile), and 12221 (Chile) contained an embedded *aacA4* gene cassette. The 2 Mexican strains (3438 and 3444) contained 2 aminoglycoside-modifying genes (*aacA7* and *aadA5*) and an unknown ORF (Figure 1) yet were clonally unrelated, as judged by PFGE profiling. None of the integrons were the same as those characterized from strains isolated from Argentina (10).

Detection and Location of *sul2* Genes

All 55 isolates (both TMP/SMX resistant and sensitive) were screened for *sul2* genes with the primers listed in the Online Appendix Table. Nine of the isolates gave PCR products for *sul2*. None of the TMP/SMX-susceptible *S. maltophilia* isolates displayed positive *sul2* PCR products. Sequence analysis showed 100% identity with previous *sul2* sequences.

Given that *sul2* is normally located on medium-to-large sized plasmids, plasmids were isolated and characterized for *sul2* carriage. Plasmid DNA was prepared from each isolate and used as a template for PCRs by using the

sul2 primer detection set. In every case, a product of the size expected of *sul2* sequence amplification was obtained. The purity of each plasmid preparation was evaluated by attempted PCR amplification of the host cell chromosomal *gyrA* gene. In no case was an amplification product obtained when plasmid DNA was used as template; in contrast, a *gyrA* amplification product of the correct size was obtained from genomic DNA. These data were later confirmed by Southern hybridization that used the labeled *sul2* gene as a probe (data not shown). Unsurprisingly, in most cases *sul2* was found on a large plasmid of ~120 kb; however, in 2 of 9 *sul2*-positive isolates, *sul2* gene was chromosomally encoded.

Detection of ISCR Elements in TMP/SMX-sensitive and -resistant Strains

The *sul2* gene and *dhfr* genes are often found on plasmids and in close association with class 1 integrons or ISCR mobile genetic elements (10,15,18,19). Accordingly, we investigated the 55 *S. maltophilia* isolates for ISCR elements. Seven of the 25 TMP/SMX-resistant isolates yielded PCR products of the expected size (~700 bp) when the ISCR specific primers CRF/CRFF-r were used, and 6 of 23 TMP/SMX-sensitive *S. maltophilia* isolates also yielded the correct-sized amplification products.

To determine whether the locations of the ISCR sequences in the *S. maltophilia* isolates are chromosomal or plasmid mediated, plasmid DNA was prepared from each isolate and used as a template for ISCR-PCR and Southern hybridization analysis in a similar manner as described for *sul2*. In every case, a product of the size expected of ISCR sequence amplification was obtained. Hence, in those isolates that possess an ISCR element, the element is located on a plasmid (data not shown). The PCR




Figure 1. Schematic diagram of class 1 integrons from *Stenotrophomonas maltophilia* isolates. A) isolates 1893 and 9431; B) isolates 489, 9189, and 12221; C) isolates 3438 and 3444. Arrows depict direction of transcription, and shaded boxes represent gene cassettes found within the integron. The dark circles represent the 59-bp region immediately 5' to the incorporated gene cassette.

coli and *Salmonella enterica*. A recent study by Antunes et al. found *sul1*, *sul2*, or *sul3* genes in most Portuguese isolates (18); 24 of 200 isolates contained both *sul1* and *sul2*. *sul2* has also recently been identified in *S. enterica* from Brazil (20). Similar results have been reported from *E. coli* urinary tract isolates in which $\approx 26\%$ of strains possessed both *sul1* and *sul2* genes (21). A biased study examining TMP/SMX-resistant *E. coli* recently reported that 15 of 20 isolates possessed *sul2* and that 6 of those also carried *sul1* on a class 1 integron (14). Additional studies of *E. coli* have shown the intercontinental predominance of *sul1* through class 1 integrons (22). A study by Pei et al. demonstrated the correlation of anthropogenic activity with the presence of *sul* genes in environmental samples (23). However, none of the studies demonstrated the genetic origin of the *sul2*.

In addition to *sul* genes associated with plasmids and class 1 integrons, we investigated whether the *S. maltophilia* isolates possessed ISCR elements and whether these could be linked to *dhfr* or *sul* genes, as has been shown (18). Of the 25 TMP/SMX-resistant isolates, 6 harbored *sul2* linked to ISCR2. However, we could not detect any *sul3* genes. In the isolates with ISCR2, the element was directly linked to a deleted version of a phosphoglucomutase gene, $\Delta glmM$, as has been reported on other occasions (Figure 2). This arrangement is identical to those of 5 other sequences in the EMBL database, in *E. coli* isolated from cattle in France and Germany (24), in the plasmid pRVS1 isolated from a strain of *Vibrio salmonicida* from Norway, in a plasmid from a strain of *S. enterica* isolated in Japan, and on the chromosome of *Shigella flexneri* isolated in the United States (18,24). In all cases, $\Delta glmM$ and *sul2* are linked to the end of ISCR2 that accommodates the IS91 *oriIS* equivalent (Figure 2). The dual arrangement of $\Delta glmM$ and *sul2* is also found in plasmids of marine psychrotrophic bacteria isolated in Norway (GenBank accession no. AJ306553/4), but in these cases the ISCR2 element appears not to be present.

Two of the isolates harbored a copy of the *floR* gene immediately upstream of a copy of ISCR2 (Figure 2), an arrangement identical to that reported on plasmids found in isolates of *E. coli* from cattle in France and Germany (24). The *S. maltophilia* isolates investigated in this study came from Turkey and the United States. Two isolates from Spain also carry the *floR* gene but not ISCR2. Instead, the isolates possess copies of ISCR3, which do not appear to be linked to *floR*. The finding of florfenicol-resistant traits on plasmids in different bacterial species from different countries highlights the wide geographic spread of this resistance mechanism. The location of *floR* next to ISCR2 is such that it is possible, if not probable, that the resistance gene can be cotransposed with the ISCR element.

The findings within this study are important for several reasons. First, this is, to our knowledge, the first report of ISCR elements being found in *S. maltophilia* isolates. In 6 cases, these were linked to *sul2* genes responsible for the TMP/SMX-resistant phenotype. Moreover, these isolates were unrelated strains found in different countries. Second, since TMP/SMX is the mainstay therapy for *S. maltophilia* infections, the mobilization of *sul* genes by means of class 1 integrons and ISCR elements is likely to increase with TMP/SMX consumption. Third, most *sul2* genes in this study have been found on plasmids, and *sul2*-containing plasmids can potentially confer an increase in bacterial "fitness" (25). As yet, such phenomena have only been explored in *Enterobacteriaceae*, and it has yet to be established whether *sul2*-carrying plasmids have such an additive effect in *S. maltophilia* or for that matter, other nonfermenting gram-negative bacilli.

These data suggest that microbiology laboratories need to carefully monitor *S. maltophilia* TMP/SMX resistance, which has the potential to increase by means of mobile elements. We also advocate the continued international surveillance of antimicrobial drug resistance that may act as early warning systems for this kind of resistance. Furthermore, yearly monitoring with molecular probes is advisable.

This work is funded by the EC through COBRA contract LSHM-CT-2003-503335 and a STREP DRESP2 contract. M.A.T. is currently funded by Cardiff University.

Dr Toleman is currently working as a research fellow at the Medical School, Cardiff University, Wales. His interest is the dissemination of antimicrobial-drug-resistant genes among clinical and environmental bacteria.

References

1. Avison MB, Higgins CS, Ford PJ, von Heldreich CJ, Walsh TR, Bennett PM. Differential regulation of L1 and L2 β -lactamase expression in *Stenotrophomonas maltophilia*. *J Antimicrob Chemother.* 2002;49:387–9.
2. Li X-Z, Nikaido H. Efflux-mediated drug resistance in bacteria. *Drugs.* 2004;64:159–204.
3. Sader HS, Jones RN. Antimicrobial susceptibility of uncommonly isolated non-enteric gram-negative bacilli. *Int J Antimicrob Agents.* 2005;25:95–109.
4. Hogardt M, Schmoltd S, Götzfried M, Adler K, Heesemann J. Pitfalls of polymyxin antimicrobial susceptibility testing of *Pseudomonas aeruginosa* isolated from cystic fibrosis. *J Antimicrob Chemother.* 2004;54:1057–61.
5. Fedler KA, Biedenbach DJ, Jones RN. Assessment of pathogen frequency and resistance patterns among paediatric patient isolates: report from the 2004 SENTRY Antimicrobial Surveillance Program on three continents. *Diagn Microbiol Infect Dis.* 2006;56:427–36.

6. Gales AC, Jones RN, Forward KR, Linares J, Sader HS, Verhoef J. Emerging importance of *Acinetobacter* species and *Stenotrophomonas maltophilia* as pathogens in severely ill patients: geographic patterns, epidemiological features, and trends in the SENTRY Antimicrobial Surveillance program (1997–1999). *Clin Infect Dis*. 2001;32(Suppl):S104–13.
7. Al-Jasser AM. *Stenotrophomonas maltophilia* resistant to trimethoprim sulfamethoxazole: an increasing problem. *Ann Clin Microbiol Antimicrob*. 2006;5:23–6.
8. Zelenitsky SA, Iacovides H, Ariano RE, Harding GK. Antibiotic combinations significantly more active than monotherapy in an in vitro infection model of *Stenotrophomonas maltophilia*. *Diagn Microbiol Infect Dis*. 2005;51:39–43.
9. Valdezate S, Vindel A, Saéz-Nieto JA, Baquero F, Canton R. Preservation of topoisomerase genetic sequences during in vivo and in vitro development of high-level resistance to ciprofloxacin in isogenic *Stenotrophomonas maltophilia* strains. *J Antimicrob Chemother*. 2005;56:220–3.
10. Barbolla R, Catalano M, Orman BE, Famiglietti A, Vay C, Smayevsky J, et al. Class 1 integrons increase trimethoprim/sulfamethoxazole MICs against epidemiologically unrelated *Stenotrophomonas maltophilia* isolates. *Antimicrob Agents Chemother*. 2004;48:666–9.
11. Beaber JW, Hochhut B, Waldor MK. Genomic and functional analyses of SXT, an integrating antibiotic resistance gene transfer element derived from *Vibrio cholerae*. *J Bacteriol*. 2002;184:4259–69.
12. National Committee for Clinical Laboratory Standards. Methods for dilution antimicrobial tests for bacteria that grow aerobically: approved standard M7–A6. Wayne (PA): The Committee; 2003.
13. Clinical Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing. Supplemental tables M100–S15. Wayne (PA): The Institute; 2005.
14. Infante B, Grape M, Larsson M, Kristiansson C, Pallecchi L, Rossolini GM, et al. Acquired sulphonamide resistance genes in faecal *Escherichia coli* from healthy children in Bolivia and Peru. *Int J Antimicrob Agents*. 2005;25:308–12.
15. Toleman MA, Bennett PM, Walsh TR. Common regions e.g. orf513 and antibiotic resistance: IS91-like elements evolving complex class 1 integrons. *J Antimicrob Chemother*. 2006;58:1–6.
16. Centers for Disease Control and Prevention. One-day (24–48 h) standardization laboratory protocol for molecular sub-typing of *Escherichia coli* O157:H7, non-typhoidal *Salmonella* serotypes, and *Shigella sonnei* by pulsed-field gel electrophoresis (PFGE). In: PulseNet PFGE manual. Atlanta: The Centers; 2002.
17. Grinsted J, Bennett PM. Preparation and electrophoresis of plasmid DNA. In: Grinsted J, Bennett PM, editors. *Plasmid technology*. London: Academic Press; 1990. pp. 129–42.
18. Antunes P, Machado J, Sousa JC, Piexe L. Dissemination of sulfonamide resistance gene (*sul1*, *sul2* and *sul3*) in Portuguese *Salmonella enterica* strains and relation with integrons. *Antimicrob Agents Chemother*. 2005;49:836–9.
19. Toleman MA, Bennett PM, Walsh TR. Common regions: novel gene capturing systems of the 21st century? *Microbiol Mol Biol Rev*. 2006;70:296–316.
20. Michael GB, Cardoso M, Schwartz S. Class 1 integron-associated gene cassettes in *Salmonella enterica* subsp. *enterica* serovar *Agona* isolated from pig carcasses in Brazil. *J Antimicrob Chemother*. 2005;55:776–9.
21. Grape M, Farra A, Kronvall G, Sundstrom L. Integrons and gene cassettes in clinical isolates of cotrimoxazole-resistant gram-negative bacteria. *Clin Microbiol Infect*. 2005;11:185–92.
22. Blahna MT, Zalewski CA, Reuer J, Kahlmeter G, Foxman B, Marrs CF. The role of horizontal gene transfer in the spread of trimethoprim-sulfamethoxazole resistance among uropathogenic *Escherichia coli* in Europe and Canada. *J Antimicrob Chemother*. 2006;57:666–72.
23. Pei R, Kim S-C, Carlson KH, Pruden A. Effect of river landscape on the sediment concentrations of antibiotics and corresponding antibiotic resistance genes (ARG). *Water Res*. 2006;40:2427–35.
24. Cloeckaert A, Baucheron S, Flaujac G, Schwarz S, Kehrenberg C, Martel JL, et al. Plasmid-mediated florfenicol resistance encoded by the *floR* gene in *Escherichia coli* isolated from cattle. *Antimicrob Agents Chemother*. 2000;44:2858–60.
25. Enne VI, Bennett PM, Livermore DM, Hall LM. Enhancement of host fitness by the *sul2*-coding plasmid p9123 in the absence of selective pressure. *J Antimicrob Chemother*. 2004;53:958–63.

Address for correspondence: Timothy R. Walsh, Department of Medical Microbiology, School of Medicine, Cardiff University, Heath Park, Cardiff CF14 4XN, Wales, U.K.; email: WalshTR@Cardiff.ac.uk

Use of trade names is for identification only and does not imply endorsement by the Public Health Service or by the U.S. Department of Health and Human Services.

EMERGING INFECTIOUS DISEASES *online*

www.cdc.gov/eid

To receive tables of contents of new issues send an email to listserv@cdc.gov with `subscribe eid-toc` in the body of your message.