

---

# Cost-effectiveness of Antiviral Stockpiling and Near-Patient Testing for Potential Influenza Pandemic

M. Ruby Siddiqui\* and W. John Edmunds\*

A decision analytical model was developed to investigate the cost-effectiveness of stockpiling antiviral (AV) drugs for a potential influenza pandemic in the United Kingdom and the possible role of near-patient testing in conserving AV drug stocks. Under base-case assumptions (including a fixed stockpile that was smaller than the clinical attack rate), the treat-only option (treating all symptomatic patients with AV drugs) would be considered cost-effective (£1,900–£13,700 per quality-adjusted life year [QALY] gained, depending on the fatality scenario), compared with no intervention (nonintervention but management of cases as they arise). The test-treat option (testing all symptomatic patients but treating those with positive tests results only) would result in moderate gains in QALYs over the treat-only option but at relatively large additional costs. Stockpiling sufficient AV drugs (but not near-patient tests) to treat all patients with clinical cases would be cost-effective, provided AV drugs are effective at preventing deaths from pandemic influenza.

Many countries are ordering stockpiles of antiviral (AV) drugs for use in a potential influenza pandemic. The United Kingdom recently announced the procurement of 14.6 million courses of oseltamivir, enough for almost 25% of the population (1).

The timing of an influenza pandemic cannot be predicted (the most recent pandemics occurred in 1918, 1957, and 1968/69). Because AV drugs have a limited shelf-life, long-term maintenance of stockpiles may constitute a significant cost. Similarly, the size (clinical attack rate [CAR]) of a pandemic cannot be accurately foreseen. Therefore, triaging of patients with influenza-like illness (ILI) may be essential to conserve limited AV drug stocks. Possible triaging methods include near-patient testing (rapid diag-

nostic tests at the point of care). This study assessed the cost-effectiveness of stockpiling AV drugs for a potential influenza pandemic and, in the event of a pandemic, also assessed the use of near-patient testing in the management of AV drugs.

## Methods

A decision analytical model (Figure 1) was constructed to compare the costs and quality-adjusted life year (QALY) loss associated with 3 potential strategies for the management of patients with ILI in the United Kingdom: 1) do not treat with AV drugs and manage complications if they arise (no intervention), 2) treat all patients with AV drugs (treat only), or 3) test then treat those who test positive for influenza with AV drugs (test-treat). Precision Tree (Palisade Corporation, Ithaca, NY, USA) running in Microsoft (Redmond, WA, USA) Excel was used to construct the model and @Risk (Palisade Corporation) was used to perform the probabilistic sensitivity analysis.

## Epidemiologic Scenarios

Baseline epidemiologic scenarios were based on the UK Department of Health Pandemic Contingency Plan (2), which assumes a cumulative CAR of 25% over 1 wave lasting 15 weeks. The demand on general practitioners (GPs) and accident and emergency departments (A&Es) would require an alternative means for AV drug distribution (2). Patients with ILI were therefore assumed to have received AV drug therapy by visiting teams or call-in centers. Of these, because of secondary complications, 5% were assumed to have consulted further with a GP and 5% with A&E departments. Also, among ILI patients, 0.55% were expected to be hospitalized. All model parameters are summarized in the online Appendix Table (available from [www.cdc.gov/EID/content/14/2/267-appT.htm](http://www.cdc.gov/EID/content/14/2/267-appT.htm)).

---

\*Health Protection Agency, London, UK




Figure 1. Decision analytical model tree of treatment strategies for patients with an influenza-like illness (ILI) during an influenza pandemic. All branches culminate in the subtree (indicated with +). QALY, quality-adjusted life year; CAR, clinical attack rate.

Background ILI rates were deduced from the mean ILI consultation rates in England and Wales observed by GP-based sentinel surveillance from 1985–2003 (4) and the estimated proportion (28%) of clinical ILI case-patients who consulted with GPs in interpandemic periods (5). Epidemic influenza is seasonal with a higher incidence in winter (wk 40–12) than summer (wk 13–39). The mean weekly ILI incidence was assumed to have remained unchanged during the pandemic, resulting in a cumulative incidence of 3.1% (winter) or 1% (summer). An additional value of 4% for midwinter (wk 49–8) was also considered in the sensitivity analyses. It is possible that a pandemic strain would out-compete the existing epidemic influenza strains and the background ILI rate would fall. However, other pathogens (e.g., respiratory syncytial virus and rhinovirus) that contribute to ILI would remain unaffected (22). The background ILI rate was therefore assumed to remain unaltered. The base-case analysis assumed a winter pandemic, giving a probability (PrF) of 89% ( $CAR/(CAR + \text{background ILI rate})$ ) that a case of ILI was pandemic influenza.

Two fatality scenarios were assumed, on the basis of the 1918 influenza pandemic (1918 scenario), which had an overall case-fatality ratio (CFR) of 2.3%, and the 1957 and 1968/69 influenza pandemics (1957/69 scenario) which had an average CFR of 0.3%, with most deaths occurring in the elderly (online Appendix Table).

Three pandemics occurred in the 20th century, which suggests that a pandemic occurs approximately every 30 years but that this is a random (Poisson) process. For the base-case analysis, we assumed that the next pandemic would take place in 30 years; this figure was varied from 1 to 50 years in the sensitivity analyses.

### Antiviral Treatments

The neuraminidase inhibitor oseltamivir was selected as the AV treatment of choice because it is cheaper and easier to stockpile than zanamivir and has been shown to dominate zanamivir in cost-utility analyses (23). Cyclic amines (amantadine and rimantadine) were not considered because resistance to these drugs has emerged in influenza A virus (H5N1) with pandemic potential (24). The rate of adverse events after the use of neuraminidase inhibitors is low and therefore was not considered in this analysis (12).

Meta-analyses of oseltamivir efficacy studies were compared, and the 2003 study by Kaiser et al. was selected for the parameterization of this model because it included the greatest number of subjects (5,15,20). No data were available for AV efficacy in reducing influenza-related deaths so the same efficacy as that for reducing influenza-related hospitalizations was assumed. Assuming all GP and accident and emergency (A&E) consultations were attributable to the development of complications (2), the probability of complications and the probability of hospitalizations (given no AV therapy) were calculated by using the odds ratios from the oseltamivir meta-analyses (15). The probability of complications for noninfluenza ILI patients was considered the same as for untreated influenza patients, but the probability of hospitalization for noninfluenza ILI patients was considered to be 40% that of untreated influenza patients (15).

We assumed that for both the treat-only and test-treat strategies all qualifying patients were given AV drugs and were tested until the stockpile ran out. No epidemiologic differences were assumed between those who received AV drugs and those who did not because the stockpile was depleted. Neuraminidase inhibitors are only recommended for use within 48 hours of symptom onset (25). We assumed that the efficiency of AV drug distribution would be such that 70% of ILI patients would receive timely AV drugs; this varied from 30% to 97% in our sensitivity analyses (online Appendix Table). Those receiving AV drugs after 48 hours were assumed to derive no benefit.

The shelf-life of oseltamivir is currently 5 years. However, tests are being conducted to increase this to 6 years (6). Shelf-life was varied in our sensitivity analyses between 4 and 6 years.

For base-case analyses, we assumed the AV drug stockpile to be 14.6 million courses (1), giving an 87% probability of case-patients receiving AV drugs (size of

Table. Total NHS costs and QALY loss (discounted at 3.5%) resulting from an influenza pandemic occurring in 30 years assuming 1918 or 1957/69 CFR\*

Treatment program	Pandemic influenza cases (millions)	Pandemic influenza deaths (millions)	Discounted NHS costs (million £)	Discounted QALY loss (millions)	Incremental cost per QALY (£)
1918 scenario					
No intervention	15	0.344	113	2.23	
Treat only	15	0.236	1,361	1.56	1,861†
Test-treat	15	0.231	2,356	1.53	31,031‡
1957–69 scenario					
No intervention	15	0.044	113	0.395	
Treat only	15	0.030	1,361	0.303	13,668†
Test-treat	15	0.030	2,356	0.299	227,896‡

\*NHS, National Health Service; QALY, quality-adjusted life year; CFR, case-fatality ratio.

†Cost per QALY gained over no intervention program.

‡Cost per QALY gained over treat-only program.

stockpile/total ILI cases). Hence, the size of the AV stockpile would be limited under the treat-only strategy.

### Near-Patient Tests

A number of near-patient influenza tests are currently available (7–10). To reflect the continual improvement in test technology, a composite test was constructed based on the best performance of currently available tests (89.5% sensitivity and 99.8% specificity) (Directigen Flu A+B, Becton Dickinson, Sparks, MD, USA) (7,12), a shelf-life of 2 years (Quickvue, Quidel Corporation, San Diego, CA, USA) (9), and a cost of £7 per test (Biostar Flu OIA, Inverness Medical-Biostar Inc, Louisville, CO, USA) (8). Parameter distributions for test sensitivity and specificity were deduced from a meta-analysis of near-patient tests (12), but distributions were assumed for the shelf-life and cost.

Although the United Kingdom has not accumulated a stockpile of near-patient tests, we assumed it to be the same size as the AV drug stockpile in the base-case analysis. The probability of being tested (test-treat option) was 87% (size of stockpile/total ILI cases).

### Costs

As recommended in the United Kingdom, costs were analyzed from the perspective of the healthcare provider, the National Health Service (NHS). Future costs and benefits were discounted at the current rate of 3.5% per annum and all costs were in 2004 pounds sterling (£1 = ≈US\$1.8) (14,26).

Mean unit costs per GP consultation, hospitalization, and A&E attendance for ILI were deduced from standard sources (17,19). We assumed that no additional costs were associated with death. The costs of complications leading to GP consultation or hospitalization were assumed to be the same for pandemic influenza and nonpandemic ILI.

The unit cost of an AV treatment course was assumed to be £16 (16) for the treat-only option and £16.87 (because of excess AV drugs) for the test-treat option (fixed stock-

piles). The unit cost of a near-patient test was assumed to be £7 (7), and storage was £1 per unit (AV drug course or test) per year for both programs. Administration costs for the distribution of AV drugs or near-patient testing by visiting teams or call-in centers were assumed equivalent to the mean cost of a home visit by a district nurse, health visitor, health care assistant, or practice nurse (£15.75 per test or AV drug course) (17). Units were assumed to be procured >2 years (1) and replenished at expiration of shelf-life.

### Health Benefits

Health benefits were assessed by using QALYs. QALY loss associated with uncomplicated ILI were calculated study by O'Brien et al. (20) assuming a normal health score of 0.85 (online Appendix Table). QALY loss associated with the development of complications was assumed to be the sum of QALY loss associated with uncomplicated ILI and that associated with pneumococcal pneumonia (outpatient) because this was the most likely complication (21). Similarly, QALY loss associated with hospitalization was considered the sum of QALY loss associated with uncomplicated ILI and that associated with pneumococcal pneumonia (inpatient) (21).

The mean discounted QALY loss associated with pandemic influenza death was estimated by using age-specific CFR under the 2 death scenarios and background life expectancy by age weighed by age-adjusted quality-of-life scores (online Appendix Table) (27). QALY loss associated with noninfluenza death was assumed to be the same as that associated with *S. pneumoniae* death (a mean discounted QALY loss per noninfluenza death of 6.09 from 1980–2000) (3).

### Base-Case Assumptions

In the base-case analysis, the cost-effectiveness of the potential strategies was compared under the assumptions of fixed AV and test stockpiles (14.6 million units), a CAR of 25%, and a time to pandemic of 30 years. There is no threshold for cost-effectiveness in the United Kingdom, although the National Institute for Clinical Excellence will probably

reject an intervention on cost-effectiveness grounds if the cost per QALY gained is in excess of £25,000–£35,000 (28). For ease of exposition, we used a simple threshold of £30,000 per QALY to define a cost-effective intervention.

## Results

### Base-Case Analysis (Fixed Stockpiles)

In our base-case model, we estimated that in a pandemic, ILI would develop in 28.1% of the population (16.8 million persons) (including 15 million pandemic influenza patients). Under the high CFR conditions of the 1918 scenario, ≈344,000 deaths would occur compared with ≈44,000 deaths under the 1957/69 scenario (Table). These scenarios would result in the loss of ≈2.2 or 0.4 million discounted QALYs, respectively, with a total discounted cost to the NHS of £113 million if no treatment program were initiated (no intervention). The treat-only program would reduce this loss by 700,000 or 90,000 QALYs at a cost of ≈£1,900 or £13,700 per QALY gained for the 1918 and 1957/69 scenarios, respectively, well below the £30,000 threshold. The test-treat program would further reduce this loss slightly by 30,000 or 4,000 QALYs but at a high cost of ≈£31,000 or £228,000 per QALY gained over the treat-only alternative. The test-treat option would be unlikely to be considered because cost-effectiveness is highly dependent on the fatality scenario.

### Univariate Sensitivity Analysis of the Treat-Only Program (Fixed Stockpile)

Because the treat-only program was the most cost-effective program under both fatality scenarios, we carried out a univariate sensitivity analysis of the incremental cost-effectiveness of this program to variability in model parameters (online Appendix Table). AV drug efficacy for reducing complications and hospitalizations had minimal effect on the cost-effectiveness of the treat-only program, but this strategy was highly sensitive to AV drug efficacy for reducing death (Figure 2). This was due to the relatively high QALY loss associated with pandemic influenza death (94% and 69% of the total QALY loss for the 1918 and 1957/69 scenarios, respectively). Because the value of this parameter is unclear, further studies of the potential protective effect of AV drugs against death are essential.

The timing of the pandemic and the discount rate were influential parameters. However, variation in the timing of the epidemic is unlikely to change the recommendation that the treat-only strategy is cost-effective. If an epidemic occurs in 45 years, the costs per QALY gained would be ≈£3,800 and £28,000 for the 2 fatality scenarios (discount rate of 3.5%), still below the £30,000 threshold. At a discount rate of 6%, the treat-only option would be cost-effective for up to 30 years.




Figure 2. Univariate sensitivity analyses of the incremental cost-effectiveness of the treat only strategy over the no-intervention strategy to model parameters under the 1918 scenario (A) and 1957/69 scenario (B). OR, odds ratio; AV, antiviral; CAR, clinical attack rate; ILI, influenza-like illness; A&E, accident and emergency department; GP, general practitioner; QALY, quality-adjusted life year.

The efficiency of AV drug distribution is likely to be important. The program would however remain cost-effective if the probability of receiving AVs within 48 hours did not drop below 35% (£3,700 or £27,000 per QALY gained for the 1918 and 1957/69 scenarios, respectively).

The treat-only program was slightly more cost-effective in the summer than the winter or midwinter as the probability of pandemic influenza being the cause of ILI was higher (96%, 89%, and 84%, respectively). Wastage (wasted quantities, fraud, theft) of AV drug supplies as high as 25% had little effect on cost-effectiveness of the treat-only option (Figure 2).

A lower CAR (15%) reduced the cost-effectiveness of AV drugs because some of the stockpile would not be used (surplus). Increasing the CAR above 25% had no effect on programs with fixed stockpiles because the same number of deaths and complications would be prevented at the same cost (the proportion of deaths prevented would be reduced, but the absolute number would remain the same).

### Threshold Conditions for Test-Treat Option (Fixed Stockpiles)

For high CARs, where a fixed AV drug stockpile is less than the expected demand (as in the base-case), near-pa-



tient tests could be used to better target therapeutic courses. A univariate sensitivity analysis of the incremental cost-effectiveness of test-treat over treat only to variability in the near-patient test parameters, test sensitivity, specificity, unit cost, and shelf-life, was carried out. Under the 1918 scenario the test-treat strategy would require test sensitivity to exceed  $\approx 90\%$  (Figure 3, panel A) or a test unit cost below  $\pounds 6$  or a shelf-life above 3 years (Figure 3, panel B) to be considered cost-effective. Test specificity would have little effect on the incremental cost-effectiveness because it has no effect on QALY loss. Under the 1957/69 scenario test-treat would never cross the cost-effectiveness threshold even with a 100%-sensitive or 100%-specific test, a test cost as low as  $\pounds 0$ , or a shelf-life as high as 4 years.

### Probabilistic Sensitivity Analysis (Fixed Stockpiles)

Model parameters were varied (online Appendix Table) in a probabilistic sensitivity analysis, which suggests that for fixed AV drug and test stockpiles, the probability is high that the treat-only option would be cost-effective, irrespective of the fatality scenario (Figure 4). The test-treat option would result in small QALY gains (and often losses) but at substantial additional costs. The probability of this strategy being cost-effective is low compared with the treat-only option, particularly for the 1957/69 fatality scenario.

### Incremental Cost-effectiveness during a Pandemic Wave (Fixed Stockpiles)

The probability that an ILI case will be due to pandemic influenza will vary over the time course of a pandemic (assumed to peak between wk 6 and 7 for a wave lasting 15 wk) (2). Therefore, near-patient testing may be useful during early stages of a pandemic when clinical judgment is low and inappropriate AV administration is high. The cost-effectiveness of test-treat over a pandemic wave was analyzed.

The AV drug stockpile was assumed to remain fixed at 14.6 million courses (1), and the test stockpile was varied with the cumulative number of ILI cases expected per week of the pandemic wave. Figure 5 shows the total incremental cost-effectiveness of the test-treat strategy over treat only for each test stockpile for a CAR of 25%. Test-treat would be cost-effective ( $< \pounds 30,000$  per QALY gained) for test stockpiles up to 12.1 million (the expected no. of cumulative ILI cases at wk 8 of a pandemic) under the 1918 scenario. Test-treat may even be considered for test stockpiles up to 13.7 million (wk 9 of a pandemic) as the cost-effectiveness was  $\approx \pounds 32,700$  per QALY gained. However, under the 1957/69 scenario test-treat would not be cost-effective at any stage of the pandemic, although it may be considered for test stockpiles up to  $\approx 35,000$  (wk 2 of a pandemic) as the cost was  $\approx \pounds 34,000$  per QALY gained.

For a CAR of 15% (data not shown), test-treat would not be cost-effective throughout a pandemic as the AV drug




Figure 3. Univariate sensitivity analyses of the incremental cost-effectiveness of the test-treat strategy over the treat only strategy to A) near-test sensitivity and specificity and B) near-test unit cost and shelf-life. The test-treat program becomes cost-effective below the cost-effectiveness threshold ( $\pounds 30,000$  per quality-adjusted life year [QALY] gained).

stockpile would exceed demand (cumulative ILI cases). For a CAR of 35% (data not shown), although test-treat would be cost-effective for test stockpiles up to 16.4 million (wk 8) under the 1918 scenario, it would not be cost-effective at any stage of the pandemic under the 1957/69 scenario. Therefore in the short-term, stockpiling enough tests for the first 2 weeks of a potential influenza pandemic ( $\approx 35,000$  tests) could help conserve limited AV drug stockpiles. However, this is highly dependent on the CAR and CFR.

### Optimal Stockpiling

In the long term, it may be more cost-effective to increase stockpiles to cover expected demand (CAR +




Figure 4. Probabilistic sensitivity analysis of the incremental cost-effectiveness of the treat-only over the no-intervention strategy and the test-treat strategy over the treat-only strategy for the A) 1918 and B) 1957/69 death scenarios (1,000 iterations). Cost-effective strategies lie to the right of the cost-effectiveness threshold (£30,000 per quality-adjusted life year [QALY] gained).

background ILI + wastage). Figure 6 and the online Appendix Figure (available from [www.cdc.gov/EID/content/14/2/267-appG.htm](http://www.cdc.gov/EID/content/14/2/267-appG.htm)) show the expected costs and QALY losses under a range of different AV drug and test stockpiles and CARs (base-case test characteristics and AV drug efficacy assumed). Each point represents 1 scenario (test and AV drug stockpile size). Points on the efficiency line were potentially cost-effective strategies. Strategies that increase cost but reduce QALY loss (moving from left to right on the efficiency line) should be considered until the slope of the line exceeds the threshold of £30,000 per QALY (efficiency line ends). For each CAR, this process suggested that the optimal strategy was treat only, stockpiling enough AV drugs to meet demand (CAR plus background ILI plus AV drug wastage). Therefore, for a CAR of 25%, the optimum stockpile was  $\approx 20$  million AV drugs only (Figure 6) because the expected number of ILI cases would be 16.8 million (15 million of which would be pandemic influenza) and the expected AV drug wastage would be 2.2 million. The test-treat strategies were never on the efficiency line, even for a perfect test (100% sensitivity and specificity), because they resulted in similar QALY loss as treat only but at increased costs. Indeed, when the size of the AV drug stockpile exceeded the demand, test-treat resulted in

increased QALY loss (if the test is not 100% sensitive), because some true pandemic influenza case-patients would be denied treatment even though there was a surplus of AV drug courses.

## Discussion

This study demonstrates that stockpiling AV drugs for a treat-only program is likely to be a cost-effective strategy in preparation for a potential influenza pandemic, even if the pandemic occurs many years from now, assuming that AV drugs provide some protection against death. However, under current UK planning assumptions (CAR 25%), the AV stockpile would be too small (at 14.6 million courses) to treat all cases of ILI. Near-patient testing is unlikely to be a cost-effective approach to conserving AV stocks but might be considered early in a pandemic. A more cost-effective strategy would be to increase the stockpile of AV drugs. Since CARs in excess of 30% have been observed in pandemics (29), increasing the stockpile to cover this possibility may be both prudent and cost-effective. Indeed, expanding the stockpile of AV drugs to encompass the whole UK population ( $\approx 60$  million) might even be acceptable ( $\approx £6,500$  per QALY gained over a no intervention strategy for the 1918 scenario under base-case assumptions).

Stockpiling AVs is a cost-effective option, even though some benefits have been ignored (e.g., possible reduction in CAR if widespread and prompt treatment is offered) (30). Furthermore, additional benefits could be derived from using AV drug stockpiles close to their expiration dates to treat epidemic influenza patients. Finally, the reduction in illness and death that may result from widespread AV drug use is likely to bring benefits to other sectors of the economy.

This study focused on mass treatment strategies because this is the policy of the United Kingdom and other




Figure 5. Incremental cost-effectiveness of the test-treat strategy over the treat-only strategy during a pandemic wave (antiviral [AV] stockpile = 14.6 million courses, test stockpile = number of cumulative influenza-like [ILI] cases, clinical attack rate = 25%). QALY, quality-adjusted life year.




Figure 6. Optimal cost-effectiveness of antiviral (AV) and test stockpiling (0–30 million units) for a clinical attack rate (CAR) of 25% under the A) 1918 and B) 1957–69 scenarios. The composite test (Test A) and a perfect test of 100% sensitivity and 100% specificity (Test B) are included. The most cost-effective strategies lie on the efficiency line. ILI, influenza-like illness; QALY, quality-adjusted life year.

countries (31). However, strategies targeting those at higher risk of complications or death would be more cost-effective provided the delivery costs are similar and AV drugs are as effective in these groups.

The main caveat to an AV drug stockpiling strategy is the uncertainty concerning the efficacy of AV drugs against the next strain of pandemic influenza, particularly efficacy against influenza-related deaths. Clearly, estimation of this important parameter should be a priority if governments are to commit large resources to mitigating the effect of an uncertain pandemic occurring at an unknown point in the future.

Use of AV drugs on the scale anticipated may create a selective pressure (any factor that leads to preferential survival of organisms with specific traits) for the emergence of AV drug resistance. In this case, large-scale use of AVs could lead to the preferential survival of flu viruses

that are resistant to AVs. However, studies to date indicate that resistance to oseltamivir in influenza virus strains occurs rarely (32,33) and that such mutations may have a fitness cost in terms of impaired growth and transmissibility (34,35). Nevertheless, recent reports suggest reduced susceptibility to oseltamivir in some currently circulating strains of avian influenza (H5N1) with pandemic potential (36–38). Clearly this may reduce the effectiveness of an oseltamivir stockpiling program, although AV drugs might still delay pandemic spread to allow vaccine development (39). Further epidemiologic and modeling studies of the potential effect of oseltamivir resistance on viral fitness and drug effectiveness are required.

### Acknowledgments

We thank Nigel Gay for helpful discussions.

This work was funded by the Health Protection Agency, United Kingdom, and by a European Commission project funded within the Sixth Framework Program, Thematic Priority Scientific Support to Policies (Contract no. SP22-CT-2004-003824: SARS Control: Effective and acceptable strategies for the control of SARS in China and Europe).

Dr Siddiqui is an infectious disease modeler in the Modelling and Economics Unit of the Health Protection Agency Centre for Infections, London. Her research interests include designing and adapting models to study transmission and burden of infectious diseases.

Dr Edmunds is the head of the Modelling and Economics Unit at the Health Protection Agency Centre for Infections in London. His work concentrates on the design of cost-effective intervention programs against infectious diseases, taking into account the direct and indirect effects of such programs.

### References

1. United Kingdom Department of Health. Improving preparedness for possible flu pandemic. 2005 Jan 3 [cited 2005 Feb 10]. Available from <http://www.dh.gov.uk>
2. United Kingdom Department of Health. Health Department influenza pandemic contingency plan. 2005 [cited 2005 Feb 10]. Available from <http://www.dh.gov.uk>
3. United Kingdom Office for National Statistics. 2005 Aug 25 [cited 2005 Oct 2]. Available from <http://www.statistics.gov.uk>
4. Fleming DM. The prediction of epidemics of respiratory infection. *Eur J Epidemiol.* 1994;10:481–3.
5. Turner D, Wailoo A, Nicholson K, Cooper N, Sutton A, Abrams K. Systematic review and economic decision modelling for the prevention and treatment of influenza A and B. *Health Technol Assess.* 2003;7:iii–xiii, 1.
6. Roche UK. Tamiflu product characteristics. 2006 [cited 2006 Oct 10]. Available from <http://www.rocheuk.com>
7. Becton Dickinson USA. Directigen. 2005 [cited 2006 Oct 10]. Available from <http://www.bd.com>
8. Biostar. Biostar Flu OIA. 2005 [cited 2006 Oct 10]. Available from <http://www.biostar.com>



9. Quidel. QuickVue. 2005 [cited 2006 Oct 10]. Available from <http://www.quidel.com>
10. Zymetx. Z-Stat Flu. 2005 [cited 2006 Oct 10]. Available from <http://www.zymetx.com>
11. Chan KH, Maldeis N, Pope W, Yup A, Ozinskas A, Gill J, et al. Evaluation of the Directigen FluA+B test for rapid diagnosis of influenza virus type A and B infections. *J Clin Microbiol.* 2002;40:1675–80.
12. Nicholson KG, Wood JM, Zambon M. Influenza. *Lancet.* 2003;362:1733–45.
13. Reina J, Padilla E, Alonso F, Ruiz dG, Munar M, Mari M. Evaluation of a new dot blot enzyme immunoassay (directigen flu A+B) for simultaneous and differential detection of influenza a and B virus antigens from respiratory samples. *J Clin Microbiol.* 2002;40:3515–7.
14. United Kingdom Department of Health. Guidance on the change in HM Treasury discount rates from 6% to 3.5% from 1 April 2003 [cited 2003 Sep 22]. Available from <http://www.info.doh.gov.uk/doh/finman.nsf>
15. Kaiser L, Wat C, Mills T, Mahoney P, Ward P, Hayden F. Impact of oseltamivir treatment on influenza-related lower respiratory tract complications and hospitalizations. *Arch Intern Med.* 2003;163:1667–72.
16. British National Formulary. BNF 50. 2005 [cited 2006 Oct 10]. Available from <http://www.bnf.org/bnf/bnf/current/openat>
17. Curtis L, Netten A. Unit costs of health and social care. Canterbury (UK): University of Kent, Personal Social Services Research Unit; 2004.
18. United Kingdom Department of Health Hospital episode statistics. 2005 [cited 2006 Oct 10]. Available from <http://www.dh.gov.uk/PublicationsAndStatistics/Statistics/HospitalEpisodeStatistics/fs/en>
19. United Kingdom. Department of Health. NHS reference costs 2004 [cited 2006 Oct 10]. Available from [http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH\\_4105545](http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_4105545)
20. O'Brien BJ, Goeree R, Blackhouse G, Smieja M, Loeb M. Oseltamivir for treatment of influenza in healthy adults: pooled trial evidence and cost-effectiveness model for Canada. *Value Health.* 2003;6:116–25.
21. Melegaro A, Edmunds WJ. Cost-effectiveness analysis of pneumococcal conjugate vaccination in England and Wales. *Vaccine.* 2004;22:4203–14.
22. Kelly H, Birch C. The causes and diagnosis of influenza-like illness. *Aust Fam Physician.* 2004;33:305–9.
23. Sander B, Gyldmark M, Hayden FG, Morris J, Mueller E, Bergemann R. Influenza treatment with neuraminidase inhibitors: cost-effectiveness and cost-utility in healthy adults in the United Kingdom. *Eur J Health Econ.* 2005;6:244–52.
24. Ilyushina NA, Govorkova EA, Webster RG. Detection of amantadine-resistant variants among avian influenza viruses isolated in North America and Asia. *Virology.* 2005;341:102–6.
25. National Institute for Clinical Excellence (NICE). Guidance on the use of zanamivir, oseltamivir and amantadine for the treatment of influenza. 2003.
26. Her Majesty's Treasury. The green book—appraisal and evaluation in central government. 2003 [cited 2006 Oct 10]. Available from [http://www.hm-treasury.gov.uk/economic\\_data\\_and\\_tools/greenbook/data\\_greenbook\\_index.cfm](http://www.hm-treasury.gov.uk/economic_data_and_tools/greenbook/data_greenbook_index.cfm)
27. Kind P, Dolan P, Gudex C, Williams A. Variations in population health status: results from a United Kingdom national questionnaire survey. *BMJ.* 1998;316:736–41.
28. Rawlins MD, Culyer AJ. National Institute for Clinical Excellence and its value judgments. *BMJ.* 2004;329:224–7.
29. Woodall J, Rowson KE, McDonald J. Age and Asian influenza, 1957. *BMJ.* 1958;2:1316–8.
30. Ferguson NM, Cummings DA, Fraser C, Cajka JC, Cooley PC, Burke DS. Strategies for mitigating an influenza pandemic. *Nature.* 2006;442:448–52.
31. Mounier-Jack S, Coker RJ. How prepared is Europe for pandemic influenza? Analysis of national plans. *Lancet.* 2006;367:1405–11.
32. Gubareva LV, Kaiser L, Matrosovich MN, Soo-Hoo Y, Hayden FG. Selection of influenza virus mutants in experimentally infected volunteers treated with oseltamivir. *J Infect Dis.* 2001;183:523–31.
33. Roberts NA. Treatment of influenza with neuraminidase inhibitors: virological implications. *Philos Trans R Soc Lond B Biol Sci.* 2001;356:1895–7.
34. Carr J, Ives J, Kelly L, Lambkin R, Oxford J, Mendel D, et al. Influenza virus carrying neuraminidase with reduced sensitivity to oseltamivir carboxylate has altered properties in vitro and is compromised for infectivity and replicative ability in vivo. *Antiviral Res.* 2002;54:79–88.
35. Herlocher ML, Truscon R, Elias S, Yen HL, Roberts NA, Ohmit SE, et al. Influenza viruses resistant to the antiviral drug oseltamivir: transmission studies in ferrets. *J Infect Dis.* 2004;190:1627–30.
36. Govorkova EA, Ilyushina NA, Boltz DA, Douglas A, Yilmaz N, Webster RG. Efficacy of oseltamivir therapy in ferrets inoculated with different clades of H5N1 influenza virus. *Antimicrob Agents Chemother.* 2007;51:1414–24.
37. McKimm-Breschkin JL, Selleck PW, Usman TB, Johnson MA. Reduced sensitivity of influenza A (H5N1) to oseltamivir. *Emerg Infect Dis* [serial on the internet]. 2007 [cited 2007 Dec 1]. Available from <http://www.cdc.gov/eid/content/13/9/1354.htm>
38. Rameix-Welti MA, Agou F, Buchy P, Mardy S, Aubin JT, Veron M, et al. Natural variation can significantly alter the sensitivity of influenza A (H5N1) viruses to oseltamivir. *Antimicrob Agents Chemother.* 2006;50:3809–15.
39. Lipsitch M, Cohen T, Murray M, Levin BR. Antiviral resistance and the control of pandemic influenza. *PLoS Med.* 2007;0111–21.

Address for correspondence: M. Ruby Siddiqui, Modelling and Economics Unit, Statistics, Modelling and Bioinformatics Department, Health Protection Agency (Centre for Infections), 61 Colindale Ave, London, NW9 5EQ, UK; email: [ruby.siddiqui@hpa.org.uk](mailto:ruby.siddiqui@hpa.org.uk)

All material published in *Emerging Infectious Diseases* is in the public domain and may be used and reprinted without special permission; proper citation, however, is required.

EMERGING INFECTIOUS DISEASES *online*

[www.cdc.gov/eid](http://www.cdc.gov/eid)

To receive tables of contents of new issues send an email to [listserv@cdc.gov](mailto:listserv@cdc.gov) with `subscribe eid-toc` in the body of your message.



**Appendix Table.** Pandemic influenza model parameters

Parameter	Base-case	Distribution	Distribution parameters	Source/assumptions
2004 UK population, millions	59.8	Fixed		(3)
Clinical attack rate	<b>0.25</b>	Normal	SD = 0.05	DoH pandemic plan (2)
Pandemic wave, wks	15	Fixed		DoH pandemic plan (2)
Mean weekly ILI consultation rate	<b>0.00059</b>	Discrete	p = 0.333	Winter (summer = 0.00018, midwinter = 0.00075) (4)
Proportion ILI patients who consult GP	0.28	Fixed		(5)
Overall CFR (1918 scenario)	0.023	Fixed		HPA unpublished data
Overall CFR (1957/69 scenario)	0.003	Fixed		HPA unpublished data
CFR (1957/69) 0–14 y	0.0001	Fixed		HPA unpublished data
CFR (1957/69) 15–44 y	0.0001	Fixed		HPA unpublished data
CFR (1957/69) 45–64 y	0.0019	Fixed		HPA unpublished data
CFR (1957/69) ≥65 y	0.0163	Fixed		HPA unpublished data
Time to pandemic, y	<b>30</b>	Discrete uniform	0–59	Assumption
AV drug/test stockpile (millions)	(14.6)	Fixed		Varied in optimal stockpile analysis (0–35) (1)
Wastage	<b>0.15</b>	Uniform	0.05–0.25	
Probability of receiving timely AV drugs	<b>0.70†</b>	Normal	SD = 0.5	Assumption
Shelf-life of AV drugs, y	5	Triangular	4, 5, 6	(6), assumption
Shelf-life of tests, y	2†	Normal	SD = 0.989	(7–10)
Sensitivity of test	<b>0.895</b>	Normal	SD = 0.033	(11–13)
Specificity of test	<b>0.998‡</b>	Normal	SD = 0.001	(11–13)
Discount rate for costs and benefits	<b>0.035</b>	Fixed		Varied in univariate sensitivity analysis only (0–0.06) (14)
<b>Complications</b>				
OR complication rate, AV drugs vs. none	<b>0.69</b>	Normal	SD = 0.085	(15)
Probability GP or A&E consultation (influenza, treated)	<b>0.05†</b>	Normal	SD = 0.05	DoH pandemic plan (2), assumption
Probability GP or A&E consultation (noninfluenza ILI or influenza, untreated)	0.068			(15)
OR hospitalization rate, AV drugs vs. none	<b>0.39§</b>	Lognormal	Mean = ln(0.39) SD = 0.4103	(15)
Probability of hospitalization				
Influenza, treated	<b>0.0055</b>	Normal	SD = 0.001	DoH pandemic plan (2), assumption
Influenza, untreated	0.014			(15)
Noninfluenza ILI	0.006			40% of probability for influenza, untreated (15)

OR death given AV drugs vs. none	<b>0.39</b> §	Lognormal	Mean = ln(0.73) SD = 0.4103	Assumed the same as hospitalization
Probability of death (not from influenza)	0.015	Fixed		Mean rate for <i>S. pneumoniae</i> (3)
Costs, £				
AV drug per course (treat only)	<b>16</b>	Triangular	1, 16, 17	(16), assumption
Near-patient test	<b>7</b>	Normal	SD = 2.80	(7-10)
Storage cost of course/test (per y)	<b>1</b>	Uniform	1-5	assumption
Administration (per course/test)	15.75†	Normal	SD = £5.09	(17)
Hospitalized (per patient)	<b>756</b>	Triangular	± 15%	(17,18), assumption
GP costs (per patient)	<b>33</b>	Triangular	± 15%	(17,18), assumption
A&E consultation (per patient)	<b>81</b>	Triangular	± 15%	(19), assumption
Health benefits (QALY loss)				
Uncomplicated influenza	0.0046	Normal	SD = 0.0006	0.61 QoL weight, 7 d (20)
Uncomplicated influenza, treated	0.0038¶	Normal	SD = 0.0015	0.65 QoL weight, 7 d (20)
Uncomplicated, noninfluenza	0.0043	Normal	SD = 0.0009	0.62 QoL weight, 7 d (20)
Complications, ILI	0.004	Triangular	± 15%	Pneumonia outpatient (21), assumption.
Hospitalization, ILI#	0.006	Triangular	± 15%	Pneumonia inpatient (21), assumption
QALY loss/influenza death, 1918	16.7			Based on CFR
QALY loss/influenza death, 1957/69	8.1			Based on age-specific CFR
QALY loss/noninfluenza death	6.1			Based on <i>S. pneumoniae</i> deaths (3)

\*UK, United Kingdom; SD, standard deviation; DoH, Department of Health, ILI, influenza-like illness; CFR, case-fatality ratio; HPA, Health Protection Agency; AV, antiviral; OR, odds ratio; GP, general practitioner; A&E, accident and emergency departments; OR, odds ratio; QALY, quality-adjusted life year; QoL, quality of life. **Boldface** indicates univariate sensitivity analysis.

†Truncated at 0.

‡Truncated at 1.

§Truncated at 0 and 1.

¶Total QALY loss associated with complications is the sum of that associated with pneumonia (outpatient) and the uncomplicated condition.

#Total QALY loss associated with hospitalization is the sum of that associated with pneumonia (inpatient) and the uncomplicated condition.