

Molecular Typing of Australian *Scedosporium* Isolates Showing Genetic Variability and Numerous *S. aurantiacum*

Laurence Delhaes,^{*1,2} Azian Harun,^{*†1} Sharon C.A. Chen,^{*1} Quoc Nguyen,^{‡1} Monica Slavin,[§] Christopher H. Heath,^{¶#} Krystyna Maszewska,^{*} Catriona Halliday,^{*} Vincent Robert,^{**} Tania C. Sorrell,^{*†} the Australian Scedosporium (AUSCEDO) Study Group,³ and Wieland Meyer^{*†1}

One hundred clinical isolates from a prospective nationwide study of scedosporiosis in Australia (2003–2005) and 46 additional isolates were genotyped by internal transcribed spacer–restriction fragment length polymorphism (ITS-RFLP) analysis, ITS sequencing, and M13 PCR fingerprinting. ITS-RFLP and PCR fingerprinting identified 3 distinct genetic groups. The first group corresponded to *Scedosporium prolificans* (n = 83), and the other 2 comprised isolates previously identified as *S. apiospermum*: one of these corresponded to *S. apiospermum* (n = 33) and the other to the newly described species *S. aurantiacum* (n = 30). Intraspecies variation was highest for *S. apiospermum* (58%), followed by *S. prolificans* (45%) and *S. aurantiacum* (28%) as determined by PCR fingerprinting. ITS sequence variation of 2.2% was observed among *S. apiospermum* isolates. No correlation was found between genotype of strains and their geographic origin, body site from which they were cultured, or colonization versus invasive disease. Twelve *S. prolificans* isolates from 2 suspected case clusters were examined by amplified fragment length polymorphism analysis. No specific clusters were confirmed.

Despite efforts to identify and eliminate infectious agents, they continue to emerge and reemerge (1). Among them, pathogenic fungi contribute substantially to illness and death, especially in immunocompromised patients (2,3). In contrast to the well-documented opportunists *Candida albicans*, *Cryptococcus neoformans*, and

Aspergillus fumigatus, the epidemiology and evolution of human infections caused by uncommon but emerging fungi are incompletely understood. Such pathogens include *Scedosporium apiospermum* (teleomorph *Pseudallescheria boydii*) and *S. prolificans*, which are inherently resistant to many antifungal agents (3–5).

¹These authors contributed equally to experimental work and data analysis.

²Current affiliation: Lille Pasteur Institute, Lille, France

³Members of the Australian Scedosporium Study Group of the Australasian Society for Infectious Diseases: Australian Capital Territory: Peter Collignon; The Canberra Hospital. New South Wales: Richard Benn (Royal Prince Alfred Hospital); Ian Chambers (Douglass Hanly Moir Pathology); Sharon Chen (Westmead Hospital); Nelson Dennis (Wollongong Hospital); Deo DeWit (Gosford Hospital); John Ferguson (John Hunter Hospital); Iain Gosbell (Liverpool Hospital); Thomas Gottlieb (Concord Hospital); Catriona Halliday (Westmead Hospital); Juliette Holland (Mayne Laverty Pathology); Alison Kesson (New Children's Hospital, Westmead); Richard Lawrence (St. George Hospital); Deborah Marriott (St. Vincent's Hospital, Sydney); Wieland Meyer (Westmead Hospital); Peter Newton (Wollongong Hospital); Quoc Nguyen (St. Vincent's Hospital, Sydney); Pamela Palasanthrian (Sydney Children's Hospital); Robert Pickles (infectious diseases physician, Taree); Robert Pritchard (Royal North Shore Hospital); Tania Sorrell (Westmead Hospital); Lex Tierney (John Hunter Hospital); Voula Tomasotos (Liverpool Hospital); Robert Vaz (Orange Base Hospital); Kerry Weeks (Royal North Shore Hospital). Queensland: Anthony Allworth (Royal Brisbane Hospital); Christopher Coulter (The Prince Charles Hospital); Joan Faoagali (Royal Brisbane Hospital); Barbara Johnson (Princess Alexandra Hospital); David Looke (Princess Alexandra Hospital); Joseph McCormack (The Mater Adult Hospital); Graeme Nimmo (Princess Alexandra Hospital); Gabrielle O'Kane (The Prince Charles Hospital); E. Geoffrey Playford (Princess Alexandra Hospital); Jennifer Robson (Sullivan and Nicolaidis Pathology). South Australia: David Ellis (Women's and Children's Hospital); Rosemary Handke (Women's and Children's Hospital); Karen Rowlands (Royal Adelaide Hospital); David Shaw (Royal Adelaide Hospital). Tasmania: Louise Cooley (Royal Hobart Hospital); Erica Cox (Launceston General Hospital); Alistair McGregor (Royal Hobart Hospital). Victoria: Clare Franklin (Alfred Hospital); Cathy Joseph (St Vincent's Hospital, Melbourne); Tony Korman (Monash Medical Centre); Orla Morrissey (Alfred Hospital); Monica Slavin (Peter MacCallum Cancer Centre); Denis Spelman (Alfred Hospital); Bryan Speed (Austin and Repatriation Hospitals); Harsha Sheorey (St. Vincent's Hospital, Melbourne). Western Australia: Peter Boan (Royal Perth Hospital); John Dyer (Fremantle Hospital); Christopher Heath (Royal Perth Hospital); Dianne Gardam (Royal Perth Hospital); Duncan McLennan (Fremantle Hospital); Ronan Murray (Royal Perth Hospital); Todd Pryce (Royal Perth Hospital).

*Westmead Hospital, Westmead, New South Wales, Australia; †University of Sydney, Sydney, New South Wales, Australia; ‡St. Vincent's Hospital, Sydney, New South Wales, Australia; §Alfred Hospital, Melbourne, Victoria, Australia; ¶Royal Perth Hospital, Perth, Western Australia, Australia; #University of Western Australia, Perth, Western Australia, Australia; and **Centraalbureau voor Schimmelcultures, Utrecht, the Netherlands

S. apiospermum infections occur worldwide, ranging from localized mycetomas to deep-seated disease such as cerebral abscesses (6,7). This species also colonizes the respiratory tract of $\approx 10\%$ of patients with cystic fibrosis and chronic suppurative lung disease (8–10). On the basis of genetic data, a new species, *S. aurantiacum*, was proposed for a subset of isolates previously identified as *S. apiospermum* (11). *S. prolificans* infections are geographically more restricted than those caused by *S. apiospermum*, being most prevalent in Australia, Spain, and the United States (12–15). *S. prolificans* typically causes localized infections in immunocompetent hosts but rapidly fatal disseminated infections in the immunocompromised among whom it has been associated with nosocomial outbreaks (3,12–17).

Since scedosporiosis, in particular that caused by *S. prolificans*, is often refractory to treatment (3,5), preventive strategies are of paramount importance. However, the epidemiology and mode of transmission of infection are not well understood. Furthermore, the environmental reservoir of *S. prolificans* is unknown. Molecular typing techniques now provide the means to elucidate the epidemiology of *Scedosporium* infections and to investigate potential case clusters (16,18,19). Strains recovered from patients with cystic fibrosis have demonstrated a high degree of genetic variability (10,20), although a single genetic profile predominated in 1 study (8). The degree of genetic variation within *S. prolificans* is more controversial. Two studies have reported low to no intraspecies genetic heterogeneity (16,21), while a third noted substantial genetic diversity (19). The results of these studies may be biased because they included only small numbers of isolates from specific patient populations. Genetic variability among *S. aurantiacum* has not yet been studied.

In this study, we used 4 molecular tools to examine genetic variation among a large number of Australian clinical *Scedosporium* isolates: 1) internal transcribed spacer (ITS)-based restriction fragment length polymorphism (ITS-RFLP) analysis; 2) DNA sequence analysis of the ITS region (selected isolates); 3) PCR fingerprinting using the microsatellite specific core sequence of phage M13; and 4) amplified fragment length polymorphism (AFLP) analysis (isolates from suspected case clusters). We also searched for the newly described species, *S. aurantiacum* and for genetic clustering of strains according to their geographic origin, body site from which they were cultured, and ability to cause invasive disease.

Materials and Methods

Scedosporium Isolates and Data Collection

A total of 146 *Scedosporium* isolates were studied (online Technical Appendix, available from www.cdc.gov/EID/content/14/2/282-Techapp.pdf). Forty-six were

from the culture collection at the Clinical Mycology Laboratory, Centre for Infectious Diseases and Microbiology Laboratory Services, Westmead Hospital, Sydney, Australia. For these isolates, the following data were captured: demographic information, patient coexisting conditions and risk factors (summarized in the online Technical Appendix). The remaining 100 isolates were obtained through a national, prospective, laboratory-based surveillance for scedosporiosis in Australia (the Australian *Scedosporium* [AUSCEDO] Study) from January 2003 to December 2005. The following data were collected: clinical status, risk factor (defined according to published risk factors for scedosporiosis [4,12–15]), major comorbidity (based on the International Classification of Diseases, 10th revision, Australian Modification [ICD-10 AM] diagnostic classification system [22]), isolated species, treatment and outcome. *Scedosporium* strains obtained from a single colony from the primary isolation plate from all patients were forwarded to the Molecular Mycology Research Laboratory, Westmead Hospital, for genotyping. Isolates were identified as *S. prolificans* or *S. apiospermum* by standard phenotypic methods (23). Species were confirmed as *S. prolificans* or *S. apiospermum*, and *S. aurantiacum* was identified (11) by ITS-RFLP analysis.

Definitions

An episode of scedosporiosis was defined as the incident isolation of *Scedosporium* spp. from any body site. Two or more episodes, fulfilling the case definition and occurring in different patients that were epidemiologically linked were defined as a potential case cluster. Invasive disease was defined according to the European Organization for Treatment of Cancer/Mycoses Study Group criteria for “definite” or “probable” infection (24). All other patients not fulfilling these criteria, including those with “possible” infection were considered colonized. Coincident hospital renovations or construction was considered to be a potential risk factor if major work was undertaken within 3 months before the isolation of *Scedosporium* spp. from a patient.

Description of 2 Potential Case Clusters

The first potential case cluster involved 8 patients located in the same hematology/hemopoietic stem cell transplant (HSCT) unit at the Alfred Hospital, a large university hospital in Melbourne (September 2000–October 2001; [15]). The second consisted of 3 patients located in the same hematology/HSCT ward at Westmead Hospital a major university hospital in Sydney (September 2003–January 2004; unpub. data). Details of the patients involved in these suspected case clusters are summarized in the online Technical Appendix). On each occasion, patient isolates were submitted for genetic analyses to inform infection control responses (see Results).

Genomic DNA Extraction and ITS-RFLP Analysis

Genomic DNA was isolated as described previously (18). The ITS1, 5.8S, and ITS2 regions of the rDNA gene cluster were amplified with the primers SR6R and LR1 (Table 1) as described previously (25). Amplicons were double digested with the restriction endonucleases *Sau96I* and *HhaI* (New England BioLabs, Ipswich, MA, USA) in accordance with the manufacturer's recommendations. Digested products were separated by electrophoresis in 3% agarose gels at 100 V for 3–4 h. Banding patterns were analyzed visually.

ITS Sequencing

Eleven isolates, representative of each of 3 ITS-RFLP patterns obtained, were selected for ITS sequencing: ITS-RFLP profile A (*S. prolificans*, WM 06.378, WM 06.440, and WM 06.393), ITS-RFLP profile B (*S. apiospermum*, WM 06.389, WM 06.471, and WM 06.497), and ITS-RFLP profile C (*S. aurantiacum*, WM 06.388, WM 06.482, WM 06.495, WM 06.496, and WM 06.498). The ITS region was amplified as described above and commercially sequenced in both directions by using SR6R or LR1 (Table 1) as forward and reverse primers.

PCR Fingerprinting

The minisatellite-specific core sequence of the wild-type phage M13 was used as a single primer for PCR fingerprinting (Table 1). Amplification reactions were performed as previously described (18). Blank control tubes containing all reagents except template DNA were included for each run; each sample was analyzed at least twice. PCR products were separated by electrophoresis on 1.4% agarose gels at 60 V for 14 cm. Strains were defined to be identical if their PCR fingerprinting profiles had a similarity of $\geq 97\%$ (= 1 band difference). Reproducibility of the PCR fingerprinting technique was assessed by re-amplifying 1 strain of each of the 3 *Scedosporium* spp. with all PCR amplifications carried out and re-running those on each gel.

AFLP Analysis

AFLP analysis was performed as described previously by using either *EcoRI*-GT 6-FAM-labeled and *MseI*-GT or *EcoRI*-TC 6-FAM-labeled and *MseI*-CA as selective primer pairs (QIAGEN, Valencia, CA, USA; Table 1) (26). All samples were analyzed by using the ABI Prism 3730 system (Applied Biosystems, Foster City, CA, USA). Data collation, fragment sizing, and pattern analyses were performed with GeneMapper software version 3.5 (Applied Biosystems). Only electrophoregram peaks above 1,000 fluorescent units were scored for the presence or absence of bands of the same size (range 50–500 bp) relative to the GeneScan 500 LIZ DNA size standard (Applied Biosystems). Only bands detected in duplicate AFLP experiments were included in the analysis.

Data Analysis

Clinical Data

Statistical analysis was performed by using SPSS version 10.0.07 (SPSS, Chicago, IL, USA) and EpiInfo version 6.0 (Centers for Disease Control and Prevention, Atlanta, GA, USA). Proportions were compared by using the χ^2 or Fisher exact test. A p value < 0.05 was statistically significant.

ITS Sequences

ITS sequences obtained from 11 isolates (see above) were aligned with the ITS sequences of the following reference strains obtained from GenBank: *S. apiospermum* CBS 101.22 (accession no. AJ888435), *S. aurantiacum* FMR 8630 (accession no. AJ888440), *S. aurantiacum* IHEM 15458 (accession no. AJ888441) and *S. prolificans* CBS 114.90 (accession no. AY882369) as well as 2 outgroup sequences: *Pseudallescheria africana* CBS 311.72 (accession no. AJ888425), and *Petriella setifera* CBS 164.74 (accession no. AY882352). Phylogenetic analyses were performed by using PAUP* version 4.06.10 (27).

PCR Fingerprinting Patterns and AFLP Fragments

PCR fingerprinting patterns were analyzed by using the 1D gel analysis module (BioGalaxy [BioAware, Hantut, Belgium]) in BioloMICS version 7.5.30 (BioAware). Images were normalized for lane to-lane differences in mobility by the alignment of patterns obtained on multiple loadings of the 1kb DNA size marker (GIBCO-BRL, Gaithersburg, MD, USA). The unweighted-pair group method by using arithmetic averages and the procedures of Nei and Li (28), both implemented in BioloMICS, were used to generate dendrograms based on the coefficient of similarity (29) between the isolates. In addition, principal coordinate analysis (PcoA; BioloMICS) was conducted to give an overall representation of the observed strain variation. AFLP fragments were analyzed with BioloMICS.

Results

A total of 146 *Scedosporium* isolates from 120 episodes (119 patients) were studied (online Technical Appendix). Demographic data were available for 108 (90%) episodes and coexisting conditions and risk factor data for 115 (95.8%). Most episodes were reported from New South Wales (64.2%), followed by Victoria (19.2%) and Western Australia (9.2%). The male: female ratio was 1.3: 1. The major patient coexisting conditions and known risk factors for scedosporiosis are summarized in the online Technical Appendix. Thirty-nine patients (32.7%) had no underlying medical condition. Coincident building construction was noted in 27 cases (22.5%). *Scedosporium* isolates were associated with invasive disease in 46 (38.3%) instances;

Table 1. Primer and adaptor oligonucleotide sequences used in the study

Primer or adaptor oligonucleotide	Sequence (5'→3')
rDNA primers	
SR6R	AAGTARAAGTCGTAACAAGG
LR1	GGTTGGTTTCTTTTCCT
M13 fingerprinting primer	
Phage M13	GAGGGTGGCGGTTCT
EcoRI adapters	
EA1	CTCGTAGACTGCGTACC
EA2	CATCTGACGCATGGTTAA
MseI adapters	
MA1	GACGATGAGTCCTGAG
MA2	TACTCAGGACTCAT
Preselective primers	
EcoRI-T	GACTGCGTACCAATTCT
EcoRI-G	GACTGCGTACCAATTCG
MseI-C	GATGAGTCCTGAGTAAC
MseI-G	GATGAGTCCTGAGTAAG
Selective primers	
EcoRI-TG	6 FAM-GACTGCGTACCAATTCTG
EcoRI-GT	6 FAM-GACTGCGTACCAATTCGT
MseI-CA	GATGAGTCCTGAGTAACA
MseI-GT	GATGAGTCCTGAGTAAGT

the remaining 74 (61.7%) were isolated from patients who were colonized (Table 2).

Molecular Typing of *Scedosporium* Isolates

All 146 isolates were examined by ITS-RFLP analysis and PCR fingerprinting. ITS sequencing was performed on 11 strains as described above. AFLP analysis was performed only for selected *S. prolificans* isolates, including the isolates of the suspected case clusters and isolates representative of the *S. prolificans* branches identified by PCR fingerprinting (online Appendix Figure 1, available from www.cdc.gov/EID/content/14/2/282-appG1.htm).

ITS-RFLP Analysis

RFLP analysis found 1 RFLP profile specific for *S. prolificans* isolates (ITS-RFLP profile A) and 2 profiles (ITS-RFLP profiles B and C) for isolates previously phenotypically identified as *S. apiospermum* (Figure 1, panel A). ITS-RFLP profile B corresponded to *S. apiospermum* and ITS-RFLP profile C to the newly described species, *S. aurantiacum*.

ITS Sequencing

Sequencing of the ITS 1, 5.8S, and ITS2 regions of the 11 strains, representative of each of the 3 ITS-RFLP profiles found the following results: BLAST searches against the corresponding GenBank reference sequences identified strains: WM 06.389 (accession no. EF639870), WM 06.497 (accession no. EF639872), and WM 06.471 (accession no. EF639871) (ITS-RFLP profile B) as *S. apiosper-*

mum (96%–99% sequence similarity to strain CBS 101.22). Strains WM 06.388 (accession no. EF639865), WM 06.482 (accession no. EF639866), WM 06.495 (accession no. EF639867), WM 06.496 (accession no. EF639868), and WM 06.498 (accession no. EF639869) (ITS-RFLP profile C) were identified as *S. aurantiacum* (100% sequence identity with strains FMR 8630 and IHEM 15458). Isolates WM 06.393 (accession no. EF639863), WM 06.440 (accession no. EF639864) and WM 06.378 (accession no. EF639862) (ITS-RFLP profile A) were identified as *S. prolificans* (100% identity with strain CBS 114.90).

Phylogenetic analysis of the sequences demonstrated 3 distinct clades, the first corresponding to *S. prolificans* as the basal clade. The other 2 corresponded to the 2 more closely related but clearly distinct clades, *S. apiospermum*, and *S. aurantiacum* (Figure 2). *S. apiospermum* showed intraspecies sequence variation of 2.2% compared to *S. aurantiacum* and *S. prolificans*, which displayed no variation.

Final Identification of *Scedosporium* spp. and Clinical Associations

S. prolificans accounted for 75 patient episodes (83 of 146 isolates; 56.9%), *S. apiospermum* for 25 (33 isolates; 22.6%), and *S. aurantiacum* for 23 (30 isolates; 20.6%) (online Technical Appendix). More than 1 *Scedosporium* spp. was isolated from the same patient in 3 instances: Patient 83: *S. apiospermum* (WM 06.471, WM 06.472, WM 06.474, and WM 06.475) and *S. prolificans* (WM 06.473); patient 91: *S. apiospermum* (WM 06.486) and *S. prolificans* (WM 06.485); and patient 102: *S. apiospermum* (WM 06.500) and *S. prolificans* (WM 06.501) (online Technical Appendix). In 6 episodes, the same species was recovered from more than 1 body site in the same patient at the same time (patients 57 [blood, bronchial washing, skin], 73 [blood, sputum], 80 [sputum, bone, wound fluid], 83 [bronchial washing, bronchoalveolar lavage], 118 [pleural fluid, bone, wound fluid, chest tissue], and 119 [blood, skin]; online Technical Appendix).

Approximately half (40%–52.2%) of *S. apiospermum* and *S. aurantiacum* isolates were from the respiratory tract/lung compared to 20% for *S. prolificans*. Conversely, all isolates from blood, 57.2% isolates from skin/soft tissue and 66.7% from eye were *S. prolificans* (Table 2). Invasive disease was more likely to be caused by *S. prolificans* than non-*prolificans* *Scedosporium* spp. (83% versus 17% of isolations; odds ratio (OR) 5.3, 95% confidence interval (CI) 2.0, 14.2, $p = 0.002$) (Table 2). This association was significant when compared with *S. apiospermum* as well as with *S. aurantiacum* ($p < 0.05$; data not shown). The relative proportions of invasive disease among *S. apiospermum* and *S. aurantiacum* were similar (Table 2). Coincident building construction (27 cases, 22.5%) was more likely to be




Figure 1. Internal transcribed spacer–restriction fragment length polymorphism (ITS-RFLP) patterns obtained by double digestion with the enzymes *Sau96I* and *HhaI* (A) and of the PCR fingerprinting profiles obtained with the microsatellite specific primer M13 (B) for *Scedosporium proliferans*: lane 1, WM 06.457; lane 2, WM 06.458; lane 3, WM 06.503; lane 4, WM 06.502; lane 5, WM 06.399; lane 6, WM 06.434. *S. aurantiacum*: lane 7, WM 06.495; lane 8, WM 06.496; lane 9, WM 06.386; lane 10, WM 06.385; lane 11, WM 06.482; lane 12, WM 06.390. *S. apiospermum*: lane 13, WM 06.475; lane 14, WM 06.474; lane 15, WM 06.472; lane 16, WM 06.471; lane 17, WM 06.424; lane 18, WM 06.443; lane M, 1-kb marker (GIBCO-BRL, Gaithersburg, MD, USA).

associated with isolation of *S. proliferans* compared with non-*prolificans* *Scedosporium* spp. (OR 11.5, 95% CI 2.4, 74.5; $p < 0.001$; data not shown).

Molecular Epidemiology

Strain Typing

PCR fingerprinting delineated 3 major clusters concordant with *S. apiospermum*, *S. aurantiacum*, and *S. proliferans* (online Appendix Figure 1; Figure 1, panel B; Figure 3). Clusters corresponding to *S. aurantiacum* and *S. proliferans* were substantially more densely grouped than the *S. apiospermum* cluster (Figure 3).

PCR fingerprinting profiles showed polymorphisms within each of the 3 species, allowing for a clear differentiation, by using a “cut-off point” of $\geq 97\%$ similarity. Multiple isolates from the same patient obtained from different anatomic sites (online Technical Appendix) had identical or $\geq 97\%$ similarity between their PCR fingerprints, except for 1 patient (patient 118). In 8 instances, PCR fingerprinting showed that patients were infected with 2 different strains: (patients 1, 10, 27, 57, 83, 99, 118 [online Appendix Figure 1, online Technical Appendix]). For all species, genetic profiles were independent of geographic origin, body site of isolation or whether the patient was infected or colonized (online Appendix Figure 1). Profiles were also independent of patient comorbidity and risk factors for scedosporiosis (data not shown). Intraspecies PCR fingerprinting variation was highest for *S. apiospermum* (58%) followed by *S. proliferans* (45%) and *S. aurantiacum* (28%) (online Appendix Figure 1).

Examination of Isolates from Suspected Case Clusters

Twelve isolates from 2 presumptive case clusters of *S. proliferans* infection (Alfred Hospital, Melbourne pa-




Figure 2. Rooted phylogram (outgroup *Pseudallescheria africana* CBS 311.72 and *Petriella setifera* CBS 164.74), showing the relationships among 11 selected strains representing each obtained internal transcribed spacer (ITS)–restriction fragment length polymorphism pattern and 4 reference strain sequences obtained from GenBank by using PAUP* version 4.06.10 (29).

Table 2. Selected characteristics for 120 isolations (episodes) of *Scedosporium* spp.

Characteristic*	No. <i>Scedosporium prolificans</i> ; n = 75 (%)†	No. <i>S. apiospermum</i> ; n = 25 (%)†	No. <i>S. aurantiacum</i> ; n = 23 (%)†
Male sex	40 (53.3)	12 (48)	8 (34.8)
Risk factor			
Surgery ≤30 d	3 (4)	1 (4)	—
Trauma	5 (6.7)	—	1 (4.4)
Clinical status‡			
Invasive disease	39 (52)	4 (16)	4 (17.4)
Colonization	36 (48)	21 (84)	19 (82.6)
Body site of isolation			
Blood	18 (24)	—	—
Eye	2 (2.7)	—	1 (4.4)
Skin/soft tissue	4 (5.3)	1 (4)	2 (8.7)
Lung/respiratory tract	15 (20)	10 (40)	12 (52.2)
Ear	4 (5.3)	10	5 (21.2)

*Some patients had *Scedosporium* isolated from more than 1 body site.

†Refers to no. episodes in which each species was isolated. The total no. of isolates was 146 comprising 83 *S. prolificans*, 33 *S. apiospermum*, and 30 *S. aurantiacum*.

‡More than 1 *Scedosporium* spp. was isolated from 4 patients.

tients: isolates WM 06.392, WM 06.393, WM 06.395, WM 06.399, WM 06.400, WM 06.401, WM 06.402, and WM 06.405; Westmead Hospital, Sydney patients: isolates WM 06.432, WM 06.434, WM06.457, and WM 06.458; online Technical Appendix) as well as 23 additional isolates, representative of the *S. prolificans* branches identified by PCR fingerprinting (online Appendix Figure 1) were further investigated by AFLP typing. *S. prolificans* was not isolated from the environment in either setting despite extensive sampling. The AFLP bands were found to be 50–493 bp by using the primers *EcoRI*-GT and *MseI*-GT (data not shown), and from 52–468 bp by using the primers *EcoRI*-TG and *MseI*-CA (online Appendix Figure 2, available from www.cdc.gov/EID/content/14/2/282-appG2.htm). These 35 isolates exhibited 32 different AFLP profiles, with isolates from the same patient (patients 1, 73, and 119) showing identical profiles (online Appendix Figure 2), confirming the PCR fingerprinting results (online Appendix Figure 1). PcoA of the combined AFLP and PCR fingerprinting data demonstrated no clustering of these isolates (Figure 4), which ruled out the possibility of nosocomial transmission.

Discussion

We examined genetic variation among a large number of population-derived *Scedosporium* isolates across the Australian continent. In line with previously reported genetic variability in the *S. apiospermum*/*P. boydii* species complex (30–32), we observed 2 distinct ITS-RFLP patterns among *S. apiospermum* isolates, showing the presence of the newly described species *S. aurantiacum* (11). Notably, we have identified by ITS sequencing that *S. aurantiacum* comprised 45% of the current collection of Australian “*S. apiospermum*” isolates and documents genetic variability within *S. aurantiacum*.

Epidemiologic investigation of *Scedosporium* infection requires accurate identification and typing. *S. apiospermum*, *S. aurantiacum*, and *S. prolificans* were clearly distinguished from each other by PCR fingerprinting and ITS-RFLP analysis. This is consistent with previous rDNA sequence-based studies (30,33,34). The observation of 2 distinct genetic groups, corresponding to *S. aurantiacum* and *S. apiospermum*, supports the proposal that *S. aurantiacum* be designated a separate species (11). This proposal is also supported by the 5%–10% ITS sequence variation found between *S. aurantiacum* and *S. apiospermum* compared to an absence of intraspecies variation in *S. aurantiacum* and *S. prolificans* and a 2.2% variation in *S. apiospermum* (30,32; current study).

Using PCR fingerprinting, intraspecies variation was greatest (58%) among *S. apiospermum* isolates (Figure 3). This diversity is generally consistent with the high degree of polymorphism (15–20 genotypes) previously found (10,20,32). In contrast, genetic variation was lowest (28%) among the *S. aurantiacum* isolates (online Appendix Figure 1; Figure 3). Nevertheless, PCR fingerprinting polymorphisms clearly differentiated all 30 strains (online Appendix Figure 1). Further genotyping studies of a greater number of and more geographically diverse *S. aurantiacum* isolates are warranted.

The intraspecies PCR fingerprint variation in *S. prolificans* (45%) was greater than that in *S. aurantiacum* but less than that in *S. apiospermum*. Given that *S. aurantiacum* is phylogenetically more closely related to *S. apiospermum* than to *S. prolificans* (11,33; current study), this result was unexpected. It may be due to different evolutionary pressures acting on the 3 different species or the relatively small numbers of *S. aurantiacum* isolates studied to date. The moderate genetic diversity among *S. prolificans* confirms previous findings (19). Despite the observed




Figure 3. Three-dimensional presentation of the principal coordinate analysis of the PCR fingerprinting data showing 3 distinct clusters which correspond to *Scedosporium prolificans* (black dots), *S. aurantiacum* (dark gray dots), and *S. apiospermum* (light gray dots), with *S. apiospermum* showing the highest genetic variation.

polymorphisms, PcoA of PCR fingerprint profiles showed dense clustering for *S. prolificans* (Figure 3), which is consistent with the low to absent intraspecies variability in *S. prolificans* found by others (20,21,33). These apparently contradictory findings emphasize the importance of choosing the optimum molecular typing tool with the most appropriate discriminatory power for the organism or species being studied.

The high degree of intraspecies variation detected by PCR fingerprinting and AFLP analysis supports the use of these methods to establish genetic relatedness between isolates recovered from different patients or multiple isolates from the same patient. In comparison, the variation detected by ITS-RFLP analysis and ITS sequencing corresponded to interspecies variation, which makes those techniques ideal for identification of any given isolate to the species level. Individual patients are most likely infected or colonized with genetically distinct strains (19–21; this study). Identical PCR fingerprint or AFLP profiles were noted in multiple isolates recovered simultaneously from different anatomic sites in the same patient (21; current study). However, 8 patients were infected or colonized by at least 2 strains as reflected by their different genetic profiles (online Technical Appendix). Possible explanations include concomitant infection by multiple strains from which only a restricted number were recovered, or colonization by 1 strain followed by infection or colonization with a second strain of a different genotype. Longitudinal genotyping studies are required to determine the likelihood that persistence of ≥ 1 genotypes later leads to clinically important

infection or whether the disease is more likely to be caused by an unrelated genotype. In this context, the development of a multilocus sequence typing scheme for *Scedosporium*, as has been developed for *Candida* spp. (35), would be of great advantage to overcome interlaboratory reproducibility problems, which are known to be associated with PCR fingerprinting or AFLP data. However, developing such a scheme remains cumbersome due to the current lack of genomic data of *Scedosporium* spp.

For all 3 *Scedosporium* spp., there was no clustering of strains according to their geographic or body site of origin or by their ability to cause invasive disease, which is in agreement with previous findings for *S. apiospermum* (20,30) and *S. prolificans* (16,17,21). Of note, no specific genotypes were associated with underlying medical conditions or risk factors. Compared with *S. apiospermum* and *S. aurantiacum* *S. prolificans* was more frequently associated with coincident hospital renovation, and invasive disease, had a greater predilection to cause disseminated infection and was the predominant species isolated from blood and other sterile sites (12–16,36; current study). Our preliminary observations indicate that the epidemiology and clinical relevance of recovering *S. aurantiacum* may be similar to that of *S. apiospermum*. *S. aurantiacum* has been reported to colonize the respiratory tract of at-risk patients (8).




Figure 4. Three-dimensional presentation of the principal coordinate analysis of the combined M13 PCR fingerprinting, amplified fragment length polymorphism (AFLP) primers EcoRI-GT and MseI-GT, and AFLP primers EcoRI-TG and MseI-CA data from the suspected Sydney and Melbourne case cluster isolates and 23 other Australian isolates. None of the investigated isolates showed any epidemiologic connection except 3 isolates obtained from the same patient (nos. 1, 73, 119). Blue dots, Melbourne outbreak isolates; pink dot, Melbourne-related isolate; red dots, Sydney outbreak isolates; green dots, Sydney-related isolates; yellow dots, unrelated Australian isolates.

In addition to PCR fingerprinting, we applied AFLP analysis to investigate the possibility of 2 case clusters caused by *S. prolificans*. AFLP analysis was chosen as an independent technique using 2 combinations of selective primers (Table 1), which have been previously shown to have good discriminatory power for fungal strain differentiation (26). Both techniques, previously used to identify outbreak strain clusters in the recent cryptococcosis outbreak on Vancouver Island (37), generated in the current situation distinct patterns from all *S. prolificans* isolates except serial isolates obtained from the same patient (online Appendix Figures 1, 2). These findings exclude the occurrence of nosocomial outbreaks or any close relationship with the nonoutbreak isolates, a result similar to those obtained previously (38). Overall nosocomial acquisition of infection has been demonstrated in only 2 instances (16,17). *Scedosporium* spp. have rarely been isolated from hospital air or from indoor or outdoor surface samples (13,39,40, current study), which raises questions about the mode of acquisition by patients and the mechanisms of the selection of this specific fungus as an infectious agent from among the high biodiversity of environmental molds.

In conclusion, ITS-RFLP analysis is a powerful tool for distinguishing between isolates of the new species *S. aurantiacum* and *S. apiospermum*. PCR fingerprinting and AFLP analysis are useful techniques for determining genetic relatedness between *Scedosporium* isolates and for investigating potential case clusters.

Acknowledgments

We thank all participating infectious disease physicians, clinical microbiologists, and hospital scientists. We also thank Nathalie van de Wiele for assistance in figure preparation, and Dee Carter and Tien Bui for sharing their AFLP expertise.

This study was supported by an NH&MRC project grant no. 352303 to W.M. and V.R., a Center for Infectious Diseases and Microbiology–Public Health start-up grant to W.M. and S.C.A.C., and a Merck, Sharp and Dohme, Australia grant to S.C.A.C. and W.M. This is a publication of the AUSCEDO and ECMM-ISHAM working groups on *Pseudallescheria/Scedosporium* infections.

S.C.A.C. is a member of the Antifungal Advisory Board of Gilead Sciences and Pfizer Australia. C.H.H., M.S., and T.C.S. are or have been on Antifungal Advisory Boards for Gilead Sciences; Pfizer Australia; Merck, Sharp and Dohme, Australia; and Schering-Plough, Australia. T.S. has been on a Global Advisory Board for Pfizer US. T.C.S., M.S., and S.C.A.C. have received untied project funding from Pfizer US; Pfizer Australia; Merck, Sharp and Dohme Australia; and Gilead Sciences. T.C.S. has also received funding from Merck US.

Dr Delhaes is the head of the filamentous fungal research group EA3609 at the Pasteur Institute in Lille, France. Her major

research interests are systemic mycoses due to *Aspergillus* and *Scedosporium*, including molecular epidemiology, nosocomial aspects of the diseases, and oxidative stress response.

References

- Rappuoli R. From Pasteur to genomics: progress and challenges in infectious diseases. *Nat Med*. 2004;10:1177–85.
- Nucci M. Emerging moulds: *Fusarium*, *Scedosporium* and Zygomycetes in transplant patients. *Curr Opin Infect Dis*. 2003;16:607–12.
- Walsh TJ, Groll A, Hiemenez J, Fleming R, Roilides E, Anaissie E. Infections due to emerging and uncommon medically-important fungal pathogens. *Clin Microbiol Infect*. 2004;10(Suppl 1):48–66.
- Steinbach WJ, Perfect JR. *Scedosporium* species infections and treatments. *J Chemother*. 2003;15(Suppl 2):16–27.
- Gilgado F, Serena C, Cano J, Gene J, Guarro J. Antifungal susceptibilities of the species of the *Pseudallescheria boydii* complex. *Antimicrob Agents Chemother*. 2006;50:4211–3.
- Guarro J, Kantarcioglu AS, Horre R, Rodriguez-Tudela JL, Cuenca Estrella M, Berenguer J, et al. *Scedosporium apiospermum*: changing clinical spectrum of a therapy-refractory opportunist. *Med Mycol*. 2006;44:295–327.
- Campagnaro EL, Woodside KJ, Early MG, Gugliuzza KK, Colome-Grimmer MI, Lopez FA, et al. Disseminated *Pseudallescheria boydii* (*Scedosporium apiospermum*) infection in a renal transplant patient. *Transpl Infect Dis*. 2002;4:207–11.
- Williamson EC, Speers D, Arthur IH, Harnett G, Ryan G, Inglis TJ. Molecular epidemiology of *Scedosporium apiospermum* infection determined by PCR amplification of ribosomal intergenic spacer sequences in patients with chronic lung disease. *J Clin Microbiol*. 2001;39:47–50.
- Cimon B, Carrere J, Vinatier JF, Chazalotte JP, Chabasse D, Bouchara JP. Clinical significance of *Scedosporium apiospermum* in patients with cystic fibrosis. *Eur J Clin Microbiol Infect Dis*. 2000;19:53–6.
- Defontaine A, Zouhair R, Cimon B, Carrere J, Bailly E, Symoens F, et al. Genotyping study of *Scedosporium apiospermum* isolates from patients with cystic fibrosis. *J Clin Microbiol*. 2002;40:2108–14.
- Gilgado F, Cano J, Gene J, Guarro J. Molecular phylogeny of the *Pseudallescheria boydii* species complex: proposal of two new species. *J Clin Microbiol*. 2005;43:4930–42.
- Wood GM, McCormack JG, Muir DB, Ellis DH, Ridley MF, Pritchard R, et al. Clinical features of human infection with *Scedosporium inflatum*. *Clin Infect Dis*. 1992;14:1027–33.
- Idigoras P, Perez-Trallero E, Pineiro L, Larruskain J, Lopez-Lopategui MC, Rodriguez N, et al. Disseminated infection and colonization by *Scedosporium prolificans*: a review of 18 cases, 1990–1999. *Clin Infect Dis*. 2001;32:e158–65.
- Berenguer J, Rodriguez-Tudela JL, Richard C, Alvarez M, Sanz M, Gaztelurrutia L, et al. Deep infections caused by *Scedosporium prolificans*. A report on 16 cases in Spain and a review of the literature. *Scedosporium prolificans* Spanish Study Group. *Medicine (Baltimore)*. 1997;76:256–65.
- Cooley L, Spelman D, Thursky K, Slavin M. Infection with *Scedosporium apiospermum* and *S. prolificans*, Australia. *Emerg Infect Dis*. 2007;13:1170–7.
- Ruiz-Diez B, Martin-Diez F, Rodriguez-Tudela JL, Alvarez M, Martinez-Suarez JV. Use of random amplification of polymorphic DNA (RAPD) and PCR-fingerprinting for genotyping a *Scedosporium prolificans* (*inflatum*) outbreak in four leukemic patients. *Curr Microbiol*. 1997;35:186–90.
- Guerrero A, Torres P, Duran MT, Ruiz-Diez B, Rosales M, Rodriguez-Tudela JL. Airborne outbreak of nosocomial *Scedosporium prolificans* infection. *Lancet*. 2001;357:1267–8.

18. Meyer W, Maszewska K, Amirmostofian M, Igreja RP, Hardtke C, Methling K, et al. Molecular typing of global isolates of *Cryptococcus neoformans* var. *neoformans* by PCR-fingerprinting and RAPD—a pilot study to standardize techniques on which to base a detailed epidemiological survey. *Electrophoresis*. 1999;20:1790–9.
19. Sole M, Cano J, Rodriguez-Tudela JL, Ponton J, Sutton DA, Perrie R, et al. Molecular typing of clinical and environmental isolates of *Scedosporium prolificans* by inter-simple-sequence-repeat polymerase chain reaction. *Med Mycol*. 2003;41:293–300.
20. Zouhair R, Defontaine A, Ollivier C, Cimon B, Symoens F, Hallet J-N, et al. Typing of *Scedosporium apiospermum* by multilocus enzyme electrophoresis and random amplification of polymorphic DNA. *J Med Microbiol*. 2001;50:925–32.
21. San Millan R, Quindos G, Garaizar J, Salesa R, Guarro J, Ponton J. Characterization of *Scedosporium prolificans* clinical isolates by randomly amplified polymorphic DNA analysis. *J Clin Microbiol*. 1997;35:2270–4.
22. International statistical classification of disease and related health problems, 10th revision, Australian Modification (ICD-10-AM). Sydney (Australia): National Centre for Classification in Health, University of Sydney; 1998.
23. de Hoog GS, Guarro J, Gene J, Figueras MJ. Hyphomycetes: Genus: *Scedosporium*. In: Atlas of clinical fungi. 2nd ed. Utrecht (the Netherlands): Centralbureau voor Schimmelcultures/Universitat Rovira i Virgili; 2000. p. 899–901.
24. Ascioğlu S, Rex JH, de Pauw B, Bennett JE, Bille J, Crokaert F, et al. Defining opportunistic invasive fungal infections in immunocompromised patients with cancer and hematopoietic stem cell transplants: an international consensus. *Clin Infect Dis*. 2002;34:7–14.
25. Vilgalys R, Hester M. Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *J Bacteriol*. 1990;172:4238–46.
26. Halliday CL, Carter DA. Clonal reproduction and limited dispersal in an environmental population of *Cryptococcus neoformans* var. *gattii* isolates from Australia. *J Clin Microbiol*. 2003;41:703–11.
27. Swofford DL. PAUP* 4.06.10: Phylogenetic analysis using parsimony. Sunderland (MA): Sinauer Associates; 2003.
28. Nei M, Li WH. Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proc Natl Acad Sci U S A*. 1979;76:5269–73.
29. Wetton JH, Carter RE, Parkin DT, Walters D. Demographic study of a wild house sparrow population by DNA fingerprinting. *Nature*. 1987;327:147–9.
30. Rainer J, de Hoog GS, Wedde M, Graser Y, Gilges S. Molecular variability of *Pseudallescheria boydii*, a neurotropic opportunist. *J Clin Microbiol*. 2000;38:3267–73.
31. de Hoog GS, Marvin-Sikkema FD, Lahpoor GA, Gottschall JC, Prins RA, Gueho E. Ecology and physiology of *Pseudallescheria boydii*, an emerging opportunistic fungus. *Mycoses*. 1994;37:71–8.
32. Gueho E, de Hoog GS. Taxonomy of the medical species of *Pseudallescheria* and *Scedosporium*. *J Mycol Méd*. 1991;1:3–9.
33. Rainer J, de Hoog GS. Molecular taxonomy and ecology of *Pseudallescheria*, *Petriella* and *Scedosporium prolificans* (*Microascales*) containing opportunistic agents on humans. *Mycol Res*. 2006;110:151–60.
34. Bounoux ME, Tavanti A, Bouchier C, Gow NA, Magnier A, Davidson AD, et al. Collaborative consensus for optimized multilocus sequence typing of *Candida albicans*. *J Clin Microbiol*. 2003;41:5265–6.
35. Wedde M, Muller D, Tintelnot K, de Hoog GS, Stahl U. PCR-based identification of clinically relevant *Pseudallescheria/Scedosporium* strains. *Med Mycol*. 1998;36:61–7.
36. Alvarez M, Lopez Ponga B, Rayon C, Garcia Gala J, Roson Porto MC, Gonzalez M, et al. Nosocomial outbreak caused by *Scedosporium prolificans* (*inflatum*): four fatal cases in leukemic patients. *J Clin Microbiol*. 1995;33:3290–5.
37. Kidd SE, Hagen F, Tschärke RL, Huynh M, Bartlett KH, Fyfe M, et al. A rare genotype of *Cryptococcus gattii* caused the cryptococcosis outbreak on Vancouver Island (British Columbia, Canada). *Proc Natl Acad Sci U S A*. 2004;101:17258–63.
38. Tapia M, Richard C, Baro J, Salesa R, Figols J, Zurbano F, et al. *Scedosporium inflatum* infection in immunocompromised hematological patients. *Br J Haematol*. 1994;87:212–4.
39. Idigoras P, Garcia-Arenzana JM, Saenz JR, Pineiro L, Marin J. Isolation of *Scedosporium prolificans* from the air in the room of a patient with leukemia and disseminated infection with this fungus. *Enferm Infecc Microbiol Clin*. 2000;18:426–7.
40. Salkin IF, McGinnis MR, Dykstra MJ, Rinaldi MG. *Scedosporium inflatum*, an emerging pathogen. *J Clin Microbiol*. 1988;26:498–503.

Address for correspondence: Wieland Meyer, Molecular Mycology Research Laboratory, Westmead Hospital, Centre for Infectious Diseases and Microbiology, Institute of Clinical Pathology and Medical Research, Level 3, Room 3114A, Darcy Rd, Westmead, New South Wales 2145, Australia; email: w.meyer@usyd.edu.au

The opinions expressed by authors contributing to this journal do not necessarily reflect the opinions of the Centers for Disease Control and Prevention or the institutions with which the authors are affiliated.

EMERGING INFECTIOUS DISEASES *online*

www.cdc.gov/eid

To receive tables of contents of new issues send an email to listserv@cdc.gov with subscribe eid-toc in the body of your message.

Technical Appendix

Table. Strains used in molecular typing of Australian *Scedosporium* isolates*

Patient	WM no.	Age/Sex	Specimen type	Comorbidity/risk factor	Status	Building renovation	City/State	Date of isolation†	Species	Comments	Strain profile
1	WM 06.373	2/M	Sputum	Cystic fibrosis	C	No	Townsville, QLD	1999	<i>S. prolificans</i>		Strain 1a
	WM 06.374	2/M	Tracheal aspirate	Cystic fibrosis	C	No	Townsville, QLD	1999	<i>S. prolificans</i>		Strain 1b
	WM 06.375	2/M	BAL	Cystic fibrosis	C	No	Townsville, QLD	1999	<i>S. prolificans</i>		Strain 1a
	WM 06.372	4/M	BAL	Cystic fibrosis	C	No	Adelaide, SA	01/04/2001	<i>S. prolificans</i>		Strain 1a
2	WM 06.376	51/M	Tissue (chest)	Trauma	I	No	Canberra, ACT	2000	<i>S. prolificans</i>		
3	WM 06.377	45/F	Bronchial wash fluid	Solid organ malignancy	C	No	Brisbane, QLD	2000	<i>S. prolificans</i>		
4	WM 06.378	61/M	Tissue (cornea)	Trauma	I	Yes	Melbourne, VIC	2000	<i>S. prolificans</i>		
5	WM 06.379	77/M	Tissue (leg)	Trauma	I	Yes	Sydney, NSW	2000	<i>S. prolificans</i>		
6	WM 06.380	20/M	Synovial fluid	HSCT	I	Yes	Melbourne, VIC	09/2000	<i>S. prolificans</i>	Suspected Melbourne outbreak	
7	WM 06.381	27/F	Blood	Hematological malignancy	I	Yes	Perth, WA	06/1994	<i>S. prolificans</i>		
8	WM 06.383	43/M	CSF	Hematological malignancy	I	No	Perth, WA	07/1991	<i>S. prolificans</i>		
9	WM 06.384	28/M	Blood	Trauma	I	No	Perth, WA	08/1989	<i>S. prolificans</i>		
10	WM 06.385	29/M	Ear swab	None	C	No	Sydney, NSW	05/11/2001	<i>S. aurantiacum</i>		Strain 10a
	WM 06.386	29/M	Ear swab	None	C	No	Sydney, NSW	06/11/2001	<i>S. aurantiacum</i>		Strain 10b
11	WM 06.387	73/M	Nasal mucosa	Chronic sinusitis	C	No	Sydney, NSW	01/12/2001	<i>S. aurantiacum</i>		
12	WM 06.388	NA†	Tissue (corneal)	Trauma	I	No	Perth, WA	09/2001	<i>S. aurantiacum</i>		
13	WM 06.389	50/M	Sputum	None	C	No	Wauchope, NSW	09/2001	<i>S. apiospermum</i>		
14	WM 06.390	64/M	Skin	Diabetes mellitus	C	No	Perth, WA	03/2001	<i>S. aurantiacum</i>		
15	WM 06.391	42/F	Skin/nail	None	C	No	Sydney, NSW	09/2001	<i>S. prolificans</i>		
16	WM 06.392	46/F	Sputum	Cystic fibrosis	C	Yes	Melbourne, VIC	10/2001	<i>S. prolificans</i>	Suspected Melbourne outbreak	
17	WM 06.393	23/M	Synovial fluid	HSCT	I	Yes	Melbourne, VIC	09/2000	<i>S. prolificans</i>		Suspected Melbourne outbreak
18	WM 06.394	38/F	Nasal fluid	None	C	Yes	Melbourne, VIC	01/2002	<i>S. prolificans</i>		
19	WM 06.395	40/M	Tissue (nasal mucosa)	Hematological malignancy	C	Yes	Melbourne, VIC	10/2001	<i>S. prolificans</i>	Suspected Melbourne outbreak	
20	WM 06.396	25/M	Blood	Hematological malignancy	I	Yes	Melbourne, VIC	05/2002	<i>S. prolificans</i>		

Patient	WM no.	Age/Sex	Specimen type	Comorbidity/risk factor	Status	Building renovation	City/State	Date of isolation†	Species	Comments	Strain profile
21	WM 06.397	85/M	Sputum	Hematological malignancy	C	Yes	Melbourne, VIC	05/2002	<i>S. prolificans</i>		
22	WM 06.398	44/F	Blood	Hematological malignancy	I	Yes	Melbourne, VIC	05/2002	<i>S. prolificans</i>		
23	WM 06.399	28/F	Blood	Hematological malignancy	I	Yes	Melbourne, VIC	11/2001	<i>S. prolificans</i>	Suspected Melbourne outbreak	
24	WM 06.400	33/M	Blood	HSCT	I	Yes	Melbourne, VIC	10/2001	<i>S. prolificans</i>	Suspected Melbourne outbreak	
25	WM 06.401	57/F	Intervertebral disk	HSCT	I	Yes	Melbourne, VIC	10/2001	<i>S. prolificans</i>	Suspected Melbourne outbreak	
26	WM 06.402	49/F	Synovial fluid	Trauma	I	Yes	Melbourne, VIC	09/2000	<i>S. prolificans</i>	Suspected Melbourne outbreak	
27	WM 06.403	66/F	Skin	None	C	No	Sydney, NSW	09/2001	<i>S. prolificans</i>		Strain 27a
	WM 06.404	66/F	Skin	None	C	No	Sydney, NSW	09/2001	<i>S. prolificans</i>		Strain 27b
28	WM 06.405	47/M	BAL	Lung transplantation	I	Yes	Melbourne, VIC	10/2001	<i>S. prolificans</i>	Suspected Melbourne outbreak	
29	WM 06.406	67/F	Blood	Hematological malignancy	I	No	Perth, WA	03/1992	<i>S. prolificans</i>		
30	WM 06.407	43/M	Cerebrospinal fluid	Hematological malignancy	I	No	Perth, WA	07/1991	<i>S. prolificans</i>		
31	WM 06.408	67/F	Blood	Hematological malignancy	I	No	Perth, WA	03/1992	<i>S. prolificans</i>		
32	WM 06.409	54/F	Blood	Hematological malignancy	I	NA	Perth, WA	09/2001	<i>S. prolificans</i>		
33	WM 06.410	NA	Ear swab	None	C	No	Brisbane, QLD	04/2001	<i>S. prolificans</i>		
34	WM 06.412	NA	Bronchial wash fluid	None	C	No	Sydney, NSW	04/2001	<i>S. prolificans</i>		
35	WM 06.413	NA	Bronchial wash fluid washings	NA	C	No	Brisbane, QLD	04/2001	<i>S. prolificans</i>		
36	WM 06.414	41/M	Bronchial wash fluid	None	C	No	Adelaide, SA	1999	<i>S. prolificans</i>		
37	WM 06.415	NA	Tissue (nasal mucoa)	None	C	No	Brisbane, QLD	04/2001	<i>S. prolificans</i>		
38	WM 06.416	58/M	Blood	Other immuno-compromise	I	No	Melbourne, VIC	2000	<i>S. prolificans</i>		
39	WM 06.417	NA	Tissue (knee)	Recent surgery	C	No	Brisbane, QLD	2000	<i>S. prolificans</i>		
40	WM 06.418	NA	Sputum	None	C	No	Sydney, NSW	04/2001	<i>S. prolificans</i>		
41	WM 06.420	56/F	Skin	None	I	No	Perth, WA	08/1991	<i>S. prolificans</i>		
42	WM 06.424	32/M	Ear swab	None	C	No	Sydney, NSW	14/04/2003	<i>S. apiospermum</i>		
43	WM 06.425	20/F	Sputum	Cystic fibrosis	C	No	Sydney, NSW	04/03/2003	<i>S. aurantiacum</i>		
44	WM 06.426	63/M	Bronchial wash fluid	Lung transplantation	I	No	Sydney, NSW	10/06/2003	<i>S. prolificans</i>		

Patient	WM no.	Age/Sex	Specimen type	Comorbidity/risk factor	Status	Building renovation	City/State	Date of isolation†	Species	Comments	Strain profile
45	WM 06.427	28/M	Sputum	Chronic lung disease	C	No	Sydney, NSW	10/06/2003	<i>S. aurantiacum</i>		
46	WM 06.428	59/F	Tissue (sphenoid sinus)	Chronic sinusitis	C	Yes	Sydney, NSW	23/06/2003	<i>S. apiospermum</i>		
47	WM 06.429	NA	Skin	None	C	No	Sydney, NSW	21/07/2003	<i>S. apiospermum</i>		
48	WM 06.430	66/M	Skin	None	I	No	Sydney, NSW	04/06/2003	<i>S. apiospermum</i>		
49	WM 06.431	19/F	Skin	None	C	No	Sydney, NSW	21/07/2003	<i>S. apiospermum</i>		
50	WM 06.432	32/M	Blood	Hematological malignancy	I	Yes	Sydney, NSW	22/09/2003	<i>S. prolificans</i>	Suspected Sydney outbreak	
51	WM 06.433	56/M	Sputum	Hematological malignancy	I	No	Sydney, NSW	22/09/2003	<i>S. prolificans</i>		
52	WM 06.434	67/M	Blood	Hematological malignancy	I	Yes	Sydney, NSW	28/11/2003	<i>S. prolificans</i>	Suspected Sydney outbreak	
53	WM 06.435	74/M	BAL	Recent surgery	C	Yes	Sydney, NSW	26/11/2003	<i>S. apiospermum</i>		
54	WM 06.436	57/M	Blood	Hematological malignancy	I	Yes	Wollongong, NSW	01/10/2003	<i>S. prolificans</i>		
55	WM 06.437	44/F	BAL	Cystic fibrosis	C	No	Sydney, NSW	15/12/2003	<i>S. apiospermum</i>		
56	WM 06.438	58/F	Sputum	Lung transplantation	C	No	Sydney, NSW	01/12/2003	<i>S. prolificans</i>		
57	WM 06.439	64/M	Blood	Hematological malignancy	I	No	Sydney, NSW	26/12/2003	<i>S. prolificans</i>		Strain 57a
	WM 06.440	64/M	Bronchial wash fluid	Hematological malignancy	I	No	Sydney, NSW	26/12/2003	<i>S. prolificans</i>		Strain 57b
	WM 06.441	64/M	Skin	Hematological malignancy	I	No	Sydney, NSW	26/12/2003	<i>S. prolificans</i>		Strain 57a
58	WM 06.442	58/M	Sputum	Hematological malignancy	I	Yes	Sydney, NSW	02/01/2004	<i>S. prolificans</i>		
59	WM 06.443	61/F	Sputum	None	C	No	Sydney, NSW	10/05/2003	<i>S. apiospermum</i>		
60	WM 06.444	80/M	Sputum	None	C	No	Sydney, NSW	28/03/2003	<i>S. aurantiacum</i>		
61	WM 06.445	62/M	Sputum	Hematological malignancy	I	No	Sydney, NSW	25/11/2003	<i>S. prolificans</i>		
62	WM 06.446	93/M	Tissue (ear)	Solid tumor malignancy, diabetes mellitus	I	No	Sydney, NSW	10/03/2003	<i>S. aurantiacum</i>		
63	WM 06.447	3/F	Tissue (brain)	None	I	Yes	Sydney, NSW	01/09/2003	<i>S. prolificans</i>		
64	WM 06.448	60/F	Bronchial washings	Chronic renal disease	C	Yes	Sydney, NSW	02/07/2003	<i>S. prolificans</i>		
65	WM 06.449	10/F	Ear swab	None	C	No	Sydney, NSW	11/07/2003	<i>S. prolificans</i>		
66	WM 06.450	58/F	Bronchial wash fluid	Chronic lung disease	C	Yes	Sydney, NSW	12/05/2003	<i>S. apiospermum</i>		
67	WM 06.451	62/F	Nose swab	None	C	Yes	Sydney, NSW	08/12/2003	<i>S. prolificans</i>		

Patient	WM no.	Age/Sex	Specimen type	Comorbidity/risk factor	Status	Building renovation	City/State	Date of isolation†	Species	Comments	Strain profile
68	WM 06.452	60/F	Sputum	None	C	Yes	Sydney, NSW	04/11/2003	<i>S. prolificans</i>		
69	WM 06.453	13/F	BAL	NA	C	No	Sydney, NSW	01/09/2003	<i>S. prolificans</i>		
70	WM 06.454	11/F	Sputum	NA	C	No	Sydney, NSW	01/09/2003	<i>S. aurantiacum</i>		
71	WM 06.455	57/M	Bronchial wash fluid	Lung transplantation	C	No	Sydney, NSW	19/12/2003	<i>S. apiospermum</i>		
72	WM 06.456	NA	Ear swab	NA	C	No	Sydney, NSW	08/01/2004	<i>S. apiospermum</i>		
73	WM 06.457	59/F	Blood	Hematological malignancy	I	Yes	Sydney, NSW	24/01/2004	<i>S. prolificans</i>	Suspected Sydney outbreak	
	WM 06.458	59/F	Sputum	Hematology malignancy	I	Yes	Sydney, NSW	24/01/2004	<i>S. prolificans</i>		Suspected Sydney outbreak
74	WM 06.459	80/F	Ear swab	None	C	No	Sydney, NSW	28/01/2004	<i>S. aurantiacum</i>		
75	WM 04.468	61/M	Bronchial wash fluid	Solid tumor malignancy	C	No	Melbourne, VIC	12/01/2004	<i>S. prolificans</i>		
76	WM 04.497	71/F	Tissue (sphenoid sinus)	Chronic sinusitis	C	No	Sydney, NSW	24/02/2004	<i>S. aurantiacum</i>		
77	WM 06.462	30/F	Sputum	Cystic fibrosis	C	No	Sydney, NSW	16/02/2004	<i>S. aurantiacum</i>		
78	WM 06.463	55/F	Bronchial wash fluid	Recent surgery	C	No	Sydney, NSW	04/03/2004	<i>S. prolificans</i>		
79	WM 06.464	63/M	Tissue (eye)	Recent eye surgery	I	Yes	Sydney, NSW	01/03/2004	<i>S. prolificans</i>		
80	WM 06.465	29/F	Sputum	Diabetes mellitus	I	No	Sydney, NSW	15/04/2004	<i>S. aurantiacum</i>		Strain 80a
	WM 06.466	29/F	Tissue (bone)	Diabetes mellitus	I	No	Sydney, NSW	07/07/2004	<i>S. aurantiacum</i>		Strain 80a
	WM 06.467	29/F	Tissue (bone)	Diabetes mellitus	I	No	Sydney, NSW	12/08/2004	<i>S. aurantiacum</i>		Strain 80a
	WM 06.468	29/F	Wound fluid	Diabetes mellitus	I	No	Sydney, NSW	19/10/2004	<i>S. aurantiacum</i>		Strain 80b
81	WM 06.469	66/M	Ear swab	Diabetes mellitus	I	No	Sydney, NSW	19/04/2004	<i>S. apiospermum</i>		
82	WM 06.470	37/F	Sputum	Other immuno-compromise	C	No	Sydney, NSW	02/03/2004	<i>S. prolificans</i>		
83	WM 06.471	43/M	BAL	Lung transplantation	I	No	Sydney, NSW	15/05/2004	<i>S. apiospermum</i>		Strain 83a
	WM 06.472	43/M	Bronchial wash fluid	Lung transplantation	I	No	Sydney, NSW	13/05/2004	<i>S. apiospermum</i>		Strain 83a
	WM 06.473	43/M	BAL	Lung transplantation	I	No	Sydney, NSW	04/08/2004	<i>S. prolificans</i>		
	WM 06.474	43/M	BAL	Lung transplantation	I	No	Sydney, NSW	04/08/2004	<i>S. apiospermum</i>		Strain 83b
	WM 06.475	43/M	BAL	Lung transplantation	I	No	Sydney, NSW	05/08/2004	<i>S. apiospermum</i>		Strain 83b

Patient	WM no.	Age/Sex	Specimen type	Comorbidity/risk factor	Status	Building renovation	City/State	Date of isolation†	Species	Comments	Strain profile
84	WM 06.476	18/M	Sputum	Chronic lung disease	C	No	Sydney, NSW	08/03/2004	<i>S. aurantiacum</i>		
85	WM 06.477	NA	Ear swab	None	C	No	Sydney, NSW	09/05/2004	<i>S. apiospermum</i>		
86	WM 06.478	NA	Blood	NA	I	No	Sydney, NSW	03/06/2004	<i>S. prolificans</i>		
87	WM 06.479	18/F	Sputum	Cystic fibrosis	C	No	Sydney, NSW	09/07/2004	<i>S. aurantiacum</i>		
	WM 06.480	18/F	Sputum	Cystic fibrosis	C	No	Sydney, NSW	18/10/2004	<i>S. aurantiacum</i>		
	WM 06.481	18/F	Sputum	Cystic fibrosis	C	No	Sydney, NSW	18/11/2004	<i>S. aurantiacum</i>		
88	WM 06.482	24/M	Lung	None	I	No	Sydney, NSW	01/06/2004	<i>S. aurantiacum</i>		
89	WM 06.483	NA	Skin	None	C	No	Sydney, NSW	04/08/2004	<i>S. aurantiacum</i>		
90	WM 06.484	36/F	Sputum	Lung transplantation	C	No	Sydney, NSW	03/08/2004	<i>S. aurantiacum</i>		
91	WM 06.485	67/M	Sputum	Chronic lung disease	C	No	Sydney, NSW	23/08/2004	<i>S. prolificans</i>		
	WM 06.486	67/M	Sputum	Chronic lung disease	C	No	Sydney, NSW	23/08/2004	<i>S. apiospermum</i>		
92	WM 06.488	42/F	skin	None	C	No	Sydney, NSW	07/06/2004	<i>S. apiospermum</i>		
93	WM 06.489	43/M	Ear swab	None	C	No	Sydney, NSW	07/06/2004	<i>S. apiospermum</i>		
94	WM 06.490	53/F	Ear swab	Diabetes mellitus	C	No	Sydney, NSW	04/06/2004	<i>S. apiospermum</i>		
95	WM 06.491	71/M	Ear swab	None	C	No	Sydney, NSW	04/06/2004	<i>S. apiospermum</i>		
96	WM 06.492	70/F	Sputum	Chronic lung disease	C	No	Sydney, NSW	17/06/2004	<i>S. aurantiacum</i>		
97	WM 06.493	5/M	Ear swab	None	C	No	Sydney, NSW	17/06/2004	<i>S. aurantiacum</i>		
98	WM 06.494	26/F	Ear swab	None	C	No	Sydney, NSW	10/06/2004	<i>S. apiospermum</i>		
99	WM 06.495	76/F	Tissue (maxillary sinus)	None	C	No	Sydney, NSW	17/06/2004	<i>S. aurantiacum</i>		Strain 99a
	WM 06.496	76/F	Tissue (maxillary sinus)	None	C	No	Sydney, NSW	17/06/2004	<i>S. aurantiacum</i>		Strain 99b
100	WM 06.498	59/F	Ear swab	None	C	No	Sydney, NSW	20/07/2004	<i>S. aurantiacum</i>		
101	WM 06.499	15/M	Sputum	Cystic fibrosis	C	No	Sydney, NSW	15/07/2004	<i>S. prolificans</i>		
102	WM 06.500	62/F	Sputum	Chronic lung disease	C	No	Sydney, NSW	22/07/2004	<i>S. apiospermum</i>		
	WM 06.501	62/F	Sputum	Chronic lung disease	C	No	Sydney, NSW	22/07/2004	<i>S. prolificans</i>		
103	WM 06.502	34/M	Tissue (ethmoid sinus)	Chronic sinusitis	I	No	Sydney, NSW	16/08/2004	<i>S. prolificans</i>		
	WM 06.503	34/M	Tissue (ethmoid sinus)	Chronic sinusitis	I	No	Sydney, NSW	31/08/2004	<i>S. prolificans</i>		

Patient	WM no.	Age/Sex	Specimen type	Comorbidity/risk factor	Status	Building renovation	City/State	Date of isolation†	Species	Comments	Strain profile
104	WM 06.504	55/F	Ear swab	None	C	No	Sydney, NSW	27/08/2004	<i>S. apiospermum</i>	White colony	
	WM 06.505	55/F	Ear swab	None	C	No	Sydney, NSW	27/08/2004	<i>S. apiospermum</i>	Grey colony	
105	WM 06.506	56/M	Nasal swab	None	C	No	Sydney, NSW	03/09/2004	<i>S. prolificans</i>		
106	WM 06.507	30/M	Ear swab	None	C	No	Sydney, NSW	03/09/2004	<i>S. prolificans</i>		
107	WM 06.508	8/M	Ear swab	None	C	No	Sydney, NSW	01/09/2004	<i>S. prolificans</i>		
108	WM 06.509	28/M	Ear swab	None	C	No	Sydney, NSW	25/08/2004	<i>S. apiospermum</i>		
109	WM 06.511	82/F	Sputum	None	C	No	Sydney, NSW	13/08/2004	<i>S. aurantiacum</i>		
110	WM 06.512	7/M	Ear swab	None	I	No	Sydney, NSW	06/09/2004	<i>S. prolificans</i>		
111	WM 06.513	45/M	BAL	Lung transplantation	I	No	Melbourne, VIC	19/12/2003	<i>S. prolificans</i>		
112	WM 06.514	47/M	BAL	Lung transplantation	C	No	Melbourne, VIC	28/01/2004	<i>S. prolificans</i>		
113	WM 06.515	44/M	Blood	Hematological malignancy	I	No	Melbourne	09/03/2004	<i>S. prolificans</i>		
114	WM 06.516	61/M	BAL	Lung transplantation	C	No	Melbourne, VIC	27/04/2004	<i>S. prolificans</i>		
115	WM 06.517	53/F	Sputum	Cystic fibrosis	C	No	Melbourne, VIC	08/09/2004	<i>S. prolificans</i>		
116	WM 06.518	64/F	Sputum	Lung transplantation	C	No	Melbourne, VIC	12/11/2003	<i>S. prolificans</i>		
117	WM 06.519	66/M	BAL	Lung transplantation	C	No	Melbourne, VIC	29/09/2004	<i>S. prolificans</i>		
118	WM 06.520	62/M	Tissue (chest)	Lung transplantation	I	No	Sydney, NSW	15/10/2004	<i>S. apiospermum</i>		Strain 118a
	WM 06.521	62/M	Pleural fluid	Lung transplantation	I	No	Sydney, NSW	14/10/2004	<i>S. apiospermum</i>		Strain 118a
	WM 06.522	62/M	Tissue (bone)	Lung transplantation	I	No	Sydney, NSW	17/10/2004	<i>S. apiospermum</i>		Strain 118a
	WM 06.523	62/M	Wound fluid	Lung transplantation	I	No	Sydney, NSW	02/11/2004	<i>S. apiospermum</i>		Strain 118b
	WM 06.524	62/M	Tissue (bone)	Lung transplantation	I	No	Sydney, NSW	09/11/2004	<i>S. apiospermum</i>		Strain 118b
119	WM 06.525	35/M	Blood	Hematological malignancy	I	No	Perth, WA	01/12/2003	<i>S. prolificans</i>		
	WM 06.526	35/M	Skin	Hematological malignancy	I	No	Perth, WA	27/11/2003	<i>S. prolificans</i>		

*C, colonization; QLD, Queensland; BAL, bronchoalveolar lavage; SA, South Australia; I, infection; ACT, Australian Capital Territory; VIC, Victoria; NSW, New South Wales; HSCT, hematopoietic stem cell transplant; WA, Western Australia; NA, not available.

†Isolation dates are given where known; otherwise, only month, year, or both is stated.