

Phylogeny of European Bat Lyssavirus 1 in *Eptesicus isabellinus* Bats, Spain

Sonia Vázquez-Morón, Javier Juste,
Carlos Ibáñez, José M. Berciano,
and Juan E. Echevarría

To better understand the epidemiology of European bat lyssavirus 1 (EBLV-1) in Europe, we phylogenetically characterized *Lyssavirus* from *Eptesicus isabellinus* bats in Spain. An independent cluster of EBLV-1 possibly resulted from geographic isolation and association with a different reservoir from other European strains. EBLV-1 phylogeny is complex and probably associated with host evolutionary history.

The genus *Lyssavirus* comprises 3 species that can infect bats in Europe: *European bat lyssavirus 1* (EBLV-1), *European bat lyssavirus 2*, and *West-Caucasian bat virus* (1,2). Most lyssavirus-infected bats have been found in north-central Europe (Germany, the Netherlands, Denmark, Poland, and France); of these, >95% were serotine bats (*Eptesicus serotinus*) infected by EBLV-1 (3–5). EBLV-1 in other bat species has rarely been described (3,6). EBLV-1-infected bats become increasingly scarce from north to south in Europe, and no cases in northern Spain or Italy have been reported. The same trend has been consistently found within Germany (3) except for an artifact that arose from varied surveillance intensity among different countries. However, several infected serotine bats in southern Spain have been reported (7). These bats have been assigned to the species *E. isabellinus*, which has closely related populations on the African side of the Gibraltar Strait (8). This species is strongly divergent from *E. serotinus* bats (>16% of cytochrome b gene) in the northern Iberian Peninsula (9). In Spain, the distribution of EBLV-1 cases

in bats apparently coincides with the distribution of *E. isabellinus* bats; 10 cases of human exposure after contact with infected bats have been reported; each was associated with *E. isabellinus* bats.

Two subtypes have been proposed for EBLV-1: EBLV-1a, which extends from the Netherlands to Russia in a west–east axis, and EBLV-1b, which includes strains that extend south through France and the Netherlands and the only 2 published strains from Iberia (1). We phylogenetically characterized EBLV-1 strains associated with *E. isabellinus* bats, a reservoir in the Iberian Peninsula that differs from *E. serotinus* bats.

The Study

We sequenced 12 bat brains positive for *Lyssavirus* antigen detected by immunofluorescence and reverse transcription–PCR (RT-PCR) as described (10). All viruses were identified as EBLV-1. For phylogenetic analyses, the 400-bp 5' variable extreme of the nucleoprotein gene of these EBLV-1 strains was amplified by specific EBLV-1 nested RT-PCR and sequenced by using the following primers: SEQVAR1F 5'-₁ACGCTTAACAACCAGATCAAAG₂₂-3', SEQVAR2F 5'-₅₁AAAAATGTAACACYYCTACA₇₀-3', EBLVSEQVAR1R 5'-₅₉₆CAGTCTCAAAGATCTGTTC AT₅₇₅-3', and EBLVSEQVAR2R 5'-₅₅₂TAGTCCCAGT ATTCTGTCC₅₃₃-3'.

All rabies-positive serotine bats came from southern Spain (Huelva, Seville, Murcia, and Badajoz) and were molecularly identified as *E. isabellinus* (8). An alignment was performed by using ClustalX (www.clustal.org) to combine the obtained sequences and other available EBLV-1 sequences from GenBank, including a Duvenhage virus used as the outgroup (online Appendix Table, www.cdc.gov/EID/content/17/3/520-appT.htm). Before conducting further analyses, we used jModel Test (<http://darwin.uvigo.es/software/jmodeltest.html>) to select the best fitting substitution model for our sequences according to the corrected Akaike information criterion. Maximum-likelihood phylogenies were reconstructed by using PHYML (<http://atgc.lirmm.fr/phyml>) software and by using a generalized time-reversible model and the γ parameter estimated in the analyses. Maximum-parsimony analyses were conducted by using PAUP* 4.0b10 (<http://paup.csit.fsu.edu/>) weighting transversions 15 \times according to the transitions/transversion ratio estimated in the jModelTest analyses. Confidence in the topologies for the maximum-likelihood and the maximum-parsimony analyses was established with 1,000 bootstrap replicates. A Bayesian phylogenetic inference was obtained by using MrBayes version 3.1 (<http://mrbayes.csit.fsu.edu/>) with random starting trees without constraints. Two simultaneous runs of 10⁷ generations were conducted, each with 4 Markov chains, and the trees were sampled

Author affiliations: Instituto de Salud Carlos III, Majadahonda, Madrid, Spain (S. Vázquez-Morón, J.M. Berciano, J.E. Echevarría); Centro de Investigación Biomédica de Epidemiología y Salud Pública, Barcelona, Spain (S. Vázquez-Morón, J.E. Echevarría); and Consejo Superior de Investigaciones Científicas Estación Biológica de Doñana, Seville, Spain (J. Juste, C. Ibáñez)

DOI: 10.3201/eid1703100894




Figure 1. European bat lyssavirus 1 (EBLV-1) phylogenetic reconstruction based on the first 400 bp of the nucleoprotein gene. The tree was obtained by Bayesian inference run for 10^7 generations; trees were sampled every 100 generations. The first 25% of trees were excluded from the analysis as burn-in. Black numbers indicate posterior probabilities. Bootstrap supports after 1,000 replicates for each node are also shown for maximum-parsimony (green numbers) and maximum-likelihood (blue numbers) analyses. Net *p*-distance values (as percentages) between groups are indicated by arrows. A parsimony-based network is presented for each major lineage; sizes of yellow circles are proportional to the number of individuals sharing a given haplotype, and reconstructed haplotypes (median vectors) are shown in red. DUVV, Duvnag virus.

every 100 generations. Net *p*-distances between groups were calculated by using MEGA4 (www.megasoftware.net/) (Figure 1).

The genetic structure and relationships between haplotypes were examined within the main lineages through a parsimony-based network built with a median-joining algorithm implemented in the Network 4.5.1

program (11). To evaluate and compare genetic variability and polymorphism among lineages, we estimated the number of haplotypes, mutations, and segregating sites as well as haplotype diversity and nucleotide diversity by using DNAsp version 4.5 (12) for the major clades (Table). Finally, to investigate population dynamics across lineages, the *F_s* and Tajima D statistics were calculated (Table).

Table. Genetic diversity statistics for EBLV-1*

Population	n	S	Eta	Hap	Hd	VarHd	Pi	ThetaNuc	k	Tajima D	Fu Fs
EBLV-1a	52	45	48	26	0.836	0.00267	0.00664	0.02656	2.6546	-2.5693 (0.00000)	-21.676 (0.00000)
EBLV-1b	25	35	35	18	0.970	0.00038	0.02202	0.02317	8.8067	-0.1885 (0.48000)	-4.555 (0.05100)
EBLV-1Spain	13	9	9	7	0.795	0.01191	0.00538	0.00725	2.1538	-1.0138 (0.18100)	-2.067 (0.06143)

*EBLV, European bat lyssavirus; n, no. sequences; S, no. segregating sites; Eta, no. mutations; Hap, no. haplotypes; Hd, haplotype diversity; VarHd, haplotype variance; Pi, nucleotide diversity; ThetaNuc, estimated population mutation rate per site; k, average no. nucleotide differences; and neutrality tests (Tajima D and Fu Fs).




Figure 2. Geographic distribution of *Eptesicus serotinus* bats (red), *E. isabellinus* bats (blue), and cases of rabies in bats (dots), Europe, 1990–2009. Obtained from Rabies Bulletin Europe (www.who-rabies-bulletin.org/).

These 2 statistics are considered to be the most powerful tests for detecting expansion events (13).

Conclusions

All phylogenetic analyses, regardless of the reconstruction criterion used, formed a monophyletic cluster of the EBLV-1 strains from Spain (only the Bayesian inference reconstruction is shown). The Bayesian inference, maximum-likelihood, and maximum-parsimony analyses identified the cluster from Spain and EBLV-1a and EBLV-1b as being monophyletic (Figure 1), although only maximum-likelihood and maximum-parsimony analyses suggested a closer relationship between EBLV-1b and the cluster from Spain. The genetic differentiation of the EBLV-1 strains from the Iberian Peninsula matches their association with another bat species (Figure 2), which suggests that the host bat's evolutionary history plays a major role in EBLV-1 molecular epidemiology, as has been proposed for rabies virus in bats in North America (14).

The low genetic diversity and the F_u F_s and Tajima D statistics (Table) all suggest rapid population expansion of EBLV-1a, which is consistent with the star-like structure of the network for this lineage (Figure 1). Conversely, haplotype and nucleotide diversity descriptors (Table) have the highest values for EBLV-1b and a complex network structure with differentiated subnetworks. All these elements indicate that this lineage has a complex evolutionary history. The lineage from Spain also has low diversity and a star-shaped network, but neutral evolution cannot be rejected on the basis of the F_s and D statistics. Net distances are similar within and between lineages,

except for EBLV-1a, which is slightly more differentiated (Figure 1). Consequently, the suggested EBLV-1 expansion from Spain into Europe (15) is not supported by our results, which record the highest variability and most complex phylogenetic structure for France and the Netherlands (Figure 1). This complex structure suggests either a longer evolutionary history in these areas or a recent contact of distinct bat lineages in this zone.

The results of this study show that the strains from Spain do not belong to subtype 1b because of their association with a different reservoir (*E. isabellinus* bats). Moreover, what is currently considered to be EBLV-1b seems to include at least 4 lineages that are more genetically diverse and have a complex history. EBLV-1a, however, has low genetic diversity despite its extensive geographic distribution, suggesting a relatively recent and successful expansion of this lineage. These results call into question the current classification of EBLV-1 into 2 single subtypes. To provide a better understanding of EBLV-1 molecular epidemiology in Europe, additional studies that consider different genes should be conducted and the current classification should be revised accordingly.

Acknowledgments

We thank the Genomics Unit of the Instituto de Salud Carlos III for analyses of the genomic sequences and Enrique Royuela Casamayor for his involvement in the daily work.

This project was financially supported by an agreement between the Public Health Department of the Spanish Ministry of Health and the Instituto de Salud Carlos III for the development of "Rabies Surveillance in Spain" and by projects SAF 2006-12784-C02-01 and SAF 2006-12784-C02-02 of the General Research Programme of the Spanish Ministry of Science and Education.


Dr Vázquez-Morón is a PhD candidate at the Instituto de Salud Carlos III and Complutense University of Madrid. Her main research interests are the epidemiology and public health implications of rabies and emerging viruses in bats.

References

- Amengual B, Whitby JE, King A, Cobo JS, Bourhy H. Evolution of European bat lyssaviruses. *J Gen Virol*. 1997;78:2319–28.
- Botvinkin AD, Poleschuk EM, Kuzmin IV, Borisova TI, Gazaryan SV, Yager P, et al. Novel lyssaviruses isolated from bats in Russia. *Emerg Infect Dis*. 2003;9:1623–5.
- Müller T, Johnson N, Freuling CM, Fooks AR, Selhorst T, Vos A. Epidemiology of bat rabies in Germany. *Arch Virol*. 2007;152:273–88. DOI: 10.1007/s00705-006-0853-5
- Van der Poel WH, Van der Heide R, Verstraten ER, Takumi K, Lina PH, Kramps JA. European bat lyssaviruses, the Netherlands. *Emerg Infect Dis*. 2005;11:1854–9.
- Picard-Meyer E, Barrat J, Tissot E, Verdot A, Patron C, Barrat MJ, et al. Bat rabies surveillance in France, from 1989 through May 2005. *Dev Biol (Basel)*. 2006;125:283–8.

6. Serra-Cobo J, Amengual B, Abellan C, Bourhy H. European bat lyssavirus infection in Spanish bat populations. *Emerg Infect Dis.* 2002;8:413–20. DOI: 10.3201/eid0804.010263
7. Vázquez-Morón S, Juste J, Ibáñez C, Ruiz-Villamor E, Avellón A, Vera M, et al. Endemic circulation of European bat lyssavirus type 1 in serotine bats, Spain. *Emerg Infect Dis.* 2008;14:1263–6. DOI: 10.3201/1408.080068
8. Juste J, Bilgin R, Muñoz J, Ibáñez C. Mitochondrial DNA signatures at different spatial scales: from the effects of the Straits of Gibraltar to population structure in the meridional serotine bat (*Eptesicus isabellinus*). *Heredity.* 2009;103:178–87. DOI: 10.1038/hdy.2009.47
9. Ibáñez C, Garcia-Mudarra JL, Ruedi M, Stadelmann B. The Iberian contribution to cryptic diversity in European bats. *Acta Chiropterologica.* 2006;8:277–97. DOI: 10.3161/1733-5329(2006)8[277:TICTCD]2.0.CO;2
10. Vázquez-Morón S, Avellón A, Echevarría JE. RT-PCR for detection of all seven genotypes of *Lyssavirus* genus. *J Virol Methods.* 2006;135:281–7. DOI: 10.1016/j.jviromet.2006.03.008
11. Bandelt HJ, Forster P, Rohlf A. Median-joining networks for inferring intraspecific phylogenies. *Mol Biol Evol.* 1999;16:37–48.
12. Rozas J, Sánchez-DelBarrio JC, Messeguer X, Rozas R. Dna SP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics.* 2003;19:2496–7. DOI: 10.1093/bioinformatics/btg359
13. Ramos-Onsins SE, Rozas J. Statistical properties of new neutrality tests against population growth. *Mol Biol Evol.* 2002;19:2092–100.
14. Hughes GJ, Orciari LA, Rupprecht CE. Evolutionary timescale of rabies virus adaptation to North American bats inferred from the substitution rate of the nucleoprotein gene. *J Gen Virol.* 2005;86:1467–74. DOI: 10.1099/vir.0.80710-0
15. Davis PL, Holmes EC, Larrous F, Van der Poel WH, Tjornehoj K, Alonso WJ, et al. Phylogeography, population dynamics, and molecular evolution of European bat lyssaviruses. *J Virol.* 2005;79:10487–97. DOI: 10.1128/JVI.79.16.10487-10497.2005

Address for correspondence: Sonia Vázquez-Morón, Diagnostic Microbiology Service, National Center for Microbiology, Instituto de Salud Carlos III, Carretera de Majadahonda-Pozuelo Km 2, Majadahonda 28220, Madrid, Spain; email: svazquez@isciii.es



Search
past issues

EID
online
www.cdc.gov/eid