

Tuberculosis among Health Care Workers

Iacopo Baussano, Paul Nunn, Brian Williams, Emanuele Pivetta, Massimiliano Bugiani, and Fabio Scano

Medscape[®]CME[™] ACTIVITY

Medscape, LLC is pleased to provide online continuing medical education (CME) for this journal article, allowing clinicians the opportunity to earn CME credit.

This activity has been planned and implemented in accordance with the Essential Areas and policies of the Accreditation Council for Continuing Medical Education through the joint sponsorship of Medscape, LLC and Emerging Infectious Diseases. Medscape, LLC is accredited by the ACCME to provide continuing medical education for physicians.

Medscape, LLC designates this Journal-based CME activity for a maximum of 1 *AMA PRA Category 1 Credit(s)*[™]. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

All other clinicians completing this activity will be issued a certificate of participation. To participate in this journal CME activity: (1) review the learning objectives and author disclosures; (2) study the education content; (3) take the post-test and/or complete the evaluation at www.medscape.org/journal/eid; (4) view/print certificate..

Release date: February 25, 2011; Expiration date: February 25, 2012

Learning Objectives

Upon completion of this activity, participants will be able to:

- Describe the risk for health care workers (HCWs) of developing incident latent tuberculosis infection (LTBI)
- Describe the risk for HCWs of developing tuberculosis (TB) disease
- Describe TB infection control measures that may be effective in healthcare settings

Editor

Karen L. Foster, MA, Technical Writer/Editor, *Emerging Infectious Diseases*. *Disclosure: Karen L. Foster, MA, has disclosed no relevant financial relationships.*

CME Author

Laurie Barclay, MD, freelance writer and reviewer, Medscape, LLC. *Disclosure: Laurie Barclay, MD, has disclosed no relevant financial relationships.*

Authors

Disclosure: Iacopo Baussano, MD, MSc, PhD, SpID; Brian Williams, PhD; Emanuele Pivetta, MD; Massimiliano Bugiani, MD; and Fabio Scano, MD, have disclosed no relevant financial relationships. Paul Nunn, MA, FRCP(UK), has disclosed the following relevant financial relationship: served as a speaker or a member of a speakers bureau for Howard Hughes Foundation.

To assess the annual risk for latent tuberculosis infection (LTBI) among health care workers (HCWs), the incidence rate ratio for tuberculosis (TB) among HCWs worldwide, and the population-attributable fraction of TB to exposure of HCWs in their work settings, we reviewed the literature. Stratified pooled estimates for the LTBI rate

Author affiliations: University "A. Avogadro," Novara, Italy (I. Baussano); Centro per la Prevenzione Oncologica–Piemonte, Novara (I. Baussano); Imperial College, London, UK (I. Baussano); World Health Organization, Geneva, Switzerland (P. Nunn, B. Williams, F. Scano); University of Turin, Turin, Italy (E. Pivetta); Centro per la Prevenzione Oncologica–Piemonte, Turin (E. Pivetta); and Regional Reference Centre for Tuberculosis Prevention, Turin (M. Bugiani)

DOI: 10.3201/eid1703.100947

for countries with low (<50 cases/100,000 population), intermediate (50–100/100,000 population), and high (>100/100,000 population) TB incidence were 3.8% (95% confidence interval [CI] 3.0%–4.6%), 6.9% (95% CI 3.4%–10.3%), and 8.4% (95% CI 2.7%–14.0%), respectively. For TB, estimated incident rate ratios were 2.4 (95% CI 1.2–3.6), 2.4 (95% CI 1.0–3.8), and 3.7 (95% CI 2.9–4.5), respectively. Median estimated population-attributable fraction for TB was as high as 0.4%. HCWs are at higher than average risk for TB. Sound TB infection control measures should be implemented in all health care facilities with patients suspected of having infectious TB.

Transmission of tuberculosis (TB) in health care settings to both patients and health care workers (HCWs) has been reported from virtually every country of the world,

regardless of local TB incidence. TB transmission occurs through droplet nuclei aerosolized by patients with infectious pulmonary TB and inhaled by other persons. Transmission is most likely to occur from unrecognized or inappropriately treated TB. The risk for transmission varies by setting, occupational group, local prevalence of TB, patient population, and effectiveness of TB infection control measures (1).

In resource-rich countries TB began to be recognized as an occupational hazard in the 1950s (2); since then, effective infection control measures have been implemented to reduce the risk for nosocomial TB (3). Poor implementation of infection control measures, the effects of the HIV epidemic on TB, and the emergence of multidrug-resistant (MDR) strains resulted in the reemergence of hospital-based transmission of TB and MDR TB in the early 1990s (2).

Most recently, the emergence of extensively drug-resistant (XDR) TB strains is posing a major public health threat in contexts characterized by a lack of proper infection control measures (4). Although the risk for latent TB infection (LTBI) and TB disease has generally been considered to be higher among HCWs than in the general population, some studies from countries with low TB incidence and others from countries with high TB incidence failed to show a higher estimated risk for TB among HCWs than among the corresponding local community.

We had 2 main objectives in conducting our study. First, by reviewing the literature, we assessed the rate of LTBI and relative risk and risk difference for the incidence of TB among HCWs worldwide, compared with the incidence of TB in the corresponding local general population. Second, we assessed the population-attributable fraction (PAF) (percentage) of TB in the general population associated with exposure to HCWs in their working settings.

Methods

Literature Search for Incidence among HCWs

An initial search of the literature for systematic reviews or reviews of meta-analyses estimated LTBI and TB incidence among HCWs worldwide allowed us to identify 4 systematic reviews (1,5–7). Most of the studies analyzed by Menzies et al. in 1995 were performed in the United States, United Kingdom, or Canada (6). In 2005, Seidler et al. focused on the risk for TB in low-incidence areas (7). In 2006, Joshi et al. analyzed low- and middle-income countries (5), and in 2007 Menzies et al. expanded their search and analyses, accounting also for high-income countries (1). We identified, retrieved, and considered for inclusion in our study all studies reported in the above-mentioned systematic reviews. To integrate and update the search, we searched 3 electronic databases

for primary studies (i.e., PubMed, www.ncbi.nlm.nih.gov/pubmed; EMBASE, <http://embase.com>; and Web of Knowledge, www.isiwebofknowledge.com) to identify studies that assessed the incidence of LTBI and/or TB among HCWs published during January 2005–July 2010. The combination of terms adopted to perform the search is reported in the online Technical Appendix (www.cdc.gov/EID/content/17/3/488-Techapp.pdf). We restricted our search to articles published in English, French, Spanish, and Italian.

Study Selection

Studies were eligible for inclusion if they reported the incidence and the corresponding confidence intervals (CIs) of LTBI and TB disease among HCWs or if they reported the number of incident LTBI and TB cases identified in the study along with the overall number of HCWs investigated or the person-years of follow-up. LTBI incidence has been defined as tuberculin conversion after a documented negative-baseline tuberculin skin test (TST) as reported in the original study (3). We considered only studies adopting the intradermal TST technique. For cases of TB disease, we have included definitive (microbiologically confirmed) and presumptive (based on clinical, imaging, or pathology) diagnoses.

We excluded studies for a variety of reasons. Studies reporting only case series, only outbreak investigations, or only prevalence of LTBI and TB among HCWs were excluded, as were studies reporting investigations only of MDR TB and of only self-reported LTBI or TB. We also excluded studies that adopted interferon- γ release assay because of paucity of data and potential problems in pooling such data with those obtained from studies that adopted the TST technique. Furthermore, we excluded studies started before 1990 to decrease possible heterogeneity between studies associated with different awareness of TB as an occupational hazard for HCWs (2) and studies reporting data from studies investigating only HCWs working in outpatient clinics. All duplicate citations were eliminated from the initial database. Two of the authors (I.B., E.P.) screened these citations by reviewing titles and abstracts to identify potentially relevant studies and discussed disagreements until reaching a consensus. The database was then screened again to include only primary articles, and the full text of each citation was obtained and reviewed.

Data Extraction

Two of the authors (I.B., E.P.) designed a data extraction form. Then, all articles were independently reviewed and data extraction cross-checked.

The following data were collected from each study: country where the study was performed, study period, and health care setting under investigation. We also collected the

estimated incidence of LTBI and/or TB and corresponding CIs and/or number of incident LTBI and/or TB cases identified and the overall number of HCWs investigated or the person-years of follow-up. If reported, the incidence of LTBI and/or TB in a comparison group, such as the local general population or administrative workers not exposed to the health care setting under investigation, was also included. We used estimates provided by the World Health Organization (WHO) for the corresponding study period as estimates of TB incidence among the general populations in the host countries (8).

Data Collation and Meta-analysis

For each study, we considered the incidence for LTBI or TB among HCWs as reported in the original study. Adjusted estimates, when available, were preferred to crude estimates. Alternatively, we calculated the annual incidence of LTBI (hereafter referred to as annual risk for LTBI [ARTI]) and TB among HCWs. Then we estimated the incidence rate ratio (IRR), the risk difference of TB among HCWs compared with the corresponding general population, and the corresponding percentage of TB cases among HCWs attributable to exposure in health care settings. We used WHO estimates for the corresponding period (8) for TB incidence in the general population. Three studies could not be located and were excluded from the analysis.

We performed a random effects meta-analysis to account for the expected between-study variability for each study, i.e., we drew pooled estimates under the assumption that each study had different characteristics and measured different, though related, underlying yields (9). We used STATA version 11.0 software (StataCorp., College Station, TX, USA) for statistical analysis. The studies included in the meta-analysis were weighted by the inverse variance of their effect-size estimate (10).

To assess the fraction of TB in the population attributable to the exposure to health care settings, we calculated the PAF using Levin's formula (online Technical Appendix) (11). As a proxy for the proportion of the population working in health care settings, we considered the population of doctors, laboratory workers, nurses, and midwives as reported in WHO's Global Health Atlas (8). Community health workers were not included in this population because no study reported TB cases from these categories. We assessed heterogeneity across studies by the conventional χ^2 test for heterogeneity and by calculating the I^2 statistic, which accounts for the number of studies included in the meta-analysis and directly measures the variability not explained by information in the analysis (12).

To investigate possible sources of heterogeneity, we stratified the analysis accounting for TB incidence estimated

in the country where the study had been conducted. In particular, we defined 3 strata: countries with low TB incidence (i.e., <50 cases/100,000 persons), countries with intermediate TB incidence (i.e., 50–99 cases/100,000 persons), and countries with high TB incidence (i.e., \geq 100 cases/100,000 persons). Furthermore, to formally assess the effect of selected variables, such as the country income level as classified by the World Bank (13), the density of HCWs as reported by WHO (8), and the year the study was published, as possible sources of heterogeneity, we performed meta-regression analyses in which variance results from 2 additive components, 1 representing the variance within units, and the other the variance between units. Finally, we qualitatively assessed publication bias by drawing a funnel plot (online Technical Appendix Figure 1).

Results

We identified 846 potentially relevant unique citations from all literature searches and 74 from previous systematic reviews (Figure 1). Forty-three studies were eligible for inclusion (online Technical Appendix); these studies accounted for 1,176 cases of LTBI among HCWs and 49,576 person-years of follow-up and for 1,942 TB cases and >1,010,047 person-years of follow-up. Twenty-five and 18 studies reported LTBI and TB incident cases among HCWs, respectively (online Technical Appendix Tables 1, 2).

LTBI among HCWs

A median of 23 (interquartile range [IQR] 14–59) LTBI cases occurred among HCWs. These LTBI cases accounted for a median of 731 (IQR 111–2,144) person-years in each study.

The median estimated ARTI among HCWs was 2.9% (IQR 1.8%–8.2%) for studies from countries with low TB incidence, 8.7% (IQR 3.9%–10.5%) for studies from countries with intermediate TB incidence (in the present study only Brazil), and 7.2% (IQR 4.1%–14.3%) for studies from countries with high TB incidence. Consistently, the stratified pooled estimates for the ARTI were 3.8% (95% CI 3.0%–4.6%, $I^2 = 98\%$), 6.9% (95% CI 3.4%–10.3, $I^2 = 78\%$), and 8.4% (95% CI 2.7%–14.0%, $I^2 = 89\%$), respectively (Figure 2). Given that most studies in the meta-analysis were from countries with low TB incidence and that the variability of estimates was higher from countries with intermediate and high TB incidence, the overall (pooled) estimates of ARTI was 4.6% (95% CI 4.1%–5.6%, $I^2 = 97\%$).

TB among HCWs

A median of 34 (IQR 18–108) TB cases occurred among HCWs; each study accounted for a median of




Figure 1. Flow diagram for selection of studies of tuberculosis (TB) in health care workers (HCWs) published during January 2005–July 2010. INF, interferon.

12,689 (IQR 2,979–57,279) person-years. The median estimated annual incidence of TB among HCWs was 67 cases/100,000 persons (IQR 40–142), 91/100,000 persons (IQR 81–723), and 1,180/100,000 persons (IQR 91–3,222) for studies from countries with low, intermediate, and high TB incidence, respectively. The median estimates for TB incidence for the general population adopted as comparisons were 33/100,000 persons (IQR 27–37), 82/100,000 persons (IQR 58–223), and 311/100,000 persons (IQR 168–405), respectively. Median estimated annual TB IRRs were 2.0 (IQR 1.5–4.1), 1.4 (IQR 0.4–8.8), and 5.4 (IQR 1.7–9.1), respectively.

The median differences in TB incidence for countries with low, intermediate, and high TB incidence were 34 cases/100,000 persons (IQR 12–105), 33/100,000 persons (IQR 41 to 641), and 409/100,000 persons (IQR 166–2817), respectively. These findings suggest that 49%, 27%, and 81% of TB cases, respectively, among HCWs were attributable to exposure in health care settings.

For countries with low and intermediate TB incidence, stratified pooled estimates for annual TB IRRs were similar: 2.42 (95% CI 1.20–3.64, $I^2 = 85\%$) and 2.45 (95% CI 1.05–3.84, $I^2 = 98\%$), respectively (Figure 3). The TB IRRs for countries with high TB incidence was higher: 3.68 (95% CI 2.89–4.48, $I^2 = 96\%$). The overall estimate of annual TB IRR was 2.97 (95% CI 2.43–3.51, $I^2 = 98\%$).

For countries with low TB incidence, the proportion of HCWs in the population ranged from 1.7 to 8.2 HCWs per 1,000 residents. For countries with intermediate and high TB incidence, the proportion of HCWs in the population ranged from 3.5 to 12.8 HCWs per 1,000 residents and from 0.6 to 4.8 HCWs per 1,000 residents, respectively.

Using the estimated annual TB IRR and the reported population numbers of HCWs for each country of interest, we estimated the PAF. The median estimated PAF for TB was 0.36% (IQR 0.13–2.24) for studies from countries with low TB incidence, 0.43% (IQR 0.13–2.24) for studies from countries with intermediate TB incidence, and 0.38% (IQR 0.14–0.78) for studies from countries with high TB incidence.

For each level of IRR, a nonlinear relationship existed between the PAF and the percentage of HCWs in the population (online Technical Appendix Figure 2). For




Figure 2. Forrest plot showing study-specific and pooled estimates of the annual incidence of latent *Mycobacterium tuberculosis* infection (LTBI) among health care workers (HCWs), by tuberculosis incidence groups according to the World Health Organization. Studies are sorted by incidence category: low, <50 cases/100,000 population; intermediate, 50–99 cases/100,000 population; high, ≥100 cases/100,000 population. CI, confidence interval. A complete list of references is provided in the online Technical Appendix (www.cdc.gov/EID/content/16/3/488-Techapp.pdf).




Figure 3. Forrest plot showing the study-specific and pooled estimates of incidence rate ratios (RR) for tuberculosis (TB) among health care workers (HCWs) compared with the corresponding general populations, by TB incidence group according to the World Health Organization. Studies are sorted by incidence category: low, <50 cases/100,000 population; intermediate, 50–99 cases/100,000 population; high, ≥ 100 cases/100,000 population. CI, confidence interval. A complete list of references is provided in the online Technical Appendix (www.cdc.gov/EID/content/16/3/488-Techapp.pdf).

example, Harries et al. (14) and Rao et al. (15) reported similar IRRs (7.9 and 7.6, respectively), but because of the different proportion of HCWs working in health care settings in Malawi and India, respectively, the estimated PAF varied widely (0.42% and 1.21%, respectively). On the other hand, Gopinath et al. (16) and Rao et al. (15) reported from India; thus, the proportion of population working in health care settings was assumed to be the same (1.9 HCWs/1,000 persons, respectively) but the large difference in IRR reported in the 2 studies produced a substantial shift in the estimated PAF (from 0.04% to 1.2%).

The meta-regression model, which accounts for both the effect of country income levels as classified by the World Bank (13) and for the HCWs densities as reported by WHO (8), has shown that the between-studies variance was significantly reduced (τ^2 decreased from 1.2–0.71). In particular, the IRR for TB among HCWs decreased by $\approx 10\%$ as the HCW density increased by 1/1,000 persons and by about two thirds in middle-income countries, compared with low-income countries. However, the variability not explained by the information included in the analysis decreased to only 96%. Incorporating the effect of time (i.e., year of publication of the paper) in the analysis did not significantly decrease the heterogeneity between the studies.

Finally, visual inspection of the funnel plot suggested a possible publication bias for the studies reporting TB incidence among HCWs from countries with high TB incidence (online Technical Appendix). A formal testing

procedure that used the Begg test, however, failed to confirm such a bias (17).

Discussion

The results of our analysis show that the risk for TB among HCWs is consistently higher than the risk among the general population worldwide. This finding confirms that TB is an occupational disease. The introduction of TB transmission control measures, essential to protect HCWs, may decrease TB annual incidence among HCWs by as much as 49%, 27%, and 81% in countries with low, intermediate, and high TB incidence, respectively.

The annual risk for LTBI among HCWs and the TB IRRs, excluding the study by Dimitrova et al. (18), increase according to the background TB incidence in the population. The exceptionally low IRR estimated from the study by Dimitrova et al. may be a consequence of our decision to calculate TB IRR by pooling HCWs working in general health services and in TB facilities and considering as a reference TB incidence in the general population of Russia (95/100,000), which was higher than the incidence reported in the Samara Oblast (75/100,000), where the study was conducted. The reported incidence among HCWs working in general health services was 69/100,000, whereas the incidence among staff at the TB services was 741/100,000, which is 10 \times higher than among the general population.

The high PAF represents the effect of an increase in the HCW population in a specific setting, as may occur in countries undergoing socioeconomic transition, without

the concomitant introduction of TB transmission control measures, and the consequential increased effects of health care settings on the total incidence of TB in the population. In contrast, low PAF estimates might represent the effect of the introduction of TB transmission control measures in health care settings, and the consequential reduced effects of health care settings on total incidence of TB in the population.

Our study suggests that the effect of TB in HCWs (estimated by PAF) on global TB incidence does not exceed 5% and is comparable across countries with different background TB incidence, possibly because the proportion of HCWs and the TB IRRs counterbalance each other. Such estimated PAF figures call for urgent improvement of TB control measures in health care settings, particularly in settings that have a high IRR for TB.

The substantial heterogeneity between studies could be due to differences in methodologic quality, study design, sampling variability, and study populations across studies. In particular, some studies focused specifically on high-risk groups among HCWs, such as laboratory workers (19–21) and nurses (22). Furthermore, we could not discriminate between settings where infection transmission control measures were or were not implemented. It was equally difficult to classify settings according to their risk for admitting a person with unsuspected TB or according to the characteristics of the population attending the health care facilities, such as homeless persons, drug abusers, or migrants from countries with high TB prevalence.

The analysis restricted to the studies reporting cases among HCWs, regardless of their specific professional role, did not show any significant variation of the IRR estimates. Whether the HCWs investigated were representative of the national population of HCWs for each reported study was impossible to know; thus, IRR estimates resulting from the comparison of their risk for TB with the risk for the general population could be nonrepresentative of the national estimates. The density of HCWs and the level of income of each country, considered simultaneously in the analysis, accounted for some between-study variance. The IRR for TB decreased with the proportion of the population working in health care settings and in middle-income countries compared with high-income countries. Unfortunately, however, much of the heterogeneity still remained unexplained.

The method adopted to estimate the PAF typically measures the effects on a population of risk factors for noncommunicable diseases (11), ignoring the specific component of TB transmission dynamics between local community and health care settings (23). Thus, the PAF estimates presented here should be considered a lower limit for the real effects of controlling TB transmission in

health care settings, accounting for the remaining 2 main components acting on such an effect: the proportion of the population working in health care settings and the role of transmission control measures. Furthermore, PAF estimates do not account for patient-to-patient transmission of TB in health care facilities. Patient-to-patient transmission could constitute a major pathway of TB transmission, especially in settings of high HIV prevalence. Control of TB transmission in such a group could appreciably reduce the overall incidence of TB in the general population. However, data on TB transmission among patients are scanty and difficult to analyze in the context of this systematic review.

Although the cost-effectiveness assessment of the introduction of TB transmission control measures, encapsulated in the WHO Stop TB Strategy (24), is beyond the scope of this article, such a potential reduction of TB incidence among HCWs would make attractive a range of infection transmission control strategies. Managerial activities, administrative controls, adoption of N95 respirators, and engineering controls all represent potentially effective measures. The implementation of a sound TB infection control package based on the facility assessment, is hampered by constraints in poor resource settings; nevertheless, administrative controls alone have been proven effective in decreasing the risk for TB among HCWs (25–27).

In conclusion, our findings show that HCWs are at higher than average risk for infection with *Mycobacterium tuberculosis* and of developing TB disease. For this reason, sound TB infection control measures, including early diagnosis and prompt treatment of infectious cases, should be prioritized. Drug resistance emphasizes the urgency for implementing such measures, which would also benefit reduction of patient-to-patient transmission in health care facilities.

Acknowledgments

We thank WHO for their support in the study design, the collection, analysis and interpretation of data; in the writing of the report, and in the decision to submit the article for publication.

I.B. has been supported by the “Regione Piemonte” Italy, Assessorato Sanità, Progetti di Ricerca Sanitaria Finalizzata, 2008.

Dr Baussano is an infectious diseases physician and epidemiologist at the Unit of Cancer Epidemiology of the University ‘A. Avogadro’ and CPO–Piemonte in Novara, Italy, and at the School of Public Health of the Imperial College in London, UK. His research focuses on modeling the transmission dynamics of infectious diseases and on TB epidemiology, particularly among immigrants from countries with a high incidence of TB.

References

- Menzies D, Joshi R, Pai M. Risk of tuberculosis infection and disease associated with work in health care settings. *Int J Tuberc Lung Dis.* 2007;11:593–605.
- Sepkowitz KA. Tuberculosis and the health care worker: a historical perspective. *Ann Intern Med.* 1994;120:71–9.
- Jensen PA, Lambert LA, Iademarco MF, Ridzon R. Guidelines for preventing the transmission of *Mycobacterium tuberculosis* in health care settings, 2005. *MMWR Recomm Rep.* 2005;54(RR-17):1–141.
- Wright A, Zignol M, Van Deun A, Falzon D, Gerdes SR, Feldman K, et al. Epidemiology of antituberculosis drug resistance 2002–07: an updated analysis of the Global Project on Anti-Tuberculosis Drug Resistance Surveillance. *Lancet.* 2009;373:1861–73. DOI: 10.1016/S0140-6736(09)60331-7
- Joshi R, Reingold AL, Menzies D, Pai M. Tuberculosis among health care workers in low- and middle-income countries: a systematic review. *PLoS Med.* 2006;3:e494.
- Menzies D, Fanning A, Yuan L, Fitzgerald M. Tuberculosis among health care workers. *N Engl J Med.* 1995;332:92–8. DOI: 10.1056/NEJM199501123320206
- Seidler A, Nienhaus A, Diel R. Review of epidemiological studies on the occupational risk of tuberculosis in low-incidence areas. *Respiration.* 2005;72:431–46. DOI: 10.1159/000086261
- World Health Organization. Global health atlas [cited 2009 Feb 16]. <http://www.who.int/globalatlas/dataQuery/default.asp>
- Normand SL. Meta-analysis: formulating, evaluating, combining, and reporting. *Stat Med.* 1999;18:321–59. DOI: 10.1002/(SICI)1097-0258(19990215)18:3<321::AID-SIM28>3.0.CO;2-P
- Sterne JAC, Bradburn MJ, Egger M. Meta-analysis in Stata. In: Egger M, Davey-Smith G, Altman DG, editors. *Systematic reviews in health care: meta-analysis in context.* London: BMJ Publishing Group; 2001. p. 347–69.
- Levin ML. The occurrence of lung cancer in man. *Acta Unio Int Contra Cancrum.* 1953;9:531–41.
- Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ.* 2003;327:557–60.
- World Bank. World Bank country classification. 2008 [cited 2009 Dec 12]. <http://web.worldbank.org>
- Harries AD, Kamenya A, Namarika D, Msolomba IW, Salaniponi FM, Nyangulu DS, et al. Delays in diagnosis and treatment of smear-positive tuberculosis and the incidence of tuberculosis in hospital nurses in Blantyre, Malawi. *Trans R Soc Trop Med Hyg.* 1997;91:15–7.
- Rao KG, Aggarwal AN, Behera D. Tuberculosis among physicians in training. *Int J Tuberc Lung Dis.* 2004;8:1392–4.
- Gopinath KG, Siddique S, Kirubakaran H, Shanmugam A, Mathai E, Chandy GM. Tuberculosis among healthcare workers in a tertiary-care hospital in South India. *J Hosp Infect.* 2004;57:339–42. DOI: 10.1016/j.jhin.2004.03.029
- Sterne JAC, Egger M, Davey-Smith G. Investigating and dealing with publication and other biases. In: Egger M, Davey-Smith G, Altman DG, editors. *Systematic reviews in health care: meta-analysis in context.* London: BMJ Publishing Group; 2001. p. 189–208.
- Dimitrova B, Hutchings A, Atun R, Drobniewski F, Marchenko G, Zakharova S, et al. Increased risk of tuberculosis among health care workers in Samara Oblast, Russia: analysis of notification data. *Int J Tuberc Lung Dis.* 2005;9:43–8.
- Boudreau AY, Baron SL, Steenland NK, Van Gilder TJ, Decker JA, Galson SK, et al. Occupational risk of *Mycobacterium tuberculosis* infection in hospital workers. *Am J Ind Med.* 1997;32:528–34.
- Kim SJ, Lee SH, Kim IS, Kim HJ, Kim SK, Rieder HL. Risk of occupational tuberculosis in National Tuberculosis Programme laboratories in Korea. *Int J Tuberc Lung Dis.* 2007;11:138–42.
- Menzies D, Fanning A, Yuan L, FitzGerald JM. Factors associated with tuberculin conversion in Canadian microbiology and pathology workers. *Am J Respir Crit Care Med.* 2003;167:599–602.
- Babus V. Tuberculosis morbidity risk in medical nurses in specialized institutions for the treatment of lung diseases in Zagreb. *Int J Tuberc Lung Dis.* 1997;1:254–8.
- Angie B, Ann A, Malgosia G, Michael K, Hans K, Michael L, et al. *Tuberculosis control in prisons: a manual for programme managers.* Geneva: World Health Organization; 2000.
- Raviglione MC, Uplekar MW. WHO's new Stop TB Strategy. *Lancet.* 2006;367:952–5.
- Scano F. WHO policy on TB infection control in health care facilities, congregate settings and households. Geneva: World Health Organization; 2009.
- Fica CA, Cifuentes DM, Ajenjo HM, Jemenao PM, Zambrano OA, Febre VN, et al. Tuberculosis in health care workers [in Spanish]. *Rev Chilena Infectol.* 2008;25:243–55.
- Yanai H, Limpakarnjanarat K, Uthavivoravit W, Mastro TD, Mori T, Tappero JW. Risk of *Mycobacterium tuberculosis* infection and disease among health care workers, Chiang Rai, Thailand. *Int J Tuberc Lung Dis.* 2003;7:36–45.

Address for correspondence: Iacopo Baussano, School of Public Health, Faculty of Medicine, Rm 512 (5th Floor). St Mary's Campus, Norfolk Place, London W2 1PG, UK; email: i.baussano06@imperial.ac.uk

Like our podcasts?

Sign up to receive email announcements
when a new podcast is available.

www.cdc.gov/ncidod/eid/subscrib.htm



Tuberculosis among Health Care Workers

Technical Appendix

Search Terms

(“tuberculosis” or “tuberculosis, multidrug-resistant” or “extensively drug-resistant tuberculosis”) and (“occupational exposure” or “health personnel” or “outpatients” or “nursing homes” or “caregivers” or “physicians” or “medical staff, hospital” or “nurses” or “nurses” aides” or “nurse practitioners” or “students, medical” or “students, nursing” or “health and worker” or “health and care”) and (“cross infection” or “disease transmission” or “disease transmission, professional-to-patient” or “disease transmission, patient-to-professional” or “occupational diseases” or “occupational exposure” or “nosocomial tuberculosis” or “occupational tuberculosis” or “hospital exposure” or “occupational hazard”).

Levin’s Formula

To assess the fraction of tuberculosis in the population that was attributable to the exposure to health care settings, we calculated the PAF using Levin’s formula:

$$PAF\% = \frac{Pe \cdot (IRR - 1)}{1 + Pe \cdot (IRR - 1)} \cdot 100$$

where IRR is the TB IRR measured from each study, and *Pe* is the proportion of the population working in health care settings. As

List of Studies included in the Systematic Review

Twenty-five studies reported latent *Mycobacterium tuberculosis* infection (LTBI) cases among health care workers (HCWs) (Technical Appendix Table 1) (1–25). Eighteen studies reported tuberculosis cases among HCWs (Technical Appendix Table 2) (26–43). Three studies could not be located and were excluded from the analysis (44–46).

References

1. Adal KA, Anglim AM, Palumbo CL, Titus MG, Coyner BJ, Farr BM. The use of high-efficiency particulate air-filter respirators to protect hospital workers from tuberculosis. A cost-effectiveness analysis. *N Engl J Med.* 1994;331:169–73. [Medline DOI: 10.1056/NEJM199407213310306](#)
2. Baussano I, Bugiani M, Carosso A, Mairano D, Pia Barocelli A, Tagna M, et al. Risk of tuberculin conversion among healthcare workers and the adoption of preventive measures. *Occup Environ Med.* 2007;64:161–6. [Medline DOI: 10.1136/oem.2006.028068](#)
3. LoBue PA, Catanzaro A. Effectiveness of a nosocomial tuberculosis control program at an urban teaching hospital. *Chest.* 1998;113:1184–9. [Medline DOI: 10.1378/chest.113.5.1184](#)
4. Liss GM, Khan R, Koven E, Simor AE. Tuberculosis infection among staff at a Canadian community hospital. *Infect Control Hosp Epidemiol.* 1996;17:29–35. [Medline DOI: 10.1086/647185](#)
5. Miller AK, Tepper A, Sieber K. Historical risks of tuberculin skin test conversion among non-physician staff at a large urban hospital. *Am J Ind Med.* 2002;42:228–35. [Medline DOI: 10.1002/ajim.10108](#)
6. Behrman AJ, Shofer FS. Tuberculosis exposure and control in an urban emergency department. *Ann Emerg Med.* 1998;31:370–5. [Medline DOI: 10.1016/S0196-0644\(98\)70349-X](#)
7. Larsen NM, Biddle CL, Sotir MJ, White N, Parrott P, Blumberg HM. Risk of tuberculin skin test conversion among health care workers: occupational versus community exposure and infection. *Clin Infect Dis.* 2002;35:796–801. [Medline DOI: 10.1086/342333](#)
8. Blumberg HM, Sotir M, Erwin M, Bachman R, Shulman JA. Risk of house staff tuberculin skin test conversion in an area with a high incidence of tuberculosis. *Clin Infect Dis.* 1998;27:826–33. [Medline DOI: 10.1086/514963](#)
9. Zahnaw K, Matts JP, Hillman D, Finley E, Brown LS Jr, Torres RA, et al. Rates of tuberculosis infection in healthcare workers providing services to HIV-infected populations. Terry Beirn Community Programs for Clinical Research on AIDS. *Infect Control Hosp Epidemiol.* 1998;19:829–35. [Medline DOI: 10.1086/647740](#)
10. Warren DK, Foley KM, Polish LB, Seiler SM, Fraser VJ. Tuberculin skin testing of physicians at a midwestern teaching hospital: a 6-year prospective study. *Clin Infect Dis.* 2001;32:1331–7. [Medline DOI: 10.1086/319993](#)

11. Menzies D, Fanning A, Yuan L, FitzGerald JM. Factors associated with tuberculin conversion in Canadian microbiology and pathology workers. *Am J Respir Crit Care Med.* 2003;167:599–602. [Medline DOI: 10.1164/rccm.200208-873BC](#)
12. Menzies D, Fanning A, Yuan L, FitzGerald JM. Hospital ventilation and risk for tuberculous infection in Canadian health care workers. Canadian Collaborative Group in Nosocomial Transmission of TB. *Ann Intern Med.* 2000;133:779–89. [Medline](#)
13. Levy MZ, Medeiros EA, Shang N, Soares MC, Homenko AS, Almeida RM, et al. TST reversion in a BCG-revaccinated population of nursing and medical students, Sao Paulo, Brazil, 1997–2000. *Int J Tuberc Lung Dis.* 2005;9:771–6. [Medline](#)
14. Silva VM, Cunha AJ, Kritski AL. Tuberculin skin test conversion among medical students at a teaching hospital in Rio de Janeiro, Brazil. *Infect Control Hosp Epidemiol.* 2002;23:591–4. [Medline DOI: 10.1086/501976](#)
15. Maciel EL, Viana MC, Zeitoune RC, Ferreira I, Fregona G, Dietze R. Prevalence and incidence of *Mycobacterium tuberculosis* infection in nursing students in Vitoria, Espirito Santo. *Rev Soc Bras Med Trop.* 2005;38:469–72. [Medline DOI: 10.1590/S0037-86822005000600004](#)
16. Lopes LK, Teles SA, Souza AC, Rabahi MF, Tipple AF. Tuberculosis risk among nursing professionals from Central Brazil. *Am J Infect Control.* 2008;36:148–51. [Medline DOI: 10.1016/j.ajic.2007.01.013](#)
17. Hohmuth BA, Yamanija JC, Dayal AS, Nardell E, Salazar JJ, Smith Fawzi MC. Latent tuberculosis infection: risks to health care students at a hospital in Lima, Peru. *Int J Tuberc Lung Dis.* 2006;10:1146–51. [Medline](#)
18. Pai M, Gokhale K, Joshi R, Dogra S, Kalantri S, Mendiratta DK, et al. *Mycobacterium tuberculosis* infection in health care workers in rural India: comparison of a whole-blood interferon gamma assay with tuberculin skin testing. *JAMA.* 2005;293:2746–55. [Medline DOI: 10.1001/jama.293.22.2746](#)
19. Corbett EL, Muzangwa J, Chaka K, Dauya E, Cheung YB, Munyati SS, et al. Nursing and community rates of *Mycobacterium tuberculosis* infection among students in Harare, Zimbabwe. *Clin Infect Dis.* 2007;44:317–23. [Medline DOI: 10.1086/509926](#)
20. Bonifacio N, Saito M, Gilman RH, Leung F, Cordova Chavez N, Chacaltana Huarcaya J, et al. High risk for tuberculosis in hospital physicians, Peru. *Emerg Infect Dis.* 2002;8:747–8. [Medline](#)

21. Yanai H, Limpakarnjanarat K, Uthaiworavit W, Mastro TD, Mori T, Tappero JW. Risk of Mycobacterium tuberculosis infection and disease among health care workers, Chiang Rai, Thailand. *Int J Tuberc Lung Dis.* 2003;7:36–45. [Medline](#)
22. Rullán JV, Herrera D, Cano R, Moreno V, Godoy P, Peiro EF, et al. Nosocomial transmission of multidrug-resistant Mycobacterium tuberculosis in Spain. *Emerg Infect Dis.* 1996;2:125–9. [Medline](#) DOI: [10.3201/eid0202.960208](https://doi.org/10.3201/eid0202.960208)
23. Roth VR, Garrett DO, Laserson KF, Starling CE, Kritski AL, Medeiros EA, et al. A multicenter evaluation of tuberculin skin test positivity and conversion among health care workers in Brazilian hospitals. *Int J Tuberc Lung Dis.* 2005;9:1335–42. [Medline](#)
24. Lee SS, Liu YC, Huang TS, Chen YS, Tsai HC, Wann SR, et al. Comparison of the interferon-gamma release assay and the tuberculin skin test for contact investigation of tuberculosis in BCG-vaccinated health care workers. *Scand J Infect Dis.* 2008;40:373–80. [Medline](#) DOI: [10.1080/00365540701730743](https://doi.org/10.1080/00365540701730743)
25. Louthier J, Rivera P, Feldman J, Villa N, DeHovitz J, Sepkowitz KA. Risk of tuberculin conversion according to occupation among health care workers at a New York City hospital. *Am J Respir Crit Care Med.* 1997;156:201–5. [Medline](#)
26. Krüüner A, Danilovitsh M, Pehme L, Laisaar T, Hoffner SE, Katila ML. Tuberculosis as an occupational hazard for health care workers in Estonia. *Int J Tuberc Lung Dis.* 2001;5:170–6. [Medline](#)
27. Cuhadaroglu C, Erelel M, Tabak L, Kilicaslan Z. Increased risk of tuberculosis in health care workers: a retrospective survey at a teaching hospital in Istanbul, Turkey. *BMC Infect Dis.* 2002;2:14. [Medline](#) DOI: [10.1186/1471-2334-2-14](https://doi.org/10.1186/1471-2334-2-14)
28. Skodric-Trifunovic V, Markovic-Denic L, Nagorni-Obradovic L, Vlajinac H, Woeltje KF. The risk of occupational tuberculosis in Serbian health care workers. *Int J Tuberc Lung Dis.* 2009;13:640–4. [Medline](#)
29. Jo KW, Woo JH, Hong Y, Choi CM, Oh YM, Lee SD, et al. Incidence of tuberculosis among health care workers at a private university hospital in South Korea. *Int J Tuberc Lung Dis.* 2008;12:436–40. [Medline](#)
30. Fica CA, Ramonda CP, Jemenao PM, Zambrano GA, Cifuentes DM, Febre VN, et al. [Tuberculosis in health care workers from a public health service in Santiago, Chile]. *Rev Chilena Infectol.* 2009;26:34–8. [Medline](#)

31. Dimitrova B, Hutchings A, Atun R, Drobniewski F, Marchenko G, Zakharova S, et al. Increased risk of tuberculosis among health care workers in Samara Oblast, Russia: analysis of notification data. *Int J Tuberc Lung Dis.* 2005;9:43–8. [Medline](#)
32. Gopinath KG, Siddique S, Kirubakaran H, Shanmugam A, Mathai E, Chandy GM. Tuberculosis among healthcare workers in a tertiary-care hospital in South India. *J Hosp Infect.* 2004;57:339–42. [Medline DOI: 10.1016/j.jhin.2004.03.029](#)
33. Naidoo S, Jinabhai CC. TB in health care workers in KwaZulu-Natal, South Africa. *Int J Tuberc Lung Dis.* 2006;10:676–82. [Medline](#)
34. Alonso-Echanove J, Granich RM, Laszlo A, Chu G, Borja N, Blas R, et al. Occupational transmission of Mycobacterium tuberculosis to health care workers in a university hospital in Lima, Peru. *Clin Infect Dis.* 2001;33:589–96. [Medline DOI: 10.1086/321892](#)
35. Jelip J, Mathew GG, Yusin T, Dony JF, Singh N, Ashaari M, et al. Risk factors of tuberculosis among health care workers in Sabah, Malaysia. *Tuberculosis (Edinb).* 2004;84:19–23. [Medline DOI: 10.1016/j.tube.2003.08.015](#)
36. Harries AD, Kamenya A, Namarika D, Msolomba IW, Salaniponi FM, Nyangulu DS, et al. Delays in diagnosis and treatment of smear-positive tuberculosis and the incidence of tuberculosis in hospital nurses in Blantyre, Malawi. *Trans R Soc Trop Med Hyg.* 1997;91:15–7. [Medline DOI: 10.1016/S0035-9203\(97\)90376-X](#)
37. Rao KG, Aggarwal AN, Behera D. Tuberculosis among physicians in training. *Int J Tuberc Lung Dis.* 2004;8:1392–4. [Medline](#)
38. Harries AD, Hargreaves NJ, Gausi F, Kwanjana JH, Salaniponi FM. Preventing tuberculosis among health workers in Malawi. *Bull World Health Organ.* 2002;80:526–31. [Medline](#)
39. Harries AD, Nyirenda TE, Banerjee A, Boeree MJ, Salaniponi FM. Tuberculosis in health care workers in Malawi. *Trans R Soc Trop Med Hyg.* 1999;93:32–5. [Medline DOI: 10.1016/S0035-9203\(99\)90170-0](#)
40. Laniado-Laborín R, Cabrales-Vargas N. Tuberculosis in healthcare workers at a general hospital in Mexico. *Infect Control Hosp Epidemiol.* 2006;27:449–52. [Medline DOI: 10.1086/503641](#)
41. Kanyerere HS, Salaniponi FM. Tuberculosis in health care workers in a central hospital in Malawi. *Int J Tuberc Lung Dis.* 2003;7:489–92. [Medline](#)
42. Wilkinson D, Gilks CF. Increasing frequency of tuberculosis among staff in a South African district hospital: impact of the HIV epidemic on the supply side of health care. *Trans R Soc Trop Med Hyg.* 1998;92:500–2. [Medline DOI: 10.1016/S0035-9203\(98\)90889-6](#)

43. Torres Costa J, Sa R, Cardoso MJ, Silva R, Ferreira J, Ribeiro C, et al. Tuberculosis screening in Portuguese healthcare workers using the tuberculin skin test and the interferon-gamma release assay. *Eur Respir J*. 2009;34:1423–8. [Medline DOI: 10.1183/09031936.00053809](#)
44. Condos R, Schluger N, Lacouture R, Rom W. Tuberculosis infections among housestaff at Bellevue Hospital in an epidemic period. [abstract]. *Am Rev Respir Dis*. 1993;147(Suppl):A124.
45. Redwood E, Anderson V, Felton C, Findley S, Ford J. Tuberculin conversions in hospital employees in a high tuberculosis prevalence area. *Am Rev Respir Dis* ;147:Suppl:A119 abstract. 1993.
46. Lainez RM, Consul M, Olona M, Martinez-Ballarín JI, Miravittles M, Vidal R. [Tuberculous infection in nursing students: prevalence and conversion during a 3-year follow-up]. *Med Clin (Barc)*. 1999;113:685–9. [Medline](#)

Technical Appendix 1 Table 1. Studies reporting incidence of latent TB infection among HCWs, by incidence level, January 2005–July 2010*

Study, country, and reference	Study period	Types of hospital workers	No. cases (no. persons at risk)	ARTI, %†
Incidence <50/100,000				
Adal et al., United States (1)	1990–1992	All	23 (124,869)	0.21
Baussano et al., Italy (2)	1998–2004	All	146 (8,122)	1.80
LoBue et al., United States (3)	1993–1995	All	59 (9,905)	0.60
Liss et al., Canada (4)	1991–1994	All	18 (809)	2.22
Miller et al., United States (5)	1990–1992	All	138 (2,362)	5.84
Behrman et al., United States (6)	1994–1995	All	57 (2,564)	2.22
Larsen et al., United States (7)	1994–1998	All	69 (5,773)	1.19
Blumberg et al., United States (8)	1992–1997	All	52 (2,144)	2.43
Zahnow et al., United States (9)	1992–1995	All	22 (766)	2.87
Warren et al., United States (10)	1992–1998	All	21 (731)	2.87
Menzies et al., Canada (11)	Not reported	Microbiology and pathology	14 (111)	12.61
Louther et al., United States (12)	1991–1994	All	65 (898)	7.24
Menzies et al., Canada (13)	1995–1996	All	238 (1,289)	18.46
Rullán et al., Spain (14)	1991–1995	All	24 (92)	26.08
Lee et al., South Korea (15)	2007	All	16 (196)	8.16
Incidence 50–99/100,000				
Levy et al., Brazil (16)	1997–2000	All	1 (46)	2.17
Silva et al., Brazil (17)	1998–1999	All	16 (414)	3.86
Roth et al., Brazil (18)	1998–1999	All	105 (1209)	8.68
Maciel et al., Brazil (19)	1997–1999	All	8 (76)	10.53
Lopes et al., Brazil (20)	2001–2004	All	7 (61)	11.48
Incidence ≥100/100,000				
Hohmuth et al., Peru (21)	2002–03	Health care students	1 (93)	1.08
Pai et al., India (22)	2004	All	6 (147)	4.08
Corbett et al., Zimbabwe (23)	2004–2005	Student nurses	41 (213)	19.25
Bonifacio et al., Peru (24)	2001	All	5 (35)	14.29
Yanai et al., Thailand (25)	1995–1996	All	24 (332)	7.24

*Incidence levels from World Health Organization statistics, 2009 (8). TB, tuberculosis; HCW, health care worker; ARTI, annual risk for *Mycobacterium tuberculosis* infection.

†Estimates of ARTI ratio between HCWs and general population, pooled estimates of ARTI ratio (by estimated TB incidence), fraction of the population working as HCW.


Technical Appendix 1 Table 2. Studies reporting TB incidence among HCWs, January 2005–July 2010*

Author, country, and reference	Period	Hospital workers/comparison group	No. cases (no. persons at risk)	TB incidence in HCWs, ×100,000	TB incidence in the comparison, ×100,000	TB IRR† (95% CI)	Incidence difference	HCW population, ×1,000	PAF%
Incidence <50/100,000‡									
Cuhadaroglu C et al., Turkey (27)	1991–2000	All/national estimates for Turkey	31 (33,590)	92.29	29	3.18 (2.24–4.53)	63.29	4.47	0.97
Skodric-Trifunovic V et al., Serbia (28)	1992–2004	All/national estimates for Serbia§	24 (57,279)	41.9	34	1.23 (0.83–1.84)	8	6.34	0.15
Fica AC et al., Chile (30)	2003–2006	All/national estimates for Chile	12 (30,680)	39.1	23	1.71 (0.97–3.01)	16.26	1.69	0.12
Torres Costa J et al., Portugal (43)	2005–2008	All/national estimates for Portugal	31 (16,178)	191.62	32	5.99 (4.21–8.51)	159.62	8.17	3.91
Incidence 50–99/100,000‡									
Dimitrova B et al. (2005), Russia (31)	1994–2002	All/national estimates for Russia§	474 (583,695)	81.21	95	0.86 (0.78–0.94)	6.21	12.78	–0.19
Kruuner A et al., Estonia (26)	1994–1998	All/national estimates for Estonia§	67 (73,650)	90.97	58	1.38 (1.08–1.75)	32.97	1142	0.43
Jo KW et al., South Korea (29)	2001–2006	All/national estimates for South Korea	61 (8,433)	723.35	92	7.86 (6.12–10.11)	631.35	3.50	2.35
Incidence ≥100/100,000‡									
Gopinath KG et al., India (32)	1992–2001	All/national estimates for India§	125 (60,163)	207.77	168	1.24 (1.04–1.47)	39.77	1.89	0.04
Naidoo S and Jinabhai CC, South Africa (33)	2004–2005	All/national estimates for South Africa§	583 (49,392)	1180.35	911	1.30 (1.19–1.41)	462.35	4.85	0.14
Alonso-Echanove J et al., Peru (34)	1994–1998	All/national estimates for Peru§	36 (9,200)	391.30	230	1.70 (1.23–2.36)	161.30	1.84	0.13
Jelip J et al., Malaysia (35)	1990–2000	All/national estimates for Malaysia§	205 (73,120)	280.36	114	2.46 (2.14–2.82)	126.36	2.51	0.37
Harries AD et al., Malawi (36)	1993–1994	All/national estimates for Malawi§	12 (620)	1935.48	358	5.41 (3.07–9.52)	1,577.48	0.61	0.27
Rao KG et al., India (37)	2001	All/national estimates for India§	13 (1,032)	1259.69	168	7.50 (4.35–12.91)	1,091.69	1.89	1.21
Harries AD et al., Malawi (38)	1996–1999	All/national estimates for Malawi§	96 (2,979)	3222.56	405	7.96 (6.51–9.72)	2,817.56	0.61	0.42
Harries AD et al. (1999), Malawi (39)	1996	All/national estimates for Malawi§	108 (3,042)	3550.30	389	9.13 (7.56–11.02)	3,161.30	0.61	0.49
Laniado-Laborin R et al., Mexico (40)	1999–2003	All/Tijuana, general population	18 (4,095)	439.56	40	10.99 (6.92–17.44)	399.56	3.34	3.23
Wilkinson D and Gilks CF, South Africa (42)	1991–1996	All/national estimates for South Africa§	13 (2,328)	558.42	311	1.80 (1.04–3.09)	247.42	4.85	0.38
Kanyerere HS et al., Malawi (41)	2001	All/national estimates for Malawi§	33 (571)	5779.33	414	13.96 (9.92–19.64)	5,360.33	0.61	0.78


*TB, tuberculosis; HCW, health care worker; IRR, incidence rate ratio; CI, confidence interval; PAF%, fraction of TB in the population attributable to TB exposure in health care settings. WHO, World Health Organization.

†Pooled estimates of annual incidence rate ratio (by estimated TB incidence).

‡World Health Organization statistics, 2009 (8).



Technical Appendix 1 Figure 1. Funnel Plot for the studies reporting tuberculosis incidence among health care workers. Studies are labeled using the country and year of the study and the reference in parentheses. * <50 cases/100,000 population; ** 50–99 cases/100,000 population; *** ≥100 cases/100,000 population. TB, tuberculosis; HCW, health care worker; IRR, incidence rate ratio.



- \square Studies from countries with TB incidence <math><50</math> cases/100,000 population
- \blacklozenge Studies from countries with TB incidence

Technical Appendix 1 Figure 2. Contour plot of relationship between the proportion of exposed population, incidence rate ratio (IRR), and population-attributable fraction (PAF) in review of studies about tuberculosis among health care workers. The proportion of the population working in health care settings is reported on the x-axis, y-axis reports the PAF on a log scale calculated by using the Levin formula. The isoclines represent different levels of IRR. The asterisk (*) indicates the fraction (%) of tuberculosis (log scale) in the population attributable to exposure in health care settings. Points are labeled by the country and year of the study, with the reference in parentheses.