

4. De Beenhouwer H, Liang Z, De Rijk P, van Eekeren C, Portaels F. Detection and identification of mycobacteria by DNA amplification and oligonucleotide-specific capture plate hybridization. *J Clin Microbiol.* 1995;33:2994–8.
5. Reischl U, Feldmann K, Naumann L, Gaugler BJM, Ninet B, Hirschel B, et al. 16S rRNA sequence diversity in *Mycobacterium celatum* strains caused by presence of two different copies of 16S rRNA gene. *J Clin Microbiol.* 1998;36:1761–4.
6. Okazaki M, Ohkusu K, Hata H, Ohnishi H, Sugahara K, Kawamura C, et al. *Mycobacterium kyorinense* sp. nov., a novel, slow-growing species, related to *Mycobacterium celatum*, isolated from human clinical specimens. *Int J Syst Evol Microbiol.* 2009;59:1336–41. DOI: 10.1099/ijs.0.000760-0
7. McNabb A, Eisler D, Adie K, Amos M, Rodrigues M, Stephens G, et al. Assessment of partial sequencing of the 65-kiloDalton heat shock protein gene (*hsp65*) for routine identification of *Mycobacterium* species isolated from clinical sources. *J Clin Microbiol.* 2004;42:3000–11. DOI: 10.1128/JCM.42.7.3000-3011.2004
8. Kim BJ, Lee SH, Lyu MA, Kim SJ, Bai GH, Chae GT, et al. Identification of mycobacterial species by comparative sequence analysis of the RNA polymerase gene (*rpoB*). *J Clin Microbiol.* 1999;37:1714–20.
9. Adékambi T, Drancourt M. Dissection of phylogenetic relationships among 19 rapidly growing *Mycobacterium* species by 16S rRNA, *hsp65*, *sodA*, *recA*, and *rpoB* gene sequencing. *Int J Syst Evol Microbiol.* 2004;54:2095–105. DOI: 10.1099/ijs.0.63094-0
10. de Lisle GW, Kawakami RP, Yates GF, Collins DM. Isolation of *Mycobacterium bovis* and other mycobacterial species from ferrets and stoats. *Vet Microbiol.* 2008;132:402–7. DOI: 10.1016/j.vetmic.2008.05.022

Address for correspondence: Eva Ludwig, Institute of Veterinary Pathology, Ludwig-Maximilians-University Munich, Veterinärstr. 13, D-80539 Munich, Germany; email: ludwig@patho.vetmed.uni-muenchen.de



Human Infection with *Pseudoterranova azarasi* Roundworm

To the Editor: Eating raw or undercooked marine fish may lead to infection with several helminths. Members of the *Pseudoterranova decipiens* species complex are the second most common nematodes found in humans (most common are nematodes of the *Anisakis simplex* complex) (1,2). The *P. decipiens* species complex consists of at least 5 sibling species (genetically but not morphologically distinguishable): *P. decipiens sensu stricto*, *P. azarasi*, *P. cattani*, *P. krabbei*, and *P. bulbosa*. In northern Japan, human infection with *Pseudoterranova* spp. is not rare; by the mid-1990s, as many as 769 cases had been reported (3). Pseudoterranovosis has also been encountered in North and South America and Europe (4,5). However, possible biologic and geographic differences of the sibling species of genus *Pseudoterranova* in relation to human infection remain unknown. We report a case of pseudoterranovosis for which the sibling species was confirmed as *P. azarasi*.

In 2009, a woman in Japan coughed up a nematode and expelled it through her mouth. Her medical history was unremarkable, and she had not traveled abroad for the past few years. Measurements of the worm were as follows: body 35 mm long and 0.85 mm wide; esophagus 1.88 mm long; ventriculus 1.05 mm long, extending anteriorly along the ventriculus. The anterior end of the worm contained 3 lips. The tail was conical, 0.21 mm long, and had a small, knob-like process at the posterior end. On the basis of morphologic features, the worm was identified as a 4th-stage larva of *P. decipiens* (sensu lato) roundworms.

DNA was extracted from the isolate from the patient (clinical isolate) and from isolates of *P. decipiens* (sensu lato) larvae from Pacific cod purchased at a local market. PCR was performed and the amplification products were directly sequenced. Primers used were 5'-CCGGGCAAAAGTCGTAACAA-3' and 5'-ATATGCTTAAATTCAGCGGGT-3' for a region that spans the internal transcribed spacer (ITS) 1, ITS2, and 5.8S rRNA; 5'-CTACTACTAAGAATTTGCGT-3' and 5'-AATCCAAATACTTACGAGGA-3' for cytochrome oxidase subunit 1; and 5'-CAGCGTATTGGTCCTAATAA-3' and 5'-AGCATAAACAAAAGTAAACTCA-3' for NADH dehydrogenase subunit 1.

Nucleotide sequences for the clinical and Pacific cod isolates (GenBank accession nos. AB576756–AB576761) were compared with those in DNA databases available to the public. The ITS1 and ITS2 sequences of the clinical and Pacific cod isolates were identical to those of *P. azarasi* roundworms; however, the ITS sequences of these 2 isolates also showed close similarity to those of *P. decipiens* (sensu stricto) worms and differed by only 1 nt in ITS2. However, phylogenetic tree analyses of NADH dehydrogenase subunit 1 and cytochrome oxidase subunit 1 sequences showed that the clinical and Pacific cod roundworm isolates clustered with *P. azarasi* and were clearly distinguished from *P. decipiens* (sensu stricto), showing that the clinical isolate belonged to *P. azarasi* (Figure).

Geographic distribution of the 5 sibling species of the *P. decipiens* complex differs somewhat. *P. azarasi* and *P. bulbosa* are found in northwestern Pacific (including Japan), *P. decipiens* (sensu stricto) and *P. krabbei* in northeastern Atlantic, *P. decipiens* (sensu stricto) in northwestern Atlantic, and *P. cattani* in southeastern Pacific waters (1). Given this distribution, it is not surprising

that the clinical isolate and the larvae from the Pacific cod were identified as *P. azarasi*, 1 of 2 species found in water near Japan. Adult worms live in the intestines of seals and sea lions, and infective larvae live in the tissues of various marine fish, including cod, pollack, and smelt (1).

In Japan, most patients infected with *Pseudoterranova* spp. have acute or subacute abdominal pain, and larvae are extracted from the stomach endoscopically. However, for some patients, diagnosis is made when 4th-stage larvae are expelled from the mouth, indicating that the larvae developed from the 3rd to 4th stage during the time of infection, as did the worm reported here. Expulsion of *Pseudoterranova* spp. larvae from the mouth in the absence of severe gastric symptoms occurs more commonly in Chile (5). Whether the varied symptoms triggered by infection with *Pseudoterranova* spp. larvae reflect different responses of individual hosts to the worms or whether the pathogenicity of *Pseudoterranova* spp. in humans differs among worm species remains to be elucidated.

Because of the increasing worldwide popularity of eating sushi and sashimi made of raw marine fish, consumers should be made aware of the possible risk for fish-borne parasitoses. Freezing and storing fish at -20°C for 7 days or freezing

at -35°C until solid and storing at -35°C for 15 hours is sufficient to kill parasites (6).

This work was supported in part by grants-in-aids from the Ministry of Health, Labour and Welfare of Japan (H20-Shinko).

**Naoki Arizono,
Toshiyuki Miura,
Minoru Yamada,
Tatsuya Tegoshi,
and Kotaro Onishi**

Author affiliations: Kyoto Prefectural University of Medicine, Kyoto, Japan (N. Arizono, M. Yamada, T. Tegoshi, K Onishi); and The University of Tokyo, Tokyo, Japan (T. Miura)

DOI: 10.3201/eid1703.101350

References

1. Mattiucci S, Nascetti G. Advances and trends in the molecular systematics of anisakid nematodes, with implications for their evolutionary ecology and host-parasite co-evolutionary processes. *Adv Parasitol.* 2008;66:47–148. DOI: 10.1016/S0065-308X(08)00202-9
2. Zhu XQ, D'Amelio S, Palm HW, Paggi L, George-Nascimento M, Gasser RB. SSCP-based identification of members within the *Pseudoterranova decipiens* complex (Nematoda: Ascaridoidea: Anisakidae) using genetic markers in the internal transcribed spacers of ribosomal DNA. *Parasitology.* 2002;124:615–23. DOI: 10.1017/S0031182002001579

3. Ishikura H. Anisakiasis. 2. Clinical pathology and epidemiology. *Progress of Medical Parasitology in Japan.* 2003;8:451–73.
4. Audicana MT, Kennedy MW. *Anisakis simplex*: from obscure infectious worm to inducer of immune hypersensitivity. *Clin Microbiol Rev.* 2008;21:360–79 DOI: 10.1128/CMR.00012-07.
5. Torres P, Jercic MI, Weitz JC, Dobrew EK, Mercado RA. Human pseudoterranovosis, an emerging infection in Chile. *J Parasitol.* 2007;93:440–3. DOI: 10.1645/GE-946R.1
6. US Food and Drug Administration. Parasites. In: *Fish and fisheries products hazards and controls guidance*, 3rd ed. Rockville (MD): The Administration; 2001. p. 65–72.

Address for correspondence: Naoki Arizono, Department of Medical Zoology, Kyoto Prefectural University of Medicine, Kawaramachi-Hirokoji, Kyoto 602-8566, Japan; email: arizonon@koto.kpu-m.ac.jp

Mycobacterium mageritense Pulmonary Disease in Patient with Compromised Immune System

To the Editor: *Mycobacterium mageritense* is one of the rapidly growing mycobacteria (RGM). It was first isolated in Spain in 1987, described as a new species in 1997 by Domenech et al. (1), and first described and associated with disease in the United States in 2002 (2). In the 2002 report, 6 isolates were recovered from sputum, a bronchoscopy sample, a wound infection after liposuction, the blood of an immunosuppressed patient with a central catheter and sepsis, a patient with severe sinusitis, and from a wound infection in a patient who had probable osteomyelitis after fixation of an open fracture. It has since




Figure. Phylogenetic analysis of members of *Pseudoterranova decipiens* species complex roundworms. Genetic relationships between NADH dehydrogenase subunit 1 sequences in clinical and Pacific cod isolates and species were inferred by using the neighbor-joining method. Bootstrap values (1,000 replicates) are shown next to the branches. The final dataset contained 498 positions. **P. decipiens* sensu stricto. Scale bar indicates nucleotide substitutions per site.