

Our reply supplies supplementary clinical and laboratory findings that could be useful for discussion.

During our patient's hospitalization, we were able to investigate 6/8 diagnostic criteria for HLH proposed by the Histiocyte Society (3). Of these, only 2 or 3 were found: persistent fever at 38.5°C; hypertriglyceridemia at 267 mg/dL (analysis performed at day 7); and cytopenias, which preferentially affected erythrocytes and lymphocytes. (Thrombocytopenia was moderate at $>100 \times 10^9$ platelets/L, and no neutropenia was found [$>2 \times 10^9$ neutrophils/L].) Liver function was not affected; no hepatomegaly was found, and alanine aminotransferase, aspartate aminotransferase, and bilirubin levels remained within reference ranges. Physical examinations did not detect splenomegaly, and laboratory findings did not show hypofibrinogenemia or ferritin level exceeding 500 ng/mL. A bone marrow biopsy performed on day 2 of hospitalization did not show hemophagocytosis. Studies of natural killer cell function and soluble CD25, which are also proposed diagnostic criteria for HLH, were not performed.

Overall, we found that the arguments in favor of HLH were too limited to consider this diagnosis and initiate an aggressive therapeutic approach based on immunosuppressive drugs. Even if, in the event of HLH, an early and appropriate treatment can be life-saving, the destruction of the remaining immune functions might also be lethal for the patient.

**Julien Lupo,
Jean-François Timsit,
and Patrice Morand**

Author affiliation: Centre Hospitalier Universitaire de Grenoble and Université Joseph Fourier, Grenoble, France

DOI: <http://dx.doi.org/10.3201/eid1809.120749>

References

1. Iaria C, Leonardi MS, Buda A, Toro ML, Cascio A. Measles and secondary hemophagocytic lymphohistiocytosis [letter]. *Emerg Infect Dis.* 2012;18:1529.
2. Lupo J, Bernard S, Wintenberger C, Baccard M, Vabret A, Antona D, et al. Fatal measles without rash in immunocompetent adult, France. *Emerg Infect Dis.* 2012;18:521–3. <http://dx.doi.org/10.3201/eid1803.111300>
3. Janka GE. Familial and acquired hemophagocytic lymphohistiocytosis. *Annu Rev Med.* 2012;63:233–46. <http://dx.doi.org/10.1146/annurev-med-041610-134208>

Address for correspondence: Julien Lupo, Laboratoire de Virologie, Institut de Biologie et Pathologie Centre Hospitalier Universitaire de Grenoble, BP 217, Grenoble Cedex 9 38043, France; email: jlupo@chu-grenoble.fr

Contaminated Soil and Transmission of Influenza Virus (H5N1)

To the Editor: Highly pathogenic avian influenza (HPAI) virus (H5N1) has been responsible for 603 confirmed human cases worldwide, including 356 that resulted in death, and for >7,000 epizootic outbreaks (1,2). Direct contact between hosts is the main mechanism of transmission for avian influenza viruses, but the possible role of the environment as a source of HPAI virus (H5N1) infection has been rarely studied, particularly in the context of countries where the virus is enzootic or epizootic (3–7). To determine if contaminated soil contributes to the transmission cycle of HPAI virus (H5N1), we used experimental and simulated field conditions to assess possible transmission in chickens.

All experiments were conducted by using HPAI virus (H5N1) strain A/chicken/Cambodia/LC1AL/2007 (GenBank accession nos. HQ200574–HQ200581). All animal experiments were conducted in the biosafety level 3 laboratory of Institut Pasteur in Cambodia (IPC), in compliance with the European Community 86/609/CEE directive and approved by the Animal Ethics Committee of IPC (permit: AEC/IPC/003/2010). Specific pathogen-free (SPF) chickens were provided by the National Veterinary Research Institute of Cambodia.

We used 3 types of soil: 1) sandy topsoil collected from around rice fields in Phnom Penh Province, Cambodia; 2) building sand purchased from a local building company; and 3) soil-based compost purchased from a local tree nursery. Physicochemical and microbiologic parameters were measured for water extracts obtained for each type of soil (online Technical Appendix Table, www.cdc.gov/EID/pdfs/12-0402-Techapp.pdf), and low- and high-dose contamination protocols (online Technical Appendix Figure) were used to experimentally contaminate each soil type. In brief, we seeded the soil samples with 1–56 infectious units of contaminated feces; 1 infectious unit was defined as 1 g feces from an SPF duck mixed with $1 \times 10^{7.8}$ 50% egg infective dose of HPAI virus (H5N1) particles. The contaminated soil was then sprinkled on the bottom of an isolator (surface area 0.2 m²) in which the chickens were housed. Oropharyngeal and cloacal swab samples and feathers were collected daily from the chickens and underwent quantitative reverse transcription PCR (qRT-PCR) testing targeting the H5 hemagglutinin gene (8). Surviving birds were killed humanely at the end of the experiments, and postmortem examination and collection of serum and organ samples were conducted on all animals. Organ samples were tested by using qRT-PCR, and serum samples were tested

by using hemagglutination inhibition assay (9).

No clinical symptoms, deaths, or seroconversion for HPAI virus (H5N1) were observed in chickens exposed to contaminated sandy topsoil, regardless of the dose protocol used (Table). However, for building sand and soil-based compost, the high-dose contamination protocol, starting with 8 infectious units, resulted in a 100% mortality rate by day 4. Low-dose protocols, starting at 1 infectious unit, resulted in survival of all birds at day 24, with no clinical symptoms and no virus detected in the samples collected postmortem. However, seroconversion for HPAI virus (H5N1) was observed in 33% and 50% of the chickens exposed to building sand and compost, respectively (Table).

Soil-based compost and building sand, although existing in natural settings, are not the most common substrates found in places where free-ranging poultry are raised in Cambodia. Therefore, despite the high mortality rate observed in our study after exposure to highly contaminated soils, the role of these soil types in transmission of HPAI virus (H5N1) infection to poultry or other species, including humans, appears limited when replaced in actual epizootic or enzootic field conditions. Our results

also show that exposure of chickens to moderately contaminated soil may result in a protective immune response.

Sandy topsoil, on the other hand, did not allow any transmission of HPAI virus (H5N1) from the environment to chickens. This type of soil, which covers $\approx 40\%$ of the rice-growing areas of Cambodia (10) and is abundant in neighboring countries of the Mekong region, is the most common ground on which local poultry are found wandering and the typical soil found at the sites of HPAI virus (H5N1) outbreaks. This sandy topsoil is acidic and poorly buffered, which explains the differences observed between our indirect pH measures and the direct measures reported in specialized literature (10). The soil's low pH may inactivate enveloped viral particles, as well as bacteria (online Technical Appendix Table).

In Cambodia, as in several other countries affected by HPAI virus (H5N1), decontamination of the environment after an outbreak is recommended by authorities; for example, disinfectants such as cresols are sprayed over environmental surfaces. However, because of resource limitations, only limited areas can be treated. Our results provide evidence that, even when abundantly

contaminated, some soil types are unlikely to allow transmission of the virus to poultry and, consequently, probably not to other animals or to humans. These results suggest that limited resources could be better concentrated in high-risk areas, where the nature of the soils would be more likely to lead to poultry infection after natural contamination. These data may aid in the design of more cost-effective and solid-based decontamination measures for preventing transmission of HPAI virus (H5N1) to humans and animals.

This work was supported by the Office of the Assistant Secretary for Preparedness and Response of the US Department of Health and Human Services.

Ramona A. Gutiérrez and Philippe Buchy

Author affiliation: Institut Pasteur in Cambodia, Phnom Penh, Cambodia

DOI: <http://dx.doi.org/10.3201/eid1809.120402>

References

- World Organisation for Animal Health. Outbreaks of highly pathogenic avian influenza (subtype H5N1) in poultry— from the end of 2003 to 7 May 2012 [cited 2012 May 24]. http://www.oie.int/fileadmin/Home/eng/Animal_Health_in_the_World/docs/pdf/graph_avian_influenza/graphs_HPAI_07_05_2012.pdf
- World Health Organization. Cumulative number of confirmed human cases for avian influenza A(H5N1) reported to WHO, 2003–2012—as of 02 May 2012 [cited 2012 May 24]. http://www.who.int/influenza/human_animal_interface/EN_GIP_20120502CumulativeNumber_H5N1cases.pdf
- Horm SV, Gutiérrez RA, Sorn S, Buchy P. Environment: a potential source of animal and human infection with influenza A (H5N1) virus. *Influenza Other Respi Viruses*. 2012. Epub ahead of print. <http://dx.doi.org/10.1111/j.1750-2659.012.00338.x>
- Indriani R, Samaan G, Gultom A, Loth L, Indryani S, Adjid R, et al. Environmental sampling for avian influenza virus A (H5N1) in live-bird markets, Indonesia. *Emerg Infect Dis*. 2010;16:1889–95.

Table. Clinical, virologic, and serologic results obtained from chickens exposed to soils experimentally contaminated with influenza virus (H5N1)*

Protocol	Sandy topsoil (rice fields)	Building sand	Soil-based compost
Low-dose contamination			
Clinical signs	None	None	None
Mortality rate	None	None	None
Virus in postmortem samples	NA	NA	NA
Seroconversion rate	None	33% at day 24	50% at day 24
High-dose contamination			
Clinical signs	None	None	None
Mortality rate	None	100% by day 4	100% by day 4
Virus in postmortem samples	NA	Yes	Yes
Seroconversion rate	None	NA	NA

*Low-dose contamination protocol, exposure of the chickens to soil inoculated with 1 infectious unit on day 0, 2 on day 6, 4 on day 12, and 8 on day 18. High-dose contamination protocol, exposure of the chickens to soil inoculated with 8 infectious units on day 0, 12 on day 6, 16 on day 12, and 20 on day 18. In each experiment, 10–20 chickens were exposed to contaminated soils. Each experiment was repeated twice. For each experiment, a control group was established (same number of chickens exposed to noncontaminated soils; no deaths were observed in control groups). NA, not applicable.

5. Leung YHC, Zhang L-J, Chow C-K, Tsang C-L, Ng C-F, Wong C-K, et al. Poultry drinking water for avian influenza surveillance. *Emerg Infect Dis.* 2007;13:1380–2. <http://dx.doi.org/10.3201/eid1309.070517>
6. Stallknecht DE, Goekjian VH, Wilcox BR, Poulson RL, Brown JD. Avian influenza virus in aquatic habitats: what do we need to learn? *Avian Dis.* 2010;54(Suppl):461–5. <http://dx.doi.org/10.1637/8760-033109-Reg.1>
7. Vong S, Ly S, Mardy S, Holl D, Buchy P. Environmental contamination during influenza A virus (H5N1) outbreaks, Cambodia, 2006. *Emerg Infect Dis.* 2008;14:1303–5. <http://dx.doi.org/10.3201/eid1408.070912>
8. World Health Organization. Recommendations and laboratory procedures for detection of avian influenza A(H5N1) virus in specimens from suspected human cases. Revised August 2007 [cited 2012 Jan 16]. <http://www.who.int/influenza/resources/documents/RecAllabtestsAug07.pdf>
9. World Organisation for Animal Health. Manual of diagnostics tests and vaccines for terrestrial animals 2010. Chapter 2.3.4: avian influenza [cited 2012 Jan 16]. http://www.oie.int/fileadmin/Home/eng/Health_standards/tahm/2.03.04_AI.pdf
10. Nesbitt HJ. Rice production in Cambodia. Manila: International Rice Research Institute; 1997.

Address for correspondence: Philippe Buchy, Institut Pasteur in Cambodia, Virology Unit, 5 Monivong Blvd, PO Box 983, 12152 Phnom Penh, Cambodia; email: pbuchy@pasteur-kh.org



Rickettsia raoultii-like Bacteria in *Dermacentor* spp. Ticks, Tibet, China

To the Editor: *Rickettsia raoultii* is an obligate intracellular gram-negative bacterium belonging to the spotted fever group (SFG) of the genus *Rickettsia*. Genotypes RpA4, DnS14, and DnS28, originally isolated from ticks from Russia in 1999 (1), were designated as *Rickettsia raoultii* sp. nov. on the basis of phylogenetic analysis (2). *R. raoultii* has been found mainly in *Dermacentor* spp. ticks in several countries in Europe (3). It was detected in a *Dermacentor marginatus* tick from the scalp of a patient with tick-borne lymphadenitis in France (2), which suggests that it might be a zoonotic pathogen. We determined the prevalence of *R. raoultii*-like bacteria in *Dermacentor* spp. in highland regions in Tibet.

Ticks from sheep (*Ovis aries*) near Namuco Lake (a popular tourist destination 4,718 m above sea level) were collected and identified morphologically as *D. everestianus* and *D. niveus* ticks (4). Genomic DNA was extracted from individual specimens by using the QIAamp DNA Mini Kit (QIAGEN, Hilden, Germany). All DNA samples were amplified by using PCRs specific for the citrate synthase (*gltA*, 770 bp) gene (5) and the outer membrane protein A (*ompA*, 629 bp) gene (6). Some samples were amplified by using a PCR specific for the *ompB* (2,479 bp) gene (7).

Randomly selected amplicons for *gltA* (n = 27), *ompA* (n = 31), and *ompB* (n = 7) were cloned into the pGEM-T Easy vector (Promega, Shanghai, China) and subjected to bidirectional sequencing (Sangon Biotech, Shanghai, China). Sequences obtained were deposited in GenBank under accession nos. JQ792101–JQ792105, JQ792107, and JQ792108–

JQ792166. Phylogenetic analysis was conducted for sequences we identified and sequences of recognized SFG rickettsial species available in Genbank by using the MegAlign program (DNASTAR, Inc., Madison, WI, USA) and MEGA 4.0 (8).

Of 874 tick specimens, 86 were *D. everestianus* ticks (13 male and 73 female), and 788 were *D. niveus* ticks (133 male and 655 female). Samples positive for *gltA* and *ompA* were considered SFG rickettsial species. Using this criterion, we found that 739 tick specimens (84.6%) were positive for *Rickettsia* spp. Of 86 *D. everestianus* ticks, 85 (98.8%) were positive for *Rickettsia* spp. and of 788 *D. niveus* ticks, 654 (83.0%) were positive. Infection rates for male and female *D. niveus* ticks were 87.9% and 82.1%, respectively. We found an overall prevalence of 84.6% for *R. raoultii*-like bacteria in *Dermacentor* spp. in the highland regions in Tibet.

Nucleotide sequence identities ranged from 99.2% to 100% (except for isolate WYG55, which had an identity of 98.6%) for the *ompA* gene and from 99.2% to 99.9% (except for isolate XG86, which had an identity of 98.5%) for the *ompB* gene. These results indicated that homology levels of most isolates were within species thresholds (*ompA* ≈98.8% and *ompB* ≈99.2%) (9). Isolate WYG55 showed the lowest identity (98.2%) among *gltA* gene sequences and the lowest identity (98.6%) among *ompA* gene sequences. Isolate XG86 showed lowest identity (98.5%) among *ompB* gene sequences. These results suggest that other *Rickettsia* spp. were among the investigated samples.

A BLASTn search (www.ncbi.nlm.nih.gov/) for the obtained sequences was conducted. The best matches (highest identities) detected were with sequences of *R. raoultii*. However, comparison of our sequences with corresponding sequences of *R. raoultii* in GenBank showed identity ranging from 98.0% to 99.0% for