

Equine Influenza A(H3N8) Virus Isolated from Bactrian Camel, Mongolia

Myagmarsukh Yondon, Batsukh Zayat,
Martha I. Nelson, Gary L. Heil,
Benjamin D. Anderson, Xudong Lin,
Rebecca A. Halpin, Pamela P. McKenzie,
Sarah K. White, David E. Wentworth,
and Gregory C. Gray

Because little is known about the ecology of influenza viruses in camels, 460 nasal swab specimens were collected from healthy (no overt illness) Bactrian camels in Mongolia during 2012. One specimen was positive for influenza A virus (A/camel/Mongolia/335/2012[H3N8]), which is phylogenetically related to equine influenza A(H3N8) viruses and probably represents natural horse-to-camel transmission.

Since the first isolation in 1963 of an avian-origin influenza A(H3N8) virus from horses (1), subtype H3N8 influenza viruses have continued to circulate panzootically among horses, causing severe outbreaks of equine influenza respiratory disease. In the United States, these viruses jumped from horses to dogs and continue to circulate among dogs (2,3). In Mongolia, the site of some of the world's largest epizootics of equine influenza A(H3N8) virus (EIV) infection, transmission of this virus is sustained among 2.1 million free-ranging horses, causing significant economic losses among rural herders. Major epizootics of EIV infection occurred in Mongolia during 2007–2008 (459,000 cases, 24,600 deaths) and again in 2011 (74,608 cases, 40 deaths) (4).

Over previous decades in Mongolia, outbreaks of respiratory disease, thought to be influenza, among camels have been reported. In the 1980s, the virus was characterized, and researchers speculated that it was related to a reassortant influenza A(H1N1) virus vaccine strain,

A/PR-8/34 + A/USSR/77, generated in a Soviet laboratory and administered to humans in Mongolia and possibly transmitted from vaccinated humans to camels in a reactivated form (5,6). However, only 1 genetic sequence from this outbreak among camels is available in GenBank: A/camel/Mongolia/1982/H1N1. Despite reports of serologic activity against influenza A virus among camels in several African countries (7,8), the lack of isolated virus from these populations highlights how little is known about the ecology of influenza viruses in camels. Questions about the potential role of camels in human cases of Middle East respiratory syndrome (9) further highlight our lack of knowledge of infectious diseases in camels and the merits of increased surveillance at this unique human–animal interface.

Since January 2011, surveillance of equine influenza viruses has been enhanced in 3 Mongolian aimags (provinces). Surveillance among camels was also initiated in response to anecdotal reports of signs of respiratory illness in Bactrian camels (*Camelus bactrianus*). We describe the isolation, full-genome sequencing, and phylogenetic characterization of an influenza A(H3N8) virus of equine lineage isolated from a Bactrian camel, thereby identifying a novel route of influenza virus interspecies transmission and raising further questions about influenza A virus ecology in understudied regions such as Mongolia.

The Study

During January–January 2013, a total of 460 nasal swab specimens were collected through active surveillance of horses and camels in 3 Mongolian aimags (Figure 1) known for high densities of free-ranging horses and camels (Table). Specimens were collected monthly, as weather permitted, from 50 free-ranging horses and 20 free-ranging Bactrian camels that were safely and carefully restrained with halters, ropes, and by hand, according to a protocol approved by the Department of Veterinary and Animal Breeding, Government of Mongolia. During sampling, camels were in a crouched or take-down position. Horse and camel specimens were carefully stored and shipped in separate containers; to prevent cross-contamination with EIV, specimens were separated during laboratory analyses.

All specimens were first screened at the Institute of Veterinary Medicine laboratory (Ulaanbaatar, Mongolia) by using the World Health Organization (WHO) influenza A quantitative reverse transcription PCR (qRT-PCR) protocol (10). Six specimens collected from camels without respiratory signs were positive for influenza A and were double-blind passaged in embryonated chicken eggs. Subsequent testing revealed hemagglutination activity in all 6 specimens. Allantoic fluid of the 6 cultured specimens was then shipped to the University of Florida for confirmation testing and sequencing.

Author affiliations: Institute of Veterinary Medicine, Ulaanbaatar, Mongolia (M. Yondon, B. Zayat); National Institutes of Health, Bethesda, Maryland, USA (M.I. Nelson); University of Florida, Gainesville, Florida, USA (G.L. Heil, B.D. Anderson, S.K. White, G.C. Gray); J. Craig Venter Institute, Rockville, Maryland, USA (X. Lin, R.A. Halpin, D.E. Wentworth); and St. Jude Children's Research Hospital, Memphis, Tennessee, USA (P.P. McKenzie)

DOI: <http://dx.doi.org/10.3201/eid2012.140435>



Figure 1. The 3 aimags from which nasal swab specimens were collected from healthy Bactrian camels, for influenza A virus testing, Mongolia, 2012. 1, Töv; 2, Khentii; 3, Dundgovi.

Only 1 specimen was confirmed positive by influenza A-specific qRT-PCR (cycle threshold [C_t] <35), suggesting possible virus degradation during shipment, despite specimens being shipped on dry ice and carefully handled upon receipt (10). The original swab specimen from a camel was later shared with the Mongolia National Influenza Center for confirmation in an anonymized panel of 10 camel swab specimens. Using WHO qRT-PCR procedures, staff identified the specimen as having the strongest evidence (by C_t) of influenza A virus. Staff further studied the specimen with conventional RT-PCR primers and probes for the hemagglutinin and neuraminidase genes. These reactions yielded amplicons of the expected size, which were sequenced and found to be 100% identical to the corresponding portions of the J. Craig Venter Institute (Rockville, MD, USA) sequences described below.

Sanger sequence data for the hemagglutinin and neuraminidase genes of this isolate demonstrated extremely

high levels of identity with recent EIV from Asia isolated under this project in 2011 (4). Full-genome sequencing of the 8 genome segments amplified by multisegment RT-PCR (11) was performed at the J. Craig Venter Institute by using Ion Torrent PGM technology (Life Technologies, Grand Island, NY, USA) with a 314v2 chip, and the sequences were validated by using the MiSeq platform (Illumina, Inc., San Diego, CA, USA). Full-genome sequence data for this virus, named A/camel/Mongolia/335/2012(H3N8), were deposited in GenBank (accession nos. CY164120.1, CY164121.1, CY164122.1, CY164123.1, CY164124.1, CY164125.1, CY164126.1 and CY164127.1). The isolate should soon (summer 2014) become available for other research use through BEI Resources (Manassas, VA, USA).

For each of the 8 viral genome segments, phylogenetic trees were inferred separately by use of the maximum-likelihood methods available in RAxML version 7.2.6 (12), a general time-reversible model of nucleotide

Table. Number of specimens collected from camels, by aimag, each month, and result of testing for influenza A virus, Mongolia, January 2012–January 2013*

Date	Aimag (province)					
	Töv		Khentii		Dundgovi	
	Positive	Total	Positive	Total	Positive	Total
2012						
January	0	20	–	–	–	–
February	–	–	–	–	0	20
March	0	20	–	–	–	–
April	0	20	0	20	–	–
May	0	20	–	–	0	20
June	0	20	–	–	0	20
July	–	–	0	20	0	20
August	0	20	–	–	0	20
September	0	20	–	–	0	20
October	0	20	0	20	–	–
November	–	–	0	20	4	20
December	0	20	–	–	0	20
2013						
January	2	20	0	20	–	–
Total	2	200	0	100	4	160

*Testing was performed by quantitative reverse transcription PCR. A total of 460 specimens were collected across all aimags. – denotes when sampling did not occur because of poor weather or road conditions.




Figure 2. Evolutionary relationships of 155 full-length hemagglutinin sequences from equine A(H3N8)viruses collected globally and A/camel/Mongolia/335/2012 (arrow). The 2 clades associated with most recent equine influenza A(H3N8) viruses, Florida clade 1 and Florida clade 2, are denoted as FC1 and FC2, respectively, and with nomenclature adopted previously (13). The maximum-likelihood tree is midpoint rooted for clarity, and all branch lengths are drawn to scale. High (>70) bootstrap values are provided for key nodes. Hemagglutinin sequences containing a 2aa insertion are identified with a solid black circle. Scale bar indicates nucleotide substitutions per site. An expanded version of this figure is available online (<http://wwwnc.cdc.gov/EID/article/20/12/14-0435-F2.htm>).

substitution, and a gamma-distributed rate variation among sites, with a bootstrapping resampling process (500 replicates). All 8 viral genome segments (polymerase basic protein [PB] 2, PB1, polymerase acidic protein (PA), hemagglutinin (HA), nucleocapsid protein (NP), neuraminidase (NA), matrix protein (MP), and nonstructural protein (NS) are closely related to the equine influenza A(H3N8) viruses that have recently been circulating among the horse population in Asia; this lineage is evolutionarily distinct from the influenza A(H3N8) viruses circulating among birds in Asia (online Technical Appendix Figures 1–7, <http://wwwnc.cdc.gov/EID/article/20/12/14-0435-Techapp1.pdf>). A more detailed phylogenetic analysis of the hemagglutinin segment performed by using the additional background sequence data representing the global diversity of influenza A(H3N8) viruses in horses indicates that A/camel/Mongolia/335/2012 is positioned within Florida clade 2 (FC2) (Figure 2) (13) and, more specifically, within a bootstrap-supported clade that contains 3 influenza A(H3N8) viruses isolated from

horses in Mongolia in 2011: A/equine/Mongolia/6/2011 (100% nt similarity), A/equine/Mongolia/56/2011 (99.9% nt similarity), and A/equine/Mongolia/3/2011 (100% nt similarity). A/camel/Mongolia/335/2012 also contains an insertion of 2 aa (I and F) near the beginning of the hemagglutinin sequence (hemagglutinin positions 8–9, online Technical Appendix) that was first detected among FC2 equine viruses in 2005 (A/equine/Bari/2005/H3N8) and has been detected in most FC2 viruses, including all viruses that are closely related to A/camel/Mongolia/335/2012.

Conclusions

The phylogeny indicates that A/camel/Mongolia/335/2012 probably represents a relatively recent horse-to-camel transmission event. Without additional isolates from camels or corresponding epidemiologic data, and given the close genetic relationship between A/camel/Mongolia/335/2012 and related equine viruses, it is impossible to determine at this time whether the virus has been successfully transmitted from camel to camel.

In recent years, enhanced surveillance has detected influenza A viruses across a wider range of mammalian hosts, including horses, swine, dogs (14), seals (15), cats, and now camels, providing a more complete picture of the ecology of influenza A viruses beyond their presence in birds. How influenza A viruses successfully jump from 1 host species to another, and what the constraints on inter-species transmission are, remain key questions about influenza virus ecology and assessments of pandemic threats. Our findings highlight the need to further elucidate the ecology of influenza viruses and other pathogens in free-ranging camel populations.

Acknowledgments

We thank Badarch Darmaa for her help with validating the influenza culture work.

This research was supported by multiple grants: contracts HH-SN266200700005C (St. Jude Children's Research Hospital) and HHSN272200900007C (J. Craig Venter Institute) from the National Institute of Allergy and Infectious Diseases, National Institutes of Health, Department of Health and Human Services, USA; R01 AI068803-ARRA supplement (G.C.G.) from the National Institute of Allergy and Infectious Diseases; and multiple grants (G.C.G.) from the US Department of Defense Armed Forces Health Surveillance Center, Global Emerging Infections Surveillance and Response Program. This research was conducted within the context of the Multinational Influenza Seasonal Mortality Study, an ongoing international collaborative effort to understand influenza epidemiology and evolution, led by the Fogarty International Center, National Institutes of Health, with funding from the Office of Global Affairs at the Department of Health and Human Services (M.I.N.).

Dr Yondon is a researcher in the Virology Laboratory at the Institute of Veterinary Medicine, Ulaanbaatar, Mongolia. His research interests include the development of new methods and technology for diagnosing, controlling, and preventing viral diseases in animals and their application in veterinary medical practice.

References

1. Waddell GH, Teigland MB, Sigel MM. A new influenza virus associated with equine respiratory disease. *J Am Vet Med Assoc.* 1963;143:587-90.
2. Crawford PC, Dubovi EJ, Castleman WL, Stephenson I, Gibbs EP, Chen L, et al. Transmission of equine influenza virus to dogs. *Science.* 2005;310:482-5. <http://dx.doi.org/10.1126/science.1117950>
3. Hayward JJ, Dubovi EJ, Scarlett JM, Janeczko S, Holmes EC, Parrish CR. Microevolution of canine influenza virus in shelters and its molecular epidemiology in the United States. *J Virol.* 2010;84:12636-45. <http://dx.doi.org/10.1128/JVI.01350-10>
4. Yondon M, Heil GL, Burks JP, Zayat B, Waltzek TB, Jamiyan BO, et al. Isolation and characterization of H3N8 equine influenza A virus associated with the 2011 epizootic in Mongolia. *Influenza Other Respir Viruses.* 2013;7:659-65. <http://dx.doi.org/10.1111/irv.12069>
5. Anchlan D, Ludwig S, Nymadawa P, Mendsaikhan J, Scholtissek C. Previous H1N1 influenza A viruses circulating in the Mongolian population. *Arch Virol.* 1996;141:1553-69. <http://dx.doi.org/10.1007/BF01718254>
6. Yamnikova SS, Mandler J, Bekh-Ochir ZH, Dachtzeren P, Ludwig S, Lvov DK, et al. A reassortant H1N1 influenza A virus caused fatal epizootics among camels in Mongolia. *Virology.* 1993;197:558-63. <http://dx.doi.org/10.1006/viro.1993.1629>
7. Caffar Elamin MA, Kheir SA. Detection of influenza antibody in animal sera from Kassala region, Sudan, by agar gel diffusion test. *Rev Elev Med Vet Pays Trop.* 1985;38:127-9.
8. Olaleye OD, Baba SS, Omolabu SA. Preliminary survey for antibodies against respiratory viruses among slaughter camels (*Camelus dromedarius*) in north-eastern Nigeria [in French]. *Revue Scientifique et Technique de l'OIE.* 1989;8:779-83.
9. Haagmans BL, Al Dhahiry SH, Reusken CB, Raj VS, Galiano M, Myers R, et al. Middle East respiratory syndrome coronavirus in dromedary camels: an outbreak investigation. *Lancet Infect Dis.* 2014;14:140-5. [http://dx.doi.org/10.1016/S1473-3099\(13\)70690-X](http://dx.doi.org/10.1016/S1473-3099(13)70690-X)
10. World Health Organization. WHO information for molecular diagnosis of influenza virus in humans—update. 2011 [cited 2013 Jun 10]. http://www.who.int/influenza/resources/documents/molecular_diagnosis_influenza_virus_humans_update_201108.pdf
11. Zhou B, Donnelly ME, Scholes DT, St George K, Hatta M, Kawaoka Y, et al. Single-reaction genomic amplification accelerates sequencing and vaccine production for classical and swine origin human influenza A viruses. *J Virol.* 2009;83:10309-13. <http://dx.doi.org/10.1128/JVI.01109-09>
12. Stamatakis A. RAXML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics.* 2006;22:2688-90. <http://dx.doi.org/10.1093/bioinformatics/btl446>
13. Murcia PR, Wood JL, Holmes EC. Genome-scale evolution and phylodynamics of equine H3N8 influenza A virus. *J Virol.* 2011;85:5312-22. <http://dx.doi.org/10.1128/JVI.02619-10>
14. Song D, Moon HJ, An DJ, Jeoung HY, Kim H, Yeom MJ, et al. A novel reassortant canine H3N1 influenza virus between pandemic H1N1 and canine H3N2 influenza viruses in Korea. *J Gen Virol.* 2012;93:551-4. <http://dx.doi.org/10.1099/vir.0.037739-0>
15. Anthony SJ, St Leger JA, Pugliarès K, Ip HS, Chan JM, Carpenter ZW, et al. Emergence of fatal avian influenza in New England harbor seals. *mBio.* 2012;3:e00166-12.

Address for correspondence: Gregory C. Gray, College of Public Health and Health Professions, University of Florida, Box 100188, Gainesville, FL 32610, USA; email: gcgray@php.ufl.edu



GovDelivery

Manage your email alerts so you only receive content of interest to you.

Sign up for an Online Subscription:
<http://wwwnc.cdc.gov/eid/subscribe.htm>

Equine Influenza A(H3N8) Virus Isolated from Bactrian Camel, Mongolia

Technical Appendix

Technical Appendix Table 1. Virus name, subtype, host species, collection date, and GenBank accession number (PB2 segment) for the 36 H3N8 viruses used in the whole-genome phylogenetic analysis conducted for all eight viral genome segments.

Virus name	Subtype	Host species	Collection date	Accession no.
A_camel_Mongolia_335_2012	H3N8	Camel	Nov-2012	CY164127.1
A_avian_Japan_8KI0102_2008	H3N8	Avian	Oct-08-08	CY079266
A_avian_Japan_8KI0129_2008	H3N8	Avian	Oct-08-08	CY079258
A_avian_Japan_8KI0150_2008	H3N8	Avian	Oct-08-08	CY079242
A_avian_Japan_8KI0162_2008	H3N8	Avian	Oct-08-08	CY079234
A_chicken_Laos_A0573_2007	H3N8	Avian	2007	CY040963
A_chicken_Vietnam_G14_2008	H3N8	Avian	Jan-2008	AB593452
A_donkey_Xinjiang_5_2007	H3N8	Equine	Dec-2007	EU794572
A_duck_Beijing_40_04	H3N8	Avian	2004	EU492488
A_duck_Beijing_61_05	H3N8	Avian	2005	EU492492
A_duck_Hokkaido_8_1980	H3N8	Avian	1980	AB274963
A_duck_Hunan_S1256_2012	H3N8	Avian	Mar-23-12	CY146601
A_duck_Hunan_S1824_2012	H3N8	Avian	Mar-24-12	CY146625
A_duck_Nanchang_1681_1992	H3N8	Avian	Dec-01-92	CY005475
A_duck_Vietnam_G119_2006	H3N8	Avian	Nov-2006	AB593428
A_environment_Hunan_S4350_2011	H3N8	Avian	Nov-13-11	CY146753
A_equine_Gansu_7_2008	H3N8	Equine	Jan-2008	EU794492
A_equine_Heilongjiang_1_2010_	H3N8	Equine	Apr-23-10	KF309031
A_equine_Heilongjiang_10_2008	H3N8	Equine	Apr-2008	EU794508
A_equine_Huabei_1_2007	H3N8	Equine	Dec-03-07	GU571147
A_equine_Inner_Mongolia_8_2008	H3N8	Equine	Feb-2008	EU794524
A_equine_Kyonggi_SA1_2011	H3N8	Equine	Jul-01-11	JX844143
A_equine_Liaoning_9_2008	H3N8	Equine	Apr-2008	EU794516
A_equine_Qinghai_1_1994	H3N8	Equine	1994	EU794532
A_equine_Sachiyama_1_1971	H3N8	Equine	1971	CY034941
A_equine_Tokyo_2_1971	H3N8	Equine	1971	CY096922
A_equine_Tottori_1_07	H3N8	Equine	2007	AB591847
A_equine_Xinjiang_1_2007	H3N8	Equine	Nov-2007	EU794540
A_equine_Xinjiang_2_2007	H3N8	Equine	Nov-2007	EU794548
A_equine_Xinjiang_3_2007	H3N8	Equine	Nov-2007	EU794556
A_equine_Xinjiang_4_2007	H3N8	Equine	Dec-2007	EU794564

Technical Appendix Table 1. Virus name, subtype, host species, collection date, and GenBank accession number (PB2 segment) for the 36 H3N8 viruses used in the whole-genome phylogenetic analysis conducted for all eight viral genome segments.

Virus name	Subtype	Host species	Collection date	Accession no.
A_equine_Xuzhou_01_2013	H3N8	Equine	Aug-27-13	KF806992
A_Mallard_SanJiang_90_2006_2006	H3N8	Avian	2006	CY100631
A_muscovy_duck_Vietnam_LBM240_2012	H3N8	Avian	2012	AB786912
A_swine_Anhui_01_2006	H3N8	Swine	Jan-06-06	FJ200417
A_swine_Chibi_01_2005	H3N8	Swine	Dec-15-05	FJ200425

Technical Appendix Table 2. Virus names for the hemagglutinin sequences of the 155 equine A/H3N8 viruses used in Figure 1. Viruses containing the two amino acid insertion near the beginning of the hemagglutinin are specified.

Virus name	Subtype	Insertion
A_camel_Mongolia_335_2012	H3N8	yes
A_donkey_Xinjiang_5_2007	H3N8	
A_equine_Alaska_29759_1991	H3N8	
A_equine_Algers_1_1972	H3N8	
A_equine_Almaty_26_2007	H3N8	yes
A_equine_Argentina_1_93	H3N8	
A_equine_Austria_421_1992	H3N8	
A_equine_Bari_2005	H3N8	yes
A_equine_Berlin_1_1989	H3N8	
A_equine_California_1_1980	H3N8	
A_equine_California_103_1982	H3N8	
A_equine_California_191_2003	H3N8	
A_equine_California_4537_1997	H3N8	
A_equine_California_83_1982	H3N8	
A_equine_California_8560_2002	H3N8	
A_equine_Carlow_1_2009	H3N8	
A_equine_Colorado_10_2007	H3N8	
A_equine_Cordoba_18_1985	H3N8	
A_equine_Donegal_1_2007	H3N8	yes
A_equine_Donegal_1_2009	H3N8	
A_equine_Down_1_2008	H3N8	yes
A_equine_Egypt_6066NAMRU3-VSVRI_2008	H3N8	
A_equine_Florida_1_93	H3N8	
A_equine_Florida_612_2004	H3N8	
A_equine_Florida_779_2004	H3N8	
A_equine_Fontainbleu_1_1979	H3N8	
A_equine_Fontainebleau_1976	H3N8	
A_equine_France_1_1976	H3N8	
A_equine_Gansu_7_2008	H3N8	
A_equine_Georgia_1_1981	H3N8	
A_equine_Georgia_10_1981	H3N8	

Technical Appendix Table 2. Virus names for the hemagglutinin sequences of the 155 equine A/H3N8 viruses used in Figure 1. Viruses containing the two amino acid insertion near the beginning of the hemagglutinin are specified.

Virus name	Subtype	Insertion
A_camel_Mongolia_335_2012	H3N8	yes
A_equine_Georgia_13_1981	H3N8	
A_equine_Georgia_3_1981	H3N8	
A_equine_Georgia_9_1981	H3N8	
A_equine_Guelph_06-28865_2006	H3N8	
A_equine_Guelph_G03-0250_2003	H3N8	
A_equine_Guelph_G03-55399_2003	H3N8	
A_equine_Guelph_G04-54701_2004	H3N8	
A_equine_Heilongjiang_1_2010	H3N8	yes
A_equine_Heilongjiang_10_2008	H3N8	
A_equine_Hokkaido_I828_2008	H3N8	
A_equine_Hong_Kong_J_1992	H3N8	
A_equine_Huabei_1_2007	H3N8	yes
A_equine_Hubei_6_2008	H3N8	
A_equine_Ibadan_6_91	H3N8	
A_equine_Ibadan_9_91	H3N8	
A_equine_Ibaraki_1_07	H3N8	
A_equine_Idaho_37875_1991	H3N8	
A_equine_Inner_Mongolia_8_2008	H3N8	
A_equine_Italy_1062_1991	H3N8	
A_equine_Italy_1199_1992	H3N8	
A_equine_Italy_824_1991	H3N8	
A_equine_Johannesburg_1_1986	H3N8	
A_equine_Kanazawa_1_2007	H3N8	
A_equine_Kascakew_1_1978	H3N8	
A_equine_Katra-Jammu_6_2008	H3N8	yes
A_equine_Kentucky_1_1981	H3N8	
A_equine_Kentucky_1_1986	H3N8	
A_equine_Kentucky_1_1987	H3N8	
A_equine_Kentucky_1_1990	H3N8	
A_equine_Kentucky_1_1991	H3N8	
A_equine_Kentucky_1_1992	H3N8	
A_equine_Kentucky_1_1994	H3N8	
A_equine_Kentucky_1277_1990	H3N8	
A_equine_Kentucky_2_1980	H3N8	
A_equine_Kentucky_2_1981	H3N8	
A_equine_Kentucky_2_1986	H3N8	
A_equine_Kentucky_2_1987	H3N8	
A_equine_Kentucky_211_1987	H3N8	
A_equine_Kentucky_3_1981	H3N8	

Technical Appendix Table 2. Virus names for the hemagglutinin sequences of the 155 equine A/H3N8 viruses used in Figure 1. Viruses containing the two amino acid insertion near the beginning of the hemagglutinin are specified.

Virus name	Subtype	Insertion
A_camel_Mongolia_335_2012	H3N8	yes
A_equine_Kentucky_3_1986	H3N8	
A_equine_Kentucky_4_1980	H3N8	
A_equine_Kentucky_5_2002	H3N8	
A_equine_Kentucky_692_1988	H3N8	
A_equine_Kentucky_694_1988	H3N8	
A_equine_Kentucky_698_1988	H3N8	
A_equine_Kentucky_8_1994	H3N8	
A_equine_Kentucky_bitter_boredom5_1976	H3N8	
A_equine_Kentucky_magnificent_genius1_1981	H3N8	
A_equine_Kentucky_pass_the_pepper1_1976	H3N8	
A_equine_Kentucky_Rosie100_1981	H3N8	
A_equine_Kildare_1_2007	H3N8	yes
A_equine_Kyonggi_SA1_2011	H3N8	
A_equine_Liaoning_9_2008	H3N8	
A_equine_Limerick_1_2010	H3N8	
A_equine_Lincolnshire_1_2007	H3N8	
A_equine_Lonquen_1_2006	H3N8	
A_equine_Massachussetts_213_2003	H3N8	
A_equine_Miami_1_1963	H3N8	
A_equine_Mongolia_20_2011	H3N8	yes
A_equine_Mongolia_3_2011	H3N8	yes
A_equine_Mongolia_56_2011	H3N8	yes
A_equine_Mongolia_6_2011	H3N8	yes
A_equine_Montana_9233_2007	H3N8	
A_equine_Mysore_1_2008	H3N8	
A_equine_New_Market_1_1979	H3N8	
A_equine_New_Market_1976	H3N8	
A_equine_New_Market_nasalwash1_1979	H3N8	
A_equine_New_York_1_1975	H3N8	
A_equine_New_York_1_1999	H3N8	
A_equine_New_York_146066_2007	H3N8	
A_equine_New_York_452_2003	H3N8	
A_equine_New_York_VR-297_1983	H3N8	
A_equine_Newmarket_5_2003	H3N8	
A_equine_Ohio_1_2003	H3N8	
A_equine_Ohio_113461-1_2005	H3N8	
A_equine_Ohio_113461-2_2005	H3N8	
A_equine_Ohio_113461-3_2005	H3N8	
A_equine_Otar_764_2007	H3N8	yes

Technical Appendix Table 2. Virus names for the hemagglutinin sequences of the 155 equine A/H3N8 viruses used in Figure 1. Viruses containing the two amino acid insertion near the beginning of the hemagglutinin are specified.

Virus name	Subtype	Insertion
A_camel_Mongolia_335_2012	H3N8	yes
A_equine_Qinghai_1_1994	H3N8	
A_equine_Richmond_1_2007	H3N8	yes
A_equine_Romania_1_1980	H3N8	
A_equine_Rome_5_1991	H3N8	
A_equine_Rook_93753_1989	H3N8	
A_equine_Sachiyama_1_1971	H3N8	
A_equine_Santa_Fe_1_1985	H3N8	
A_equine_Sao_Paulo_1_1969	H3N8	
A_equine_Sao_Paulo_6_1963	H3N8	
A_equine_Spain_1_2007	H3N8	yes
A_equine_Suffolk_89	H3N8	
A_equine_Sussex_1_1989	H3N8	
A_equine_Switzerland_1118_1979	H3N8	
A_equine_Switzerland_173_1993	H3N8	
A_equine_Sydney_6085_2007	H3N8	
A_equine_Taby_1991	H3N8	
A_equine_Tennessee_5_1985	H3N8	
A_equine_Tennessee_5_1986	H3N8	
A_equine_Texas_117793_2005	H3N8	
A_equine_Texas_39655_1991	H3N8	
A_equine_Tiaret_1_2011	H3N8	yes
A_equine_Tiaret_10_2011	H3N8	yes
A_equine_Tiaret_2_2011	H3N8	yes
A_equine_Tiaret_3_2011	H3N8	yes
A_equine_Tiaret_4_2011	H3N8	yes
A_equine_Tiaret_5_2011	H3N8	yes
A_equine_Tiaret_6_2011	H3N8	yes
A_equine_Tiaret_7_2011	H3N8	yes
A_equine_Tiaret_8_2011	H3N8	yes
A_equine_Tiaret_9_2011	H3N8	yes
A_equine_Tokyo_1971	H3N8	
A_equine_Tokyo_2_1971	H3N8	
A_equine_Tottori_1_07	H3N8	
A_equine_Uruguay_1_1963	H3N8	
A_equine_Virginia_131054-3_2005	H3N8	
A_equine_Wisconsin_1_03	H3N8	
A_equine_Xinjiang_1_2007	H3N8	
A_equine_Xinjiang_2_2007	H3N8	
A_equine_Xinjiang_3_2007	H3N8	


Technical Appendix Table 2. Virus names for the hemagglutinin sequences of the 155 equine A/H3N8 viruses used in Figure 1. Viruses containing the two amino acid insertion near the beginning of the hemagglutinin are specified.

Virus name	Subtype	Insertion
A_camel_Mongolia_335_2012	H3N8	yes
A_equine_Xinjiang_4_2007	H3N8	
A_equine_Xuzhou_01_2013	H3N8	yes
A_equine_Yokohama_aq13_2010	H3N8	yes
A_equine_Yokohama_aq19_2009	H3N8	
A_equine_Yokohama_aq29_2011	H3N8	
A_equine_Yokohama_aq5_2011	H3N8	
A_equine_Yokohama_aq53_2011	H3N8	
A_equine_Yokohama_aq79_2011	H3N8	

Technical Appendix Table 3. Position of the two amino acid insertion at the beginning of the hemagglutinin sequence.


Example of virus with insertion:	
A/equine/Almaty/26/2007/H3N8	MKTTIIFIFILLTHW
Example of virus without insertion:	
A/equine/Alaska/29759/1991/H3N8	MKTTIIL--ILLTHW

PB2




Technical Appendix Figure 1. Evolutionary relationships of the PB2 segment of 36 influenza A viruses of the H3N8 subtype collected in Asia from horses (n = 17), avian species (n = 16), swine (n = 2), and a camel (*A/camel/Mongolia/335/2012(H3N8)*, highlighted in bold). The tree is midpoint rooted for clarity, and all branch lengths are drawn to scale. High bootstrap values (> 70) are provided for key nodes. Scale bar indicates nucleotide substitutions per site.

PB1




Technical Appendix Figure 2. Evolutionary relationships of the PB1 segment of 36 influenza A viruses of the H3N8 subtype collected in Asia from horses (n = 17), avian species (n = 16), swine (n = 2), and a camel (*A/camel/Mongolia/335/2012(H3N8)*, highlighted in bold). The tree is midpoint rooted for clarity, and all branch lengths are drawn to scale. High bootstrap values (> 70) are provided for key nodes. Scale bar indicates nucleotide substitutions per site.

PA




Technical Appendix Figure 3. Evolutionary relationships of the PA segment of 36 influenza A viruses of the H3N8 subtype collected in Asia from horses (n = 17), avian species (n = 16), swine (n = 2), and a camel (*A/camel/Mongolia/335/2012*(H3N8), highlighted in bold). The tree is midpoint rooted for clarity, and all branch lengths are drawn to scale. High bootstrap values (> 70) are provided for key nodes. Scale bar indicates nucleotide substitutions per site.

NP




Technical Appendix Figure 4. Evolutionary relationships of the NP segment of 36 influenza A viruses of the H3N8 subtype collected in Asia from horses (n = 17), avian species (n = 16), swine (n = 2), and a camel (*A/camel/Mongolia/335/2012(H3N8)*, highlighted in bold). The tree is midpoint rooted for clarity, and all branch lengths are drawn to scale. High bootstrap values (> 70) are provided for key nodes. Scale bar indicates nucleotide substitutions per site.



0.02

NS



Technical Appendix Figure 7. Evolutionary relationships of the NS segment of 36 influenza A viruses of the H3N8 subtype collected in Asia from horses (n = 17), avian species (n = 16), swine (n = 2), and a camel (A/camel/Mongolia/335/2012(H3N8), highlighted in bold). The tree is midpoint rooted for clarity, and all branch lengths are drawn to scale. High bootstrap values (> 70) are provided for key nodes.