

Sika Deer Carrying *Babesia* Parasites Closely Related to *B. divergens*, Japan

To the Editor: Human babesiosis caused by *Babesia divergens*, a zoonotic pathogen of bovines in Europe, is an emerging tickborne disease (1). In the United States, a closely related *Babesia* sp. was identified in persons in Missouri and Kentucky and in eastern cottontail rabbits (*Sylvilagus floridanus*) on Nantucket Island, Massachusetts (2–5). We report that sika deer (*Cervus nippon*) in Japan also carry parasites genetically closely related to *B. divergens*.

During November 2007–February 2008 and November 2008–February 2009, we collected blood samples from 96 wild sika deer throughout Japan. We purified DNA from the blood, which had been stored in a freezer, by phenol/chloroform extraction and performed nested PCR for *B. divergens* 18S rRNA (*rDNA*), chaperonin-containing T-complex protein 1 eta subunit (*CCT7*, formerly described as CCT η) (6), and β -*tubulin* (7) genes. Primers for *rDNA* were designed from the sequences of related *Babesia* spp. (GenBank accession nos. U16370, U16369, and AY046575): dv101F (5'-ACAA-CAGTTATAGTTTCTTTG-GTATTTCG-3') and dv1353R (5'-GCCTTAAACTTCCTTGCG-GCTTAGAGC-3'), and dv159F (5'-GCTAATGCAAGTTCGAG-GCCTTTTGCG-3') and dv1296R (5'-CGGACGAACCTTTTACG-GACTAG-3') for the first and second rounds, respectively. *CCT7* primers were similarly designed (GenBank accession nos. AB367924 and AB367925): Bdiv/odoCCTF1 (5'-CAAAATGAGYCACCTMCT-CAACCTACC-3') and Bdiv/CCTR1 (5'-ATCTCAGCAGCTCACTA-CAGTGACCACCTC-3'), and Bdiv/odoCCTF2 (5'-CAACCTACCRAT-TCTCCTYYTGAAGGAGGG-3')

and Bdiv/CCTR2 (5'-GGCTAATA-AGTCCGATATTGCGGGGCT-CACG-3') for the first and second rounds, respectively. The β -*tubulin* PCR protocol has been described (7).

Of the 96 blood samples, 12 from 5 prefectures (Hokkaido, Iwate, Tochigi, Nagano, and Miyazaki) were positive for *Babesia rDNA* (online Technical Appendix, <http://wwwnc.cdc.gov/EID/article/20/8/13-0061-Techapp1.pdf>). The sequence for sample 08–22 from Hokkaido (GenBank accession no. KC465978) was distinct from sequences of the other 11 samples (97.5%–97.6% identity in 1,041 bp), which consisted of 7 variant sequences (GenBank accession nos. KC465973–7 and AB857845–6) and 5 identical sequences (GenBank accession nos. KC465977 and AB861504–7) (99.7%–100% identity). The 5 identical sequences varied in only 1 of 909 bp from *B. divergens rDNA* from an *Ixodes persulcatus* tick in Russia [GenBank accession no. GU057385] (8).

β -*tubulin* (900 bp) was also amplified from the 12 *Babesia rDNA*-positive samples. Ten of the sequences consisted of 3 sequence variants (99.9% identity; GenBank accession nos. KC465969, KC465970, and KC465968/AB861508–14). The 2 divergent sequences (GenBank accession nos. KC465971 [08–22] and KC465989 [08–25]) were most similar to *B. odocoilei* (GenBank accession no. KC465972; 91% identity) and *Theileria orientalis* (GenBank accession no. AP011947; 79.9% identity), respectively. Thus, at least 1 deer likely had *Babesia* and *Theileria* spp. infections.

CCT7 was amplified from 10 of the 12 *Babesia*-positive blood samples. The sequences (GenBank accession nos. KC465979–88) were more heterogeneous (98.7%–99.9% identity) than those for *rDNA* and β -*tubulin*; this finding was expected because *CCT7* evolves more quickly (6).

We generated a neighbor-joining phylogenetic tree (ClustalW,

<http://clustalw.ddbj.nig.ac.jp/index.php?lang=ja>) from the *Babesia rDNA* sequences from our study and from GenBank (Figure, panel A). The tree shows a distinct lineage (Asian) cluster for the deer parasites, except for 08–22 (GenBank accession no. KC465978), within a clade also holding the *B. divergens* strains (human and bovine) from Europe (European Union lineage). The tree also shows a cluster encompassing *Babesia* spp. (human and rabbit) from the United States; *B. divergens* (deer), *B. capreoli* (deer), and *Babesia* sp. (chamois) from Europe; and *B. divergens* (human) from Portugal. Sequence 08–22 branches with *Babesia* spp. in *Ix. ovatus* ticks from Japan (GenBank accession nos. AY190123 and AY190124) (9). The branch lengths indicate clear separation between the isolates from sika deer and ticks, suggesting that the clustering may be attributable to the limited number of available related sequences.

We also generated a phylogenetic tree of β -*tubulin* sequences (900 bp), which produced similar topology and high bootstrap support (Figure, panel B). However, the limited number of relevant β -*tubulin* sequences precludes conclusions regarding the phylogeny of *Babesia* parasites. A *CCT7* phylogenetic tree was not generated because of the paucity of sequences. Overall, the phylogenies suggest that *B. divergens*- and *B. capreoli*-related parasites are found worldwide in temperate zones of the Northern Hemisphere, including Europe (1), the United States (2–5), Russia (8), and Japan.

We showed the presence of *B. divergens*-like *rDNA*, β -*tubulin*, and *CCT7* genes in sika deer from different Japanese prefectures, confirming the presence of this parasite in Japan. *B. capreoli*, which is serologically indistinguishable from *B. divergens*, was previously reported in sika deer (10). However, no molecular data for the *B. capreoli* isolate exist, so no




Figure. Neighbor-joining phylogenetic trees generated from *Babesia* sequences from GenBank and from our study in Japan, November 2007–February 2008 and November 2008–February 2009. Bootstrap support (1,000 repetitions) is indicated at the nodes. The trees are based on the partial (1,041 bp) *Babesia* 18S rRNA gene (A) and the partial (900 bp) β -tubulin gene (B). Sequences determined in this study are in boldface. Scale bar indicates the inferred number of substitutions per site. Lineages are indicated. CDS, coding DNA sequence.

conclusion may be drawn regarding its relationship to the *B. divergens*-like parasites from sika deer in our study.

There is an overabundance of wild sika deer in Japan because these animals easily adapt to a variety of climates, vegetation, and geography. Increased human exposure to deer habitats increases the risk of exposure to tickborne zoonoses, such as those caused by *Babesia* spp. In humans, infections caused by *B. divergens* and *B. divergens*-like parasites can be life threatening; fatality rates of 42% and 33% have been reported in infected asplenic patients in Europe and the United States, respectively (1–3). The findings from our study emphasize the need for increased clinical awareness of babesiosis in Japan and globally. They also emphasize

the need for the swift diagnosis of suspected cases and prompt treatment of confirmed cases, especially in asplenic patients at high risk for the potentially deadly consequences of babesiosis.

Acknowledgments

We are grateful to Danny H.-K. Jokl for his critical reading of the manuscript. We thank Jeremy Gray for providing genomic DNA of *B. divergens* MRNK and Marek Sawczuk and Bogumiła Skotarczak for providing genomic DNA of *B. divergens* from *Ix. ricinus* tick collected in Poland.

Financial support for this study was provided in part by grants-in-aid from the Ministry of Education, Science and Culture of Japan and by the Ministry of

Health, Labor and Welfare Science of Japan (H25-Shinko-Ippan-008 and H25-Shinko-Shitei-009). Support was also provided by a Grant-in-Aid for Cooperative Research from Rakuno-Gakuen University and by Gakujutsu Frontier Cooperative Research in Rakuno-Gakuen University.

**Aya Zamoto-Niikura,
Masayoshi Tsuji,¹
Koichi Imaoka,
Masanobu Kimura,
Shigeru Morikawa,
Patricia J. Holman,
Haruyuki Hirata,
and Chiaki Ishihara**

Author affiliations: National Institute of Infectious Diseases, Tokyo, Japan (A. Zamoto-Niikura, K. Imaoka, M. Kimura, S. Morikawa); Rakuno-Gakuen University, Hokkaido, Japan (A. Zamoto-Niikura, M. Tsuji, H. Hirata, C. Ishihara); and Texas A&M University, College Station, Texas, USA (P.J. Holman)

DOI: <http://dx.doi.org/10.3208/eid2008.130061>

References

- Gorenflot A, Moubri K, Precigout E, Carcy B, Schetters TP. Human babesiosis. *Ann Trop Med Parasitol*. 1998;92:489–501. <http://dx.doi.org/10.1080/00034989859465>
- Herwaldt B, Persing DH, Precigout EA, Goff WL, Mathiesen DA, Taylor PW, et al. A fatal case of babesiosis in Missouri: identification of another piroplasm that infects humans. *Ann Intern Med*. 1996;124:643–50. <http://dx.doi.org/10.7326/0003-4819-124-7-199604010-00004>
- Beattie JF, Michelson ML, Holman PJ. Acute babesiosis caused by *Babesia divergens* in a resident of Kentucky. *N Engl J Med*. 2002;347:697–8. <http://dx.doi.org/10.1056/NEJM200208293470921>
- Goethert HK, Telford SR III. Zoonotic transmission of *Babesia divergens* among cottontail rabbits on Nantucket Island, Massachusetts. *Am J Trop Med Hyg*. 2003;69:455–60.
- Spencer AM, Goethert HK, Telford SR III, Holman PJ. In vitro host erythrocyte specificity and differential morphology of *Babesia divergens* and a zoonotic *Babesia* sp. from eastern cottontail rabbits (*Sylvilagus floridanus*). *J Parasitol*. 2006;92:333–40. <http://dx.doi.org/10.1645/GE-662R.1>

¹Deceased.

6. Nakajima R, Tsuji M, Oda K, Zamoto-Niikura A, Wei Q, Kawabuchi-Kurata T, et al. *Babesia microti*-group parasites compared phylogenetically by complete sequencing of the *CCTeta* gene in 36 isolates. *J Vet Med Sci.* 2009;71:55–68. <http://dx.doi.org/10.1292/jvms.71.55>
7. Hirata H, Kawai S, Maeda M, Jinnai M, Fujisawa K, Katakai Y, et al. Identification and phylogenetic analysis of Japanese macaque *Babesia*-1 (JM-1) detected from a Japanese macaque (*Macaca fuscata fuscata*). *Am J Trop Med Hyg.* 2011;85:635–8. <http://dx.doi.org/10.4269/ajtmh.2011.11-0035>
8. Rar VA, Epikhina TI, Livanova NN, Panov VV. Genetic diversity of *Babesia* in *Ixodes persulcatus* and small mammals from North Ural and West Siberia, Russia. *Parasitology.* 2011;138:175–82. <http://dx.doi.org/10.1017/S0031182010001162>
9. Inokuma H, Yoshizaki Y, Shimada Y, Sakata Y, Okuda M, Onishi T. Epidemiological survey of *Babesia* species in Japan performed with specimens from ticks collected from dogs and detection of new *Babesia* DNA closely related to *Babesia odocoilei* and *Babesia divergens* DNA. *J Clin Microbiol.* 2003;41:3494–8. <http://dx.doi.org/10.1128/JCM.41.8.3494-3498.2003>
10. Gray JS, Murphy TM, Waldrup KA, Wagner GG, Blewett DA, Harrington R. Comparative studies of *Babesia* spp. from white-tailed and sika deer. *J Wildl Dis.* 1991;27:86–91. <http://dx.doi.org/10.7589/0090-3558-27.1.86>


Address for correspondence: Aya Zamoto-Niikura, Division of Experimental Animal Research, National Institute of Infectious Diseases, 4-7-1 Gakuen, Musashimurayama, Tokyo 208-0011, Japan; email: zamoto@nih.go.jp

How to Correctly Remove a Tick

Grasp the tick firmly and as closely to the skin as possible. With a steady motion, pull the tick's body away from the skin.

Do not be alarmed if the tick's mouthparts remain in the skin.

Cleanse the area with an antiseptic.



For more information please contact:
Centers for Disease Control and Prevention
 1600 Clifton Road NE, Atlanta, GA 30333
 Telephone: 1-800-CDC-INFO (232-4636)
 TTY: 1-888-232-63548
 Web: www.cdc.gov/Lyme

Transcontinental Movement of Asian Genotype Chikungunya Virus

To the Editor: Chikungunya virus (CHIKV), a mosquito-transmitted virus (family *Togaviridae*, genus *Alphavirus*), was first isolated >60 years ago in Africa and is responsible for epidemics of acute polyarthralgia. During CHIKV epidemics, the transmission cycle is from humans to mosquitoes, with no intervening amplifying host, and the virus can rapidly disseminate, infecting large numbers of persons. Epidemics have been described in Africa, the Middle East, Europe, India, and Southeast Asia. On the basis of detailed clinical descriptions of the disease, chikungunya fever, it appears that CHIKV caused epidemics in the Caribbean (St. Thomas, US Virgin Islands) and the southeastern coastal United States during the early 19th century (1).

Genetic studies show that the virus has evolved into 3 distinct genotypes: West African, East/Central/South African (ECSA), and Asian (2). The genotypes likely indicate independent evolution of the virus in historically isolated areas. Phenotypic differences have been described between genotypes and between individual strains, most notably an E1 mutation among some ECSA strains, which facilitates replication in *Aedes albopictus* mosquitoes (3). However, more recently, the movement of virus genotypes has increased dramatically, probably as a direct result of increased movement of humans and increased commercial trade. Beginning in 2005 and through 2006, the ECSA genotype virus was responsible for an explosive epidemic, during which the virus moved from coastal Kenya to islands adjacent to southeastern Africa and then to India, where >1 million cases were recorded (2). During this time, imported cases were reported worldwide, and in some


instances, autochthonous transmission was detected in distal locations (4,5).

In October 2013, the arbovirus diagnostic laboratory at the Centers for Disease Control and Prevention (CDC; Fort Collins, CO, USA) detected CHIKV in human serum specimens from Yap State, Federated States of Micronesia; the specimens were collected during an epidemic of disease clinically compatible with chikungunya fever. In December 2013, the French National Reference Centre for arboviruses confirmed that CHIKV was responsible for an epidemic occurring on St. Martin Island, French West Indies, in the Caribbean (6). In January 2014, the Caribbean Public Health Agency detected CHIKV in 2 human serum specimens from the British Virgin Islands (BVI); the samples were subsequently confirmed by CDC to be positive for CHIKV.

By using next-generation sequencing, we determined the complete nucleotide sequence for 1 of the CHIKV specimens detected in BVI and for 2 of the CHIKV specimens detected in Yap. DNA libraries for next-generation sequencing were prepared directly from RNA extracted from serum, and the amplified libraries were sequenced by using the Ion Torrent Personal Genome Machine (Life Technologies, Grand Island, NY, USA). The CLC Genomics Workbench (CLC bio, Aarhus, Denmark) and Lasergene NextGen (DNASTAR, Madison, WI, USA) were used to analyze and assemble raw sequence reads. ClustalW (<http://www.ebi.ac.uk/Tools/msa/clustalw2/>) was used to align the complete genome sequences with a variety of CHIKV sequences, representing the 3 genotypes, from GenBank. Nearly identical phylogenetic trees were generated by several methods (i.e., minimum evolution, maximum likelihood, neighbor joining); a representative neighbor-joining tree generated and analyzed with 1,000 replicates for bootstrap testing is shown in the Figure.

Sika Deer Carrying *Babesia* Parasites Closely Related to *B. divergens*, Japan

Technical Appendix



Prefectures with no sika deer positive for *Babesia* spp. (N):
Chiba (5), Shizuoka (3), Hyogo (1), Hiroshima (5),
Tokushima (5), Oita (3), Kumamoto (4), Kagoshima (4)

Technical Appendix Figure. Map of Japan showing prefectures where *rDNA* genes of *Babesia divergens*-like parasites were detected by PCR in blood samples from sika deer (black areas). Sample identification numbers are listed under prefecture names; numbers in parentheses indicate the number of positive samples/total number tested samples. A list of prefectures where no positive samples were found is shown at the bottom of the figure; the number of animals tested is shown in parentheses.