

Global Avian Influenza Surveillance in Wild Birds: A Strategy to Capture Viral Diversity

Catherine C. Machalaba, Sarah E. Elwood, Simona Forcella, Kristine M. Smith, Keith Hamilton, Karim B. Jebara, David E. Swayne, Richard J. Webby, Elizabeth Mumford, Jonna A.K. Mazet, Nicolas Gaidet, Peter Daszak, William B. Karesh

Wild birds play a major role in the evolution, maintenance, and spread of avian influenza viruses. However, surveillance for these viruses in wild birds is sporadic, geographically biased, and often limited to the last outbreak virus. To identify opportunities to optimize wild bird surveillance for understanding viral diversity, we reviewed responses to a World Organisation for Animal Health–administered survey, government reports to this organization, articles on Web of Knowledge, and the Influenza Research Database. At least 119 countries conducted avian influenza virus surveillance in wild birds during 2008–2013, but coordination and standardization was lacking among surveillance efforts, and most focused on limited subsets of influenza viruses. Given high financial and public health burdens of recent avian influenza outbreaks, we call for sustained, cost-effective investments in locations with high avian influenza diversity in wild birds and efforts to promote standardized sampling, testing, and reporting methods, including full-genome sequencing and sharing of isolates with the scientific community.

Avian influenza is a global threat to food animal production and distribution systems, as well as human health. However, sustained, comprehensive, and coordinated global efforts to monitor the continually changing genetic diversity of avian influenza viruses circulating in nature are lacking (1,2). Two avian influenza viruses are current pandemic threats: highly pathogenic avian influenza A(H5N1) virus,

Author affiliations: EcoHealth Alliance, New York, New York, USA (C.C. Machalaba, S.E. Elwood, K.M. Smith, P. Daszak, W.B. Karesh); World Organisation for Animal Health, Paris, France (S. Forcella, K. Hamilton, K.B. Jebara, W.B. Karesh); Ministry of Environment and Water, Dubai, United Arab Emirates (K.B. Jebara); US Department of Agriculture, Athens, Georgia, USA (D.E. Swayne); St. Jude Children's Research Hospital, Memphis, Tennessee, USA (R.J. Webby); World Health Organization, Geneva, Switzerland (E. Mumford); University of California, Davis, California, USA (J.A.K. Mazet); French Agricultural Research Centre for International Development, Montpellier, France (N. Gaidet); International Union for the Conservation of Nature Species Survival Commission, New York (W.B. Karesh)

DOI: <http://dx.doi.org/10.3201/eid2104.141415>

which has spilled over repeatedly to humans since its first report in 1996; and novel avian influenza A(H7N9) virus, first detected in March 2013, which has caused serious human infections across China. Wild birds played a role in the evolution of influenza A(H7N9) virus (3) and might have contributed to the spread of this virus to parts of Asia, Europe, and Africa after a 2005 outbreak in birds at Qinghai Lake in China (4,5).

Most recently, wild birds have been considered as having a role in the unexpected appearance and spread of influenza A(H5N8) virus in Europe in November 2014 and in North America in December 2014; genetically similar lineages have also been found in South Korea and Japan (6). Concern about highly pathogenic avian influenza viruses has generated sporadic attention and investments paired to specific subtypes. For example, alarm about influenza A(H5N1) virus resulted in short-term spending of hundreds of millions of dollars for wild bird–related research globally. Although interest and funding has since waned, threats from avian influenza viruses remain, and H5N1 and H7N9 subtype viruses continue to cause human infections and deaths.

Wild birds are natural reservoirs for avian influenza virus (4,7), host a wide diversity of subtypes, and provide a dynamic population for viral evolution and transmission to domestic flocks and mammals. Most hemagglutinin (HA) and neuraminidase (NA) subtypes have been detected in wild birds, although some infrequently (8,9). Highly pathogenic avian influenza viruses of poultry are defined by the World Organisation for Animal Health (OIE) as having an intravenous pathogenicity index >1.2 in 6-week-old chickens and causing $\geq 75\%$ deaths in 4 to 8-week-old chickens infected intravenously, or H5 or H7 virus isolates with a characteristic molecular sequence at the HA cleavage site (for full definition, see the OIE Terrestrial Animal Health Code [10]). Although these viruses are rarely detected in wild birds, they have been found in diverse wild bird species from disparate locations and can be potentially transmitted along transcontinental flyways (5).

Avian influenza rarely causes widespread deaths in wild birds, and influenza caused by H5N1 subtype virus represents the first major clinical avian influenza virus–associated disease recognized in wild birds since the outbreak of influenza caused by H5N3 subtype virus in South Africa in 1961 in common terns (*Sterna hirundo*) (7,11). Thus, limiting surveillance for avian influenza virus to only deaths of wild birds provides little insight into the diversity of avian influenza virus genotypes circulating globally or risk for future outbreaks in poultry or humans.

Programs implemented during sporadic periods of concern (e.g., the Global Avian Influenza Network for Surveillance of Wild Birds after influenza outbreaks caused by H5N1 subtype viruses in 2005 [12]) have provided useful data on avian influenza virus diversity for a limited number of isolates but have missed the opportunity to document how diversity changes over time (1). The situation for wild bird surveillance parallels similar constraints for surveillance of influenzas circulating in swine, as highlighted in a recent review (13).

While focusing on wild birds, we recognize the benefits of globally coordinated surveillance systems for all influenza viruses with the aims of multitype detection, whole-genome sequencing, sharing of virus isolates, and analysis of epidemiologic data to highlight changes in circulating virus subtype prevalence. In contrast to our current system, which largely emphasizes reacting to new avian influenza viruses once they are detected in poultry, more upstream tracking of this information could potentially provide a critical early warning system or at least provide a sense of the likely evolution and movement of these viruses so that more proactive action can be taken.

Wild bird surveillance information could directly benefit human and animal health through understanding of how avian influenza virus genes flow into poultry, swine, equine, and human influenza viruses and could provide a basis for strategies that reduce their risk for introduction into agricultural species and humans. The documented human case of influenza in Taiwan in 2013 caused by an H6N1 subtype virus (14) points to the potential value of broader avian influenza virus surveillance in other species because this subtype has rarely been included in influenza surveillance systems.

In recognition of the potential benefits of surveillance, the OIE–Food and Agriculture Organization global network of expertise on animal influenza (OFFLU) established a working group on wildlife influenzas in October 2014; this group has highlighted the need for wild bird surveillance to understand circulation dynamics of avian influenza virus and recommends full-genome sequencing (15). In April 2014, the Strategic Alliances for the Coordination of Research on the Major Infectious Diseases of Animals and Zoonoses identified avian influenza surveillance in wild birds as a top priority for collaboration as part of a 10-year strategic research agenda (16). We review current surveillance efforts and provide recommendations toward establishing an effective surveillance program for avian influenza in wild birds.

Assessing the Current State of Wild Bird Surveillance

A global look at multiple major data sources on wild bird surveillance for avian influenza viruses by country has not been previously reported. An excellent review by Hoye

et al. (1) analyzed 191 literature reports of wild bird surveillance initiated during 1961–2007 and identified needs for a coordinated scientific approach, including refined sampling and screening strategies. Our analysis incorporates additional surveillance and viral diversity data sources available at a global level and assesses current/recent efforts to determine capacity and actions needed for an effective avian influenza virus tracking system in wild birds (online Technical Appendix, <http://wwwnc.cdc.gov/EID/article/21/4/14-1415-Techapp1.pdf>). Other sources of information (e.g., in local languages) might provide further useful information for future analyses.

OIE Member Survey

Forty-six (25.8%) of 178 OIE member countries responded to an influenza surveillance survey (online Technical Appendix). Eleven (23.9%) of 46 responding countries reported active (live birds) and passive (dead birds) surveillance activities (whether collected specifically for the study or by other means [e.g., sampling of hunter collections]); 14 (30.4%) reported active surveillance only; 14 (30.4%) conducted passive surveillance only; and the remaining 7 (15.2%) reported conducting no surveillance activity. Several of these countries cited a lack of funding as the impediment. Of 39 countries that reported surveillance (online Technical Appendix), 23 (58.9%) specified testing for virus subtypes deemed to be a higher pathogenic risk to poultry and humans (H5, H7, and all highly pathogenic avian influenza viruses identified) and 9 (23.0%) reported that they tested to determine all subtypes found. Only 11 subtype combinations were reported through the OIE survey from 9 countries during January 2012–March 2013.

OIE–World Animal Health Information Data Interface

A total of 116 countries submitted reports during the period examined (2008–2012) (17). Reports indicated that 82 of those countries conducted some form of avian influenza viral surveillance in wild birds (online Technical Appendix). One country reported highly pathogenic avian influenza virus (H5N1 subtype), 6 reported low pathogenicity avian influenza during 2008–2012 (including 10 HA segments and 39 subtype combinations), and the remainder reported no viruses. The other countries did not report detection. Of the 682 entries of highly pathogenic and low pathogenicity viruses detected in wild birds from the 37 countries reporting detection or suspicion of avian influenza, 244 entries were at the serotype (HA type or subtype combination) level of information.

Influenza Research Database

A total of 39 countries (online Technical Appendix) reported wild bird surveillance. This surveillance, which

was based on species name, was conducted during January 2008–June 2013.

Web of Knowledge

At least 54 countries reported wild bird surveillance during 2008–2013. This surveillance was reported mainly through peer-reviewed literature.

We found that ≥ 119 countries conducted and reported some form of avian influenza virus surveillance in wild birds during January 2008–January 2013 (online Technical Appendix). We identified 3 trends in surveillance efforts and their implications for comparability of data, as well as opportunities for refining surveillance strategies to capture diversity of avian influenza viruses.

The first trend was sampling method (active versus passive surveillance; number of birds and species sampled; types of samples collected as cloacal, oropharyngeal/tracheal, fecal, blood, or tissue samples; sampling site characterization; frequency and seasonality of sampling) differs widely across surveillance programs. This finding is consistent with analysis of surveillance efforts of Hoyer et al (1) conducted during 1961–2007 and suggested that unstandardized sampling approaches remain a chronic challenge for a global avian influenza virus surveillance system. Sampling methods, although not necessarily a limiting factor in identifying the wide range of these viruses circulating in nature, pose difficulties in discerning the usefulness of a report and comparing fluctuations in subtype virus prevalence between years or areas.

The second trend was that testing and virus characterization protocols vary widely. Some countries or programs screen for influenza A virus, some selectively screen for H5 and H7 virus subtypes, and some screen only for HA and NA virus subtypes, but not subtype virus combinations. Although we acknowledge resource and capacity limitations for analysis, the high level of effort required to capture or sample wild birds argues for investment to test for virus subtypes in addition to those believed to be currently highly pathogenic, to sequence as many as economically feasible, and to share samples and genomic data widely. Sequence data are needed to effectively track viral diversity, spread, and evolution.

The third trend was that critical data are deficient in most reporting systems. Measures of sampling effort are often absent. None of the data sources reviewed confirmed that a country did not conduct surveillance, which confounded analyses of negative findings. Approximate location or date of positive and negative findings is often missing, which hinders spatiotemporal risk analyses. Because positive findings are commonly reported without denominator data, especially for H5 and H7 subtype viruses, relative risk for spillover is difficult to assess. Only the Influenza Research Database (18) enables aggregation of full

genetic sequences. Finally, analysis of which types of samples yielded more positive results (e.g., cloacal swab versus fecal specimens) is not possible from these data, which limits their value in informing future sampling strategies.

We recognize that studies have had varied aims that contribute to differences in surveillance methods. However, the lack of standardized, sustained, and targeted surveillance prevents effective tracking of avian influenza virus diversity over time in the major reservoir for this group of viruses. Opportunities for systematic avian influenza virus surveillance could have been readily implemented through existing programs. For example, regional guidance through the European Union (EU) directives on wild bird surveys has promoted some level of comparability of data through a required uniform reporting format, which led to extensive data sharing. In 2006, EU member states reported having tested 120,706 birds (active and passive surveillance) in ≥ 330 species of 22 orders (19). Unfortunately, samples were not screened beyond influenza A virus (or, in some cases, H5 or H7 subtype viruses) under the EU directive. Future regional coordinated efforts could begin to increase avian influenza virus tracking even if only requiring full genome sequencing on a subset of samples. Ongoing systematic surveillance can inform avian influenza virus ecology research, going beyond pattern descriptions or experimental conditions by generating long-term data to address complex processes around dynamics of host immunity or viral diversity (2).

Capturing Viral Diversity

Improved understanding of avian influenza virus need not involve a monumental global effort but requires a shift in screening practices to move beyond emphasis on highly pathogenic avian influenza viruses. Where available, genomic sequencing for detection of avian influenza virus may provide more robust subtype findings.

A recent study detected by sequencing different HA subtype viruses that had not been detected by antigenic subtyping approaches (20). However, we call for a phase change in surveillance programs, which includes additional measures to track avian influenza virus diversity beyond subtype, at least in a subset of samples. Capturing information on all 8 avian influenza virus gene segments by routinely sequencing wild bird isolates would enhance understanding of avian influenza virus dynamics even more. For example, influenza A(H7N9) virus that caused respiratory disease in humans evolved from genetic contributions of ≥ 4 gene segment origins, including sources from wild birds (3). These contributions exemplify complex major genetic evolution around antigenic drift and gene reassortment, which subtype characterization alone does not capture.

The 6 less well-studied influenza virus gene segments may confer major determinants of infectivity, pathogenicity,

transmissibility, and host species susceptibility (e.g., the role of the basic polymerase 2 gene segment in host range and virulence) (9). Tracking gene segments may be especially useful for wild birds, given the high reassortment rate in their avian influenza virus genome sequences (21) and the unknown potential for transient genomes to become stable and pose risks for endemic infections or transmissibility or pathogenicity in humans. As a long-term goal, surveillance should be refined to maximize understanding of transmission factors (e.g., host receptivity and susceptibility, host dispersal) and ecologic factors (e.g., climate-dependent viral persistence, migration timing and range) that drive avian influenza virus prevalence in wild birds (22) and enable spillover, emergence, and maintenance (1).

Coordinating bodies, such as OFFLU, have an opportunity to develop specific standards for avian influenza virus diversity-oriented surveillance programs to ensure a clear strategy forward for the scientific community and its funders. Similarly, funders have a key role in driving efforts to track avian influenza virus diversity but will have to embrace sustained screening efforts for highly pathogenic and low pathogenicity avian influenza viruses and promote the value of increased sharing of negative findings and require full-genome sequencing of avian influenza viruses, even if required for only a subset of samples.

Reporting Systems

Despite benefits of international sharing of avian influenza viruses, regardless of their pathogenicity, there is no standardized and comprehensive reporting requirement beyond highly pathogenic avian influenza virus and H5 and H7 subtypes of low pathogenicity avian influenza viruses, nor adequate reporting incentives. Scientific journals and online databases enable sharing of information from research projects, as well as from official government surveillance, but these sources inherently underreport negative findings, and the lack of standardization among studies reduces value to authorities responsible for prevention and control. Online databases, such as the Influenza Research Database and GenBank, provide forums for detailed and consistent reporting, including sequence data.

We urge avian influenza virus surveillance funders to require reporting of results to the Influenza Research Database to drive sharing of metadata and genetic sequence information. This sharing would expand the utility of local- and national-level data to feed into global data analysis. The limited current comparability of data can inform database directors on the need to harmonize data deposition requirements, improve interoperability, and encourage full reporting of negative and positive results to optimize tracking potential for avian influenza viruses. Ongoing discussions between coordinating bodies, such as OFFLU, database designers, and funders, might help improve

efficiency and refinement of reporting systems and also ensure resource alignment across high-priority surveillance and reporting. Overall, a coordinated surveillance system should capitalize on global sharing tools to promote access to information that will support rapid detection of these virus and ongoing analysis.

Targeted Country Participation

A coordinated, annual surveillance system with a global perspective does not necessarily require participation from every country. Rather, resource allocation could be prioritized to provide sustained surveillance in a few targeted locations and in specific seasons that maximize information on viral diversity relevant to potential spread (e.g., high-risk species, species interfaces, major staging and migration stopover sites, and reassortment hotspots [23]). Recent analysis of avian influenza virus subtype diversity and richness suggests that 75% of HA and NA subtype diversity in wild birds could be captured through targeted surveillance efforts in the Northern Hemisphere over a 4-year period (24).

In addition, findings recently reported from the largest avian influenza surveillance program in wild birds ever implemented (a study by the US Department of Agriculture and Department of Interior in collaboration with authorities in Canada and Mexico) suggested that hotspots of avian influenza virus in wild birds were primarily located in the northern latitudes of the United States (25). These findings suggest that sites in the Northern Hemisphere are high-yield starting locations for viral diversity, especially because we found that countries considered as major sources of avian influenza virus diversity (e.g., Canada, China, Germany, Mongolia, Norway, Russia, the United Kingdom, and the United States) have reported surveillance data since 2008, but to different data portals (Figure; online Technical Appendix).

Coordinating extant programs in areas with major roles in avian influenza virus diversity would be a cost-effective first measure. Although we propose an initial focus on the Northern Hemisphere to leverage current investments and target surveillance on the basis of prior avian influenza study findings reported above (24,25), we do not intend to undermine the role of efforts elsewhere. We especially acknowledge that bias in surveillance effort has limited current knowledge on avian influenza virus diversity in the Southern Hemisphere (4). Surveillance in the Southern Hemisphere, where resources are available, provides highly useful information on exchange in the avian influenza virus genetic pool through bird migration (migration–shedding dynamics may enable avian influenza virus dispersal over extensive distances [5]), poultry trade, and maintenance of some specific phylogenetic lineages of these viruses.




Figure. Feasibility of coordinating and improving avian influenza surveillance in wild birds where viral diversity is highest. Countries in red, orange, and yellow currently self-report some type of avian influenza surveillance in wild birds (For a country list, see online Technical Appendix, <http://wwwnc.cdc.gov/EID/article/21/4/14-1415-Techapp1.pdf>). Country income levels, based on gross domestic product, further suggest financial capacity to contribute to a coordinated surveillance system. The polar view emphasizes where most avian influenza viral diversity is circulating.

Study Limitations

Our analysis was intended to provide a snapshot of recent avian influenza virus surveillance effort in wild birds to explore how existing infrastructure could be optimized to capture viral diversity. There are limitations of our study, including information missed through our data compilation methods. Only a subset of OIE member countries responded to the survey, and some responses were incomplete or unclear, which was potentially caused by instructions or language barriers. In addition, because only highly pathogenic influenza viruses or influenza A H5 or H7 subtype viruses are required for reporting to OIE, data for low pathogenicity avian influenza viruses were reported voluntarily by countries. Thus, reports to OIE cannot be assumed to be comprehensive.

We targeted information specific for wild birds, but it was not always possible from the data sources reviewed to determine whether animals were truly free-ranging or captive (e.g., data from the World Animal Health Information Data Interface [WAHID] was specific for wild species but did not distinguish by setting, which might show different dynamics for avian influenza viruses). Articles reporting surveillance from the Web of Knowledge were reviewed for specific parameters (e.g., time frame, wild bird species), but this method excluded papers not explicitly reporting

this information. Additional relevant papers were likely available through other databases.

Non-reporting does not indicate lack of surveillance and likely increases the number of countries actually conducting surveillance. We also recognize that reports of surveillance are not always verified for accuracy, and surveillance at one time point might not indicate current capacity. We included sampling effort reported that occurred during 2008–2013, but not all data sources spanned that time frame (e.g., the OIE survey only reflected 1 year of activities). Last, because in some cases it was not possible to ascertain the lead institution organizing surveillance efforts in a given country (government versus in-country research organizations or outside institutions), this information was not compiled. We acknowledge that outside research might not be indicative of true in-country capacity and might have different implications for reporting. Despite several limitations, our findings suggest that investments have been recently made for surveillance in most countries and thus provide a starting infrastructure for capturing avian influenza virus diversity.

A Cost-effective Surveillance System

Establishing collaborative networks among countries would be cost-effective, reduce the need for additional

laboratory capacity in regions of interest for surveillance, and complement and support other surveillance programs in maintaining trained and operational field teams in targeted locations, instead of rebuilding local capacity and logistics for each new avian influenza virus threat. A more robust understanding of virus diversity and changing viral trends might inform biosecurity efforts at the wildlife/domestic animal interface in which virus spillover might occur. This understanding has potential value, given devastation from the death or culling of millions of birds during 1 outbreak of highly pathogenic avian influenza (26), because surveillance provides information on genes of low pathogenicity avian influenza viruses that could recombine or mutate to produce high pathogenic avian influenza viruses (27).

Stronger capacity for early detection of changing avian influenza virus dynamics in reservoir populations may provide inputs to support public health and pandemic preparedness. These inputs include potentially useful genetic material for vaccine development, detection of virus origins, refined diagnostic capacity beyond influenza A screening, improved understanding of molecular determinants of transmission and pathogenesis from gene segment characterization, and analysis of the potential for virus spread through migration and trade (28).

The cost of avian influenza virus outbreaks in poultry is substantial; outbreaks of influenza A(H5N1) virus during 2004–2009 caused US \$30 billion in damage (29), and the frequency of highly pathogenic avian influenza outbreaks in poultry appears to be increasing (11). Rather than sporadically releasing large amounts of funding for wild bird surveillance when specific avian influenza viruses emerge, sustained national, regional, and global investments can provide the targeted baseline level of systematic surveillance we propose. Many countries, especially where avian influenza virus diversity in wild birds is highest, are already investing in some form of avian influenza virus surveillance in wild birds.

Current efforts should be refined by leveling the investment roller coaster that has funded subtype-specific wild bird surveillance toward a lower-cost but long-term investment in collecting and sequencing wild bird avian influenza viruses. Global coordinating bodies, such as OFFLU (a network linking influenza experts and laboratories working with poultry and swine influenzas and World Health Organization expertise) (13), provide collaborative forums for government agencies and researchers to compile and share sequences and isolates. For wild bird avian influenza viruses specifically and all influenza viruses, we would be remiss not to work toward coordinated surveillance to support more effective assessment, preparation, and response for emerging influenza viruses that pose potential public health threats.

Acknowledgments

We thank the country delegates who responded to the survey administered by the OIE, as well as the Influenza Research Database team for their guidance.

The concept of this study was developed by W.B.K., C.C.M., S.M.E., and K.M.S. in collaboration with K.H. C.C.M. reviewed the literature base, influenza research database, and WAHID reports. S.M.E. reviewed the OIE survey. C.C.M., S.M.E., and W.B.K. drafted the paper. W.B.K. provided technical guidance throughout the data review process. S.F. and K.B.J. designed and disseminated the survey to OIE delegates and designed and collected data from WAHID. D.E.S., R.J.W., and E.M. provide oversight and editing on surveillance and virology aspects. J.A.K.M., N.G., W.B.K., and P.D. contributed to ecology and field techniques interpretation and recommendations. All authors reviewed the report and provided input.

C.C.M., S.M.E., K.M.S., J.A.K.M., P.D., and W.B.K. were supported by the US Agency for International Development Emerging Pandemic Threats Program PREDICT.

Ms. Machalaba is the Program Coordinator for Health and Policy at the EcoHealth Alliance, New York, New York. She serves as Program Officer for the IUCN Species Survival Commission Wildlife Health Specialist Group and is a doctoral student in Public Health at the City University of New York. Her research interests focus on One Health approaches to human, animal, and environmental health challenges.

References

- Hoye BJ, Munster VJ, Nishiura H, Klaassen M, Fouchier RA. Surveillance of wild birds for avian influenza virus. *Emerg Infect Dis*. 2010;16:1827–34 and <http://dx.doi.org/10.3201/eid1612.100589>
- Latorre-Margalef N, Tolf C, Grosbois V, Avril A, Bengtsson D, Wille M, et al. Long-term variation in influenza A virus prevalence and subtype diversity in migratory mallards in northern Europe. *Proc Biol Sci*. 2014;281:20140098 and <http://dx.doi.org/10.1098/rspb.2014.0098>
- Liu D, Shi W, Shi Y, Wang D, Xiao H, Li W, et al. Origin and diversity of novel avian influenza A H7N9 viruses causing human infection: phylogenetic, structural, and coalescent analyses. *Lancet*. 2013;381:1926–32 and [http://dx.doi.org/10.1016/S0140-6736\(13\)60938-1](http://dx.doi.org/10.1016/S0140-6736(13)60938-1)
- Olsen B, Munster VJ, Wallensten A, Waldenstrom J, Osterhaus AD, Fouchier RA. Global patterns of influenza a virus in wild birds. *Science*. 2006;312:384–8 and <http://dx.doi.org/10.1126/science.1122438>
- Gaidet N, Cappelle J, Takekawa JY, Prosser DJ, Iverson SA, Douglas DC, et al. Potential spread of highly pathogenic avian influenza H5N1 by wildfowl: dispersal ranges and rates determined from large-scale satellite telemetry. *J Appl Ecol*. 2010;47:1147–57. <http://dx.doi.org/10.1111/j.1365-2664.2010.01845.x>
- Food and Agriculture Organization of the United Nations. Avian influenza A(H5N8) detected in Europe... a journey to the West? November 14, 2014 [cited 2014 Nov 17]. http://www.fao.org/ag/againfo/home/en/news_archive/2014_A-H5N8_detected_in_Europe.html
- Stallknecht DE, Brown JD. Wild birds and the epidemiology of avian influenza. *J Wildl Dis*. 2007;43(suppl):S15–20.

8. Fouchier RA, Munster VJ. Epidemiology of low pathogenic avian influenza viruses in wild birds. *Rev Sci Tech*. 2009;28:49–58 .
9. Chen Y, Liang W, Yang S, Wu N, Gao H, Sheng J, et al. Human infections with the emerging avian influenza A H7N9 virus from wet market poultry: clinical analysis and characterisation of viral genome. *Lancet*. 2013;381:1916–25 and [http://dx.doi.org/10.1016/S0140-6736\(13\)60903-4](http://dx.doi.org/10.1016/S0140-6736(13)60903-4)
10. World Organisation for Animal Health. Chapter 10.4. Article 10.4.1. Infection with avian influenza viruses. In: *Terrestrial animal health code*. 23rd ed; Paris: The Organisation; 2014 [cited 2014 Dec 22]. http://www.oie.int/index.php?id=169&L=0&htmfile=chapitre_avian_influenza_viruses.htm
11. Alexander DJ. An overview of the epidemiology of avian influenza. *Vaccine*. 2007;25:5637–44 and <http://dx.doi.org/10.1016/j.vaccine.2006.10.051>
12. Wildlife Conservation Society. Avian influenza (H5N1), 2014 [cited 2014 Nov 17]. <http://www.wcs.org/conservation-challenges/wildlife-health/disease-investigations/avian-influenza-h5n1.aspx>
13. Vincent A, Awada L, Brown I, Chen H, Claes F, Dauphin G, et al. Review of influenza A virus in swine worldwide: a call for increased surveillance and research. *Zoonoses Public Health*. 2014;61:4–17 and <http://dx.doi.org/10.1111/zph.12049>
14. Shi W, Shi Y, Wu Y, Liu D, Gao GF. Origin and molecular characterization of the human-infecting H6N1 influenza virus in Taiwan. *Protein Cell*. 2013;4:846–53.
15. World Organisation for Animal Health–Food and Agriculture Organization of the United Nations OFFLU. OFFLU strategy document for surveillance and monitoring of influenzas in animals, May 2013 [cited 2014 Dec 18]. <http://www.offlu.net/fileadmin/home/en/publications/pdf/OFFLUSurveillance.pdf>
16. OFFLU and STAR-IDAZ. A consultation to develop a global animal influenza research agenda. Paris, April 8–9, 2014 [cited 2014 Dec 18]. http://www.offlu.net/fileadmin/home/en/publications/pdf/300714/OFFLU_STAR-IDAZ_Executive_Summary_final.pdf
17. World Organization for Animal Health. World Animal Health Information Database (WAHID) interface. December 30, 2013 [cited 2014 Dec 18]. http://www.oie.int/wahis_2/public/wahid.php/Wahidhome/Home
18. Squires RB, Noronha J, Hunt V, Garcia-Sastre A, Macken C, Baumgarth N, et al. Influenza research database: an integrated bioinformatics resource for influenza research and surveillance. *Influenza Other Respir Viruses*. 2012;6:404–16.
19. Hesterberg U, Harris K, Stroud D, Guberti V, Busani L, Pittman M, et al. Avian influenza surveillance in wild birds in the European Union in 2006. *Influenza Other Respir Viruses*. 2009;3:1–14.
20. Fries AC, Nolting JM, Bowman AS, Killian ML, Wentworth DE, Slemmons RD. Genomic analyses detect Eurasian-lineage H10 and additional H14 influenza A viruses recovered from waterfowl in the Central United States. *Influenza Other Respir Viruses*. 2014; 8:493–8.
21. Dugan VG, Chen R, Spiro DJ, Sengamalay N, Zaborsky J, Ghedin E, et al. The evolutionary genetics and emergence of avian influenza viruses in wild birds. *PLoS Pathog*. 2008;4:e1000076 and <http://dx.doi.org/10.1371/journal.ppat.1000076>
22. Gaidet N, Caron A, Cappelle J, Cumming GS, Balanca G, Hammoumi S, et al. Understanding the ecological drivers of avian influenza virus infection in wildfowl: a continental-scale study across Africa. *Proc Biol Sci*. 2012;279:1131–41 and <http://dx.doi.org/10.1098/rspb.2011.1417>
23. Fuller TL, Gilbert M, Martin V, Cappelle J, Hosseini P, Njabo K, et al. Predicting hotspots for influenza virus reassortment. *Emerg Infect Dis*. 2013;19:581–8 and <http://dx.doi.org/10.3201/eid1904.120903>
24. Olson SH, Parmley J, Soos C, Gilbert M, Latorre-Margalef N, Hall JS, et al. Sampling strategies and biodiversity of influenza a subtypes in wild birds. *PLoS ONE*. 2014;9:e90826 and <http://dx.doi.org/10.1371/journal.pone.0090826>
25. Bevins SN, Pedersen K, Lutman MW, Baroch JA, Schmit BS, Kohler D, et al. Large-scale avian influenza surveillance in wild birds throughout the United States. *PLoS ONE*. 2014;9:e104360 and <http://dx.doi.org/10.1371/journal.pone.0104360>
26. Chmielewski R, Swayne DE. Avian influenza: public health and food safety concerns. *Annu Rev Food Sci Technol*. 2011;2:37–57.
27. Horimoto T, Rivera E, Pearson J, Senne D, Krauss S, Kawaoka Y, et al. Origin and molecular changes associated with emergence of a highly pathogenic H5N2 influenza virus in Mexico. *Virology*. 1995;213:223–30 and <http://dx.doi.org/10.1006/viro.1995.1562>
28. Bahl J, Krauss S, Kuhnert D, Fourment M, Raven G, Pryor SP, et al. Influenza a virus migration and persistence in North American wild birds. *PLoS Pathog*. 2013;9:e1003570 and <http://dx.doi.org/10.1371/journal.ppat.1003570>
29. Newcomb J, Harrington T, Aldrich S. The economic impact of selected infectious disease outbreaks. Cambridge (MA): Bio-Economic Research Associates; 2011.

Address for correspondence: William B. Karesh, EcoHealth Alliance, 460 West 34th St, 17th Floor, New York, NY 10001, USA; email: karesh@ecohealthalliance.org

Global Avian Influenza Surveillance in Wild Birds: A Strategy to Capture Viral Diversity

Technical Appendix

Global Avian Influenza Surveillance in Wild Birds

Methods

We analyzed data from 1) a survey of the Organisation for Animal Health (OIE) member countries conducted by OIE, 2) official OIE animal health reports from member countries by using the World Animal Health Information Database (WAHID) interface (<http://oie.int/wahid>) and the WAHID-Wild interface (<http://oie.int/wahidwild>); 3) reports of avian influenza wild bird surveillance results published in the peer-reviewed literature by using the Web of Knowledge; and 4) the Influenza Research Database (<http://www.fludb.org>).

OIE Member Survey

Influenza caused by H7N5 subtype virus that is affecting humans in China is not causing clinical disease in birds. Thus, more proactive approaches to surveillance in birds are necessary to detect viral infection. To determine the scope of current surveillance activities in wild birds, in April 2013, the OIE asked all 178 member countries to provide information on surveillance activities for avian influenza (caused by low pathogenic and highly pathogenic influenza viruses) during the previous 12 months. Results received from 46 of 178 member countries were reviewed.

OIE WAHID Interface Data

We also reviewed avian influenza reporting from the disease timeline database on the WAHID interface (www.oie.int/wahid) during 2008–2012 by using data provided by OIE member countries (data for 2013 were not available at the time of analysis). Reporting of

avian influenza virus subtypes H5, H7, and highly pathogenic avian influenza virus identified in wild birds to OIE is mandated, and reporting of other subtypes of low pathogenic avian influenza virus is voluntary as part of the OIE Worldwide Monitoring System for Wild Animals. For the purposes of identifying countries reporting surveillance activities for this analysis, we included countries reporting any of the following occurrence categories: confirmed infection but not clinical disease, suspected, confirmed clinical disease, infection/infestation in ≥ 1 zones, or clinical disease limited to ≥ 1 zones, disease suspected but not confirmed limited to ≥ 1 zones, or absent.

Influenza Research Database

To determine other sources of information on avian influenza surveillance in wild birds from outside sources (or potentially in conjunction with) reporting to OIE, we collected wild avian surveillance data from the Influenza Research Database (<http://www.fludb.org/>) by using a 2-step approach. First, we conducted a search on November 26, 2013, on the Avian Surveillance Summary Search (an option under Search Data→Animal Surveillance) refined for the host type parameter Wild. The search excluded avian surveillance records categorized as captive wild or domestic host type. Total wild bird counts per country were summed. Entries for countries with $n = 0$ wild bird counts were excluded. By our definition ($n > 0$ wild bird counts per country), reports of avian influenza viruses originating from 40 countries were found. Second, by using the Animal Surveillance Search page, we set display type to customizable summary table. Country was selected as the primary y-axis attribute, and collection year was selected as the primary x-axis attribute. The remaining default search settings were used. The resulting table displayed years with sampling data for each country. For the 40 countries from which there were reports of $n > 0$ wild bird counts, data records were reviewed to determine if wild birds were sampled during January 2008–June 2013 (collection date). Individual collection records were reviewed to ensure that the records reflected $n > 0$ wild birds as opposed to captive wild or domestic birds (noted as behavior classification in the collection record) during the time frame. Most of total effort (sample collection) for countries was reported during 2008–2013.

Web of Knowledge

To explore the extent of reporting on avian influenza surveillance in wild birds in the scientific literature, we searched the Thompson Reuters Web of Knowledge system (www.webofknowledge.com) version 5.12 (all databases) on November 27, 2013, topic search term [Avian Influenza] with publication date parameters for 2008–2013. The search yielded 20,757 results. The search was refined for search term wild bird and yielded 1,220 results. The search was further refined for surveillance, which yielded 431 results. We limited inclusion of reports to those conducting surveillance during 2008–2013. Abstracts were reviewed for surveillance time frames and to confirm $n > 0$ wild bird sampling in the country(s). When results were not clear, full texts were reviewed. Papers needing more review without full-text online versions were searched by using Google Scholar (www.scholar.google.com); if not available on either system, they were not included. Publications focused on surveillance of domestic animals involving experimental infection or investigating surveillance or testing method validation were excluded. In general, information was not included from review papers, except where it was explicit that wild birds had been sampled in a given country within the specified time frame. The list of countries from which reports of conducting wild bird surveillance during January 2008–January 2013 by an article available on Web of Knowledge was compiled (online Appendix Table). Information matching the headings of the OIE survey was also extracted for a subset of 26 countries not responding to the OIE survey but still reporting in the literature available on Web of Knowledge to gain a sense of their sampling, testing, and reporting approaches.

Table. Current and recent surveillance for avian influenza virus in wild birds, by country and income level, reported surveillance by country*

Income level, country	OIE survey	Reporting to OIE	Web of Knowledge	Influenza Research Database
High				
Australia	X	NR	X (1,2)	X
Austria	NR	X	X (3)	NR
Bahrain	NR	X	NR	NR
Belgium	X	X	NR	NR
Canada	X	X	X (4,5)	X
Chile	NR	X	NR	NR
Taiwan	X	X	NR	X
Croatia	NR	X	NR	NR
Czech Republic	X	X	X (6)	NR
Denmark	X	X	X (7)	NR
Equatorial Guinea	NR	X	NR	NR
Estonia	NR	X	NR	NR
Finland	X	X	NR	NR
France	NR	X	X (8)	NR
Germany	NR	X	X (3,9)	NR
Guadeloupe	NR	NR	X (10)	NR

Income level, country	OIE survey	Reporting to OIE	Web of Knowledge	Influenza Research Database
Hong Kong	NR	NR	X (11,12)	NR
Ireland	NR	X	NR	NR
Israel	NR	X	NR	NR
Italy	X	X	X (13,14)	NR
Japan	X	X	X (15-21)	X
South Korea	X	X	X (22-28)	NR
Kuwait	NR	X	NR	NR
Latvia	NR	NR	NR	X
Lithuania	NR	NR	X (29)	X
the Netherlands	NR	X	X (30)	X
New Caledonia	NR	X	NR	NR
New Zealand	X	X	NR	NR
Norway	X	X	X (31)	NR
Oman	NR	NR	NR	X
Poland	X	X	X (32,33)	NR
Portugal	X	X	X (34,35)	NR
Russia	NR	NR	X (36-38)	X
San Marino	NR	X	NR	NR
Saudi Arabia	NR	X	NR	NR
Singapore	X	X	NR	NR
Slovakia	NR	X	X (39)	NR
Slovenia	X	X	X (40)	NR
Spain	X	X	X (41)	NR
Sweden	NR	X	NR	NR
Switzerland	NR	NR	X (3,42)	NR
United Kingdom	X	X	X (43,44)	NR
United States	X	X	X (5,44-51)	X
Uruguay	NR	X	NR	NR
Upper middle				
Algeria	NR	X	NR	NR
Argentina	X	X	X (52)	NR
Azerbaijan	NR	X	NR	NR
Bosnia and Herzegovina	X	X	NR	NR
Botswana	NR	X	X (53,54)	NR
Brazil	NR	X	NR	NR
Bulgaria	NR	X	X (55)	X
China	NR	X	X (56,57)	X
Colombia	X	NR	NR	X
Greece	X	NR	NR	X
Greenland	NR	NR	X (7)	X
Hungary	X	X	NR	X
Iceland	NR	X	NR	X
Iraq	X	NR	NR	NR
Kazakhstan	X	NR	X (38)	NR
Lebanon	NR	NR	NR	X
Libya	NR	X	NR	NR
Malaysia	X	X	NR	NR
Mauritius	X	NR	NR	NR
Mexico	NR	X	X (5,58)	NR
Montenegro	NR	X	NR	NR
Namibia	NR	X	NR	NR
Peru	X	X	X (59)	NR
Romania	NR	X	NR	X
Serbia	X	X	NR	NR
South Africa	NR	X	X (53,54)	NR
Thailand	X	X	X (60)	X
Tunisia	X	X	NR	NR
Turkey	X	NR	X (61)	X
Lower middle				
Armenia	NR	X	NR	NR
Bhutan	NR	NR	NR	X
Cameroon	NR	NR	NR	X
Côte d'Ivoire	X	X	NR	NR
Egypt	NR	NR	X (62,63)	X
El Salvador	NR	X	NR	NR
Georgia	NR	NR	X (64)	X
Ghana	NR	NR	X (65)	X
Guatemala	NR	NR	X (66)	X

Income level, country	OIE survey	Reporting to OIE	Web of Knowledge	Influenza Research Database
India	NR	X	X (67)	NR
Lesotho	NR	X	NR	NR
Mauritania	NR	NR	X (54)	NR
Mongolia	X	X	X (21,38,68,69)	X
Morocco	NR	X	NR	NR
Nigeria	NR	X	X (70)	NR
Pakistan	NR	NR	X (71)	NR
Papua New Guinea	NR	NR	NR	X
Paraguay	NR	X	NR	NR
Moldova	NR	X	NR	NR
Republic of the Congo	NR	NR	NR	X
Senegal	NR	X	X (54)	NR
Sri Lanka	NR	NR	NR	X
Sudan	NR	X	NR	X
Swaziland	NR	X	NR	NR
Ukraine	NR	NR	X (72)	NR
Vanuatu	NR	X	NR	NR
Vietnam	X	NR	X (73,74)	X
Yemen	X	NR	NR	NR
Zambia	NR	NR	X (75)	NR
Low				
Bangladesh	X	X	X (76)	X
Benin	NR	X	NR	X
Cambodia	NR	NR	X (77)	X
Central African Republic	NR	NR	NR	X
Eritrea	NR	X	NR	NR
Ethiopia	NR	X	NR	NR
Guinea	NR	X	NR	NR
Haiti	NR	X	NR	NR
Kenya	NR	NR	X (76)	NR
North Korea	X	NR	NR	NR
Madagascar	NR	X	NR	NR
Mali	NR	NR	X (54)	NR
Mozambique	NR	NR	X (53,54)	NR
Nepal	NR	X	NR	NR
Tanzania	NR	X	NR	NR
Uganda	NR	X	NR	X
Zimbabwe	NR	X	X (53,54)	X

*Reported surveillance by country as per 1) responses to OIE survey confirming surveillance activity; 2) official reports to OIE indicating surveillance activities, 2008–2012; 3) Web of Knowledge literature indicating surveillance activities during 2008–2013; and 4) information reported to the Influenza Research Database. Country income levels are based on the 2013 United Nations categorization. OIE, World Organisation for Animal Health; X, ≥1 report was found in the relevant reporting category; NR, no report. Values in parentheses are references.

References

1. Hansbro PM, Warner S, Tracey JP, Arzey KE, Selleck P, O'Riley K, et al. Surveillance and analysis of avian influenza viruses, Australia. *Emerg Infect Dis.* 2010;16:1896–904. [PubMed](https://pubmed.ncbi.nlm.nih.gov/20071111/)
<http://dx.doi.org/10.3201/eid1612.100776>
2. Hoque MA, Burgess GW, Greenhil AR, Hedlefs R, Skerratt LF. Causes of morbidity and mortality of wild aquatic birds at Billabong Sanctuary, Townsville, North Queensland, Australia. *Avian Dis.* 2012;56:249–56. [PubMed](https://pubmed.ncbi.nlm.nih.gov/22411111/) <http://dx.doi.org/10.1637/9863-072611-Case.1>

3. Knight-Jones TJD, Hauser R, Matthes D, Staerk KDC. Evaluation of Effectiveness and Efficiency of Wild Bird Surveillance for Avian Influenza. *Vet Res.* 2010;41:50. [PubMed](#)
<http://dx.doi.org/10.1051/vetres/2010023>
4. Parmley J, Lair S, Leighton FA. Canada's inter-agency wild bird influenza survey. *Integrative Zoology.* 2009;4:409–17. [PubMed](#) <http://dx.doi.org/10.1111/j.1749-4877.2009.00177.x>
5. Pasick J, Pedersen J, Hernandez MS. Avian influenza in North America, 2009–2011. *Avian Dis.* 2012;56:845–8. [PubMed](#) <http://dx.doi.org/10.1637/10206-041512-Reg.1>
6. Nagy A, Cernikova L, Krivda V, Hornickova J. Digital genotyping of avian influenza viruses of H7 subtype detected in central Europe in 2007–2011. *Virus Res.* 2012;165:126–33. [PubMed](#)
<http://dx.doi.org/10.1016/j.virusres.2012.02.005>
7. Hulsager CK, Breum SO, Trebbien R, Handberg KJ, Therkildsen OR, Madsen JJ, et al. Surveillance for avian influenza viruses in wild birds in Denmark and Greenland, 2007–10. *Avian Dis.* 2012;56:992–8. [PubMed](#) <http://dx.doi.org/10.1637/10190-041012-ResNote.1>
8. Lebarbenchon C, Albespy F, Brochet A-L, Grandhomme V, Renaud F, Fritz H, et al. Spread of avian influenza viruses by common teal (*Anas crecca*) in Europe. *PLoS ONE.* 2009;4:e7289. [PubMed](#)
<http://dx.doi.org/10.1371/journal.pone.0007289>
9. Rabl S, Rinder M, Neubauer-Juric A, Bogner K-H, Korbel R, Buettner M. Surveillance of wild birds for avian influenza a virus (AIV) in Bavaria in the years 2007 and 2008 [in German]. *Berl Munch Tierarztl Wochenschr.* 2009;122:486–93. [PubMed](#)
10. Lefrançois T, Hendrikx P, Ehrhardt N, Millien M, Gomez L, Gouyet L, et al. Surveillance of avian influenza in the Caribbean through the Caribbean Animal Health Network: surveillance tools and epidemiologic studies. *Avian Dis.* 2010;54:369–73. [PubMed](#) <http://dx.doi.org/10.1637/8787-040109-ResNote.1>
11. Leung YH, Cheung P, Zhang L-J, Wu Y-O, Chow K-C, Ho C-K, et al. Influenza viruses in wild birds in Hong Kong, 2003–2010. *Influenza Other Respi Viruses.* 2011;5:77–8.
12. Smith GJ, Vijaykrishna D, Ellis TM, Dyrting KC, Leung YH, Bahl J, et al. Characterization of avian influenza viruses A (H5N1) from wild birds, Hong Kong, 2004–2008. *Emerg Infect Dis.* 2009;15:402–7. [PubMed](#) <http://dx.doi.org/10.3201/eid1503.081190>
13. Kelvin AA, Meloni D, Sansonetti P, Borghetto I, Rowe T, Santangelo R, et al. Influenza monitoring in Sardinia, Italy identifies H3 subtype in Mediterranean wild migratory birds. *J Infect Dev Ctries.* 2012;6:786–97. [PubMed](#) <http://dx.doi.org/10.3855/jidc.2793>

14. Delogu M, Piredda I, Pintore A, Cabras P, Cotti C, Ghetti G, et al. Avian influenza and animal health risk: conservation of endemic threatened wild birds in Sardinia Island. *Avian Dis.* 2012;56:1034–9. [PubMed](#) <http://dx.doi.org/10.1637/10174-041012-ResNote.1>
15. Abao LN, Jamsransuren D, Bui VN, Ngo LH, Trinh DQ, Yamaguchi E, et al. Surveillance and characterization of avian influenza viruses from migratory water birds in eastern Hokkaido, the northern part of Japan, 2009–2010. *Virus Genes.* 2013;46:323–9. [PubMed](#) <http://dx.doi.org/10.1007/s11262-012-0868-9>
16. Bui VN, Ogawa H, Karibe K, Matsuo K, Nguyen TH, Awad SS, et al. Surveillance of avian influenza virus in migratory water birds in eastern Hokkaido, Japan. *J Vet Med Sci.* 2011;73:209–15. [PubMed](#) <http://dx.doi.org/10.1292/jvms.10-0356>
17. Fujimoto Y, Ito H, Shivakoti S, Nakamori J, Tsunekuni R, Otsuki K, et al. Avian influenza virus and paramyxovirus isolation from migratory waterfowl and shorebirds in San-in district of western Japan from 2001 to 2008. *J Vet Med Sci.* 2010;72:963–7. [PubMed](#) <http://dx.doi.org/10.1292/jvms.10-0012>
18. Ito T. Highly pathogenic avian influenza and wild birds. *Uirusu.* 2009;59:53–8. [PubMed](#) <http://dx.doi.org/10.2222/jsv.59.53>
19. Neagari Y. Challenges for the investigation of causes of wildlife mortality. *Japanese Journal of Zoo and Wildlife Medicine.* 2013;18:53–9.
20. Samad RA, Sakoda Y, Tsuda Y, Simulundu E, Manzoor R, Okamatsu M, et al. Virological surveillance and phylogenetic analysis of the PB2 genes of influenza viruses isolated from wild water birds flying from their nesting lakes in Siberia to Hokkaido, Japan in autumn. *Jpn J Vet Res.* 2011;59:15–22. [PubMed](#)
21. Yamamoto N, Sakoda Y, Motoshima M, Yoshino F, Soda K, Okamatsu M, et al. Characterization of a non-pathogenic H5N1 influenza virus isolated from a migratory duck flying from Siberia in Hokkaido, Japan, in October 2009. *Virol J.* 2011;8:65. [PubMed](#) <http://dx.doi.org/10.1186/1743-422X-8-65>
22. Baek YH, Pascua PN, Song M-S, Park KJ, Kwon HI, Lee JH, et al. Surveillance and characterization of low pathogenic H5 avian influenza viruses isolated from wild migratory birds in Korea. *Virus Res.* 2010;150:119–28. [PubMed](#) <http://dx.doi.org/10.1016/j.virusres.2010.03.002>
23. Choi JG, Kang HM, Jeon WJ, Choi KS, Kim KI, Song BM, et al. Characterization of clade 2.3.2.1 H5N1 highly pathogenic avian influenza viruses isolated from wild birds (mandarin duck and

- Eurasian eagle owl) in 2010 in Korea. *Viruses*. 2013;5:1153–74. [PubMed](#)
<http://dx.doi.org/10.3390/v5041153>
24. Kang H-M, Choi J-G, Kim K-I, Park H-Y, Park C-K, Lee Y-J. Genetic and antigenic characteristics of H4 subtype avian influenza viruses in Korea and their pathogenicity in quails, domestic ducks and mice. *J Gen Virol*. 2013;94:30–9. [PubMed](#) <http://dx.doi.org/10.1099/vir.0.046581-0>
 25. Kang HM, Jeong OM, Kim MC, Kwon JS, Paek MR, Choi JG, et al. Surveillance of avian influenza virus in wild bird fecal samples from South Korea, 2003–2008. *J Wildl Dis*. 2010;46:878–88. [PubMed](#) <http://dx.doi.org/10.7589/0090-3558-46.3.878>
 26. Kim H-R, Park C-K, Lee Y-J, Oem J-K, Kang H-M, Choi J-G, et al. Low pathogenic H7 subtype avian influenza viruses isolated from domestic ducks in South Korea and the close association with isolates of wild birds. *J Gen Virol*. 2012;93:1278–87. [PubMed](#)
<http://dx.doi.org/10.1099/vir.0.041269-0>
 27. Kwon HI, Song M-S, Pascua PN, Baek YH, Lee JH, Hong S-P, et al. Genetic characterization and pathogenicity assessment of highly pathogenic H5N1 avian influenza viruses isolated from migratory wild birds in 2011, South Korea. *Virus Res*. 2011;160:305–15. [PubMed](#)
<http://dx.doi.org/10.1016/j.virusres.2011.07.003>
 28. Lee D-H, Park J-K, Youn H-N, Lee Y-N, Lim T-H, Kim M-S, et al. Surveillance and isolation of HPAI H5N1 from wild mandarin ducks (*Aix galericulata*). *J Wildl Dis*. 2011;47:994–8. [PubMed](#)
<http://dx.doi.org/10.7589/0090-3558-47.4.994>
 29. Svazas S, Morkunas M, Verhagen J, Fouchier R, Morkunas J, Mundkur T, et al. Surveillance of wild waterbirds for avian influenza viruses in Lithuania. *Veterinarija Ir Zootechika*. 2012;60:72–8.
 30. Verhagen JH, Munster VJ, Majoor F, Lexmond P, Oanh V, Stumpel JBG, et al. Avian influenza A virus in wild birds in highly urbanized areas. *PLoS ONE*. 2012;7:e38256. [PubMed](#)
<http://dx.doi.org/10.1371/journal.pone.0038256>
 31. Tønnessen R, Kristoffersen AB, Jonassen CM, Hjortaa MJ, Hansen EF, Rimstad E, et al. Molecular and epidemiological characterization of avian influenza viruses from gulls and dabbling ducks in Norway. *Virol J*. 2013;10:112. [PubMed](#) <http://dx.doi.org/10.1186/1743-422X-10-112>
 32. Smietanka K, Minta Z, Włodarczyk R, Wyrostek K, Jozwiak M, Olszewska M, et al. Avian influenza viruses in wild birds at the Jeziorsko reservoir in Poland in 2008–2010. *Pol J Vet Sci*. 2012;15:323–8. [PubMed](#) <http://dx.doi.org/10.2478/v10181-011-0150-4>

33. Smietanka K, Pikula A, Minta Z, Meissner W. Evidence of persistence and multiple genetic modifications of H7N7 low-pathogenic avian influenza virus in wild mallards in Poland provided by phylogenetic studies. *Avian Pathol.* 2011;40:131–8. [PubMed](#)
<http://dx.doi.org/10.1080/03079457.2010.537304>
34. Tolf C, Bengtsson D, Rodrigues D, Latorre-Margalef N, Wille M, Figueiredo ME, et al. Birds and viruses at a crossroad: surveillance of influenza A virus in Portuguese waterfowl. *PLoS ONE.* 2012;7:e49002. [PubMed](#) <http://dx.doi.org/10.1371/journal.pone.0049002>
35. Henriques AM, Fagulha T, Barros SC, Ramos F, Duarte M, Luis T, et al. Multiyear surveillance of influenza A virus in wild birds in Portugal. *Avian Pathol.* 2011;40:597–602. [PubMed](#)
<http://dx.doi.org/10.1080/03079457.2011.618943>
36. Sivay MV, Baranovich T, Marchenko VY, Sharshov KA, Govorkova EA, Shestopalov AM, et al. Influenza A (H15N4) virus isolation in western Siberia, Russia. *J Virol.* 2013;87:3578–82.
[PubMed](#) <http://dx.doi.org/10.1128/JVI.02521-12>
37. Sivay MV, Sayfutdinova SG, Sharshov KA, Alekseev AY, Yurlov AK, Runstadler J, et al. Surveillance of influenza A virus in wild birds in the Asian portion of Russia in 2008. *Avian Dis.* 2012;56:456–63. [PubMed](#) <http://dx.doi.org/10.1637/9868-080111-Reg.1>
38. Marchenko VY, Alekseev AY, Sharshov KA, Petrov VN, Silko NY, Susloparov IM, et al. Ecology of influenza virus in wild bird populations in central Asia. *Avian Dis.* 2012;56:234–7. [PubMed](#)
<http://dx.doi.org/10.1637/9834-061611-ResNote.1>
39. Borovska P, Kabat P, Ficova M, Trnka A, Svetlikova D, Betakova T. Prevalence of avian influenza viruses, *Mycobacterium avium*, and *Mycobacterium avium* subsp *paratuberculosis* in marsh-dwelling passerines in Slovakia, 2008. *Biologia.* 2011;66:282–7.
<http://dx.doi.org/10.2478/s11756-011-0016-3>
40. Slavec B, Krapez U, Racnik J, Hari A, Wernig JM, Dovc A, et al. Surveillance of influenza A viruses in wild birds in Slovenia from 2006 to 2010. *Avian Dis.* 2012;56:999–1005. [PubMed](#)
<http://dx.doi.org/10.1637/10175-041012-ResNote.1>
41. Pérez-Ramirez E, Acevedo P, Allepuz A, Gerrikagoitia X, Alba A, Busquets N, et al. Ecological factors driving avian influenza virus dynamics in Spanish wetland ecosystems. *PLoS ONE.* 2012;7:e46418. [PubMed](#) <http://dx.doi.org/10.1371/journal.pone.0046418>

42. Baumer A, Feldmann J, Renzullo S, Mueller M, Thuer B, Hofmann MA. Epidemiology of avian influenza virus in wild birds in Switzerland between 2006 and 2009. *Avian Dis.* 2010;54:875–84. [PubMed http://dx.doi.org/10.1637/9119-110209-Reg.1](http://dx.doi.org/10.1637/9119-110209-Reg.1)
43. Breed AC, Irvine RM, Duncan D, Rae D, Snow L, Cook AJC, et al. An evaluation of wild bird avian influenza surveillance in Great Britain. *Avian Dis.* 2012;56:986–91. [PubMed http://dx.doi.org/10.1637/10166-040912-Reg.1](http://dx.doi.org/10.1637/10166-040912-Reg.1)
44. Siembieda JL, Johnson CK, Cardona C, Anchell N, Dao N, Reisen W, et al. Influenza A viruses in wild birds of the Pacific Flyway, 2005–2008. *Vector Borne Zoonotic Dis.* 2010;10:793–800. [PubMed http://dx.doi.org/10.1089/vbz.2009.0095](http://dx.doi.org/10.1089/vbz.2009.0095)
45. Kistler WM, Stallknecht DE, Deliberto TJ, Swafford S, Pedersen K, Van Why K, et al. Antibodies to avian influenza viruses in Canada geese (*Branta canadensis*): a potential surveillance tool? *J Wildl Dis.* 2012;48:1097–101. [PubMed http://dx.doi.org/10.7589/2011-02-046](http://dx.doi.org/10.7589/2011-02-046)
46. Hill NJ, Takekawa JY, Cardona CJ, Meixell BW, Ackerman JT, Runstadler JA, et al. Cross-seasonal patterns of avian influenza virus in breeding and wintering migratory birds: a flyway perspective. *Vector Borne Zoonotic Dis.* 2012;12:243–53. [PubMed http://dx.doi.org/10.1089/vbz.2010.0246](http://dx.doi.org/10.1089/vbz.2010.0246)
47. Ferro PJ, Khan O, Peterson MJ, Batchuluun D, Reddy SM, Lupiani B. Avian influenza virus surveillance in hunter-harvested waterfowl, Texas Coast, September 2009–January 2010. *Avian Dis.* 2012;56:1006–9. [PubMed http://dx.doi.org/10.1637/10194-041012-ResNote.1](http://dx.doi.org/10.1637/10194-041012-ResNote.1)
48. Reeves AB, Pearce JM, Ramey AM, Ely CR, Schmutz JA, Flint PL, et al. Genomic analysis of avian influenza viruses from waterfowl in western Alaska, USA. *J Wildl Dis.* 2013;49:600–10. [PubMed http://dx.doi.org/10.7589/2012-04-108](http://dx.doi.org/10.7589/2012-04-108)
49. Ely CR, Hall JS, Schmutz JA, Pearce JM, Terenzi J, Sedinger JS, et al. Evidence that life history characteristics of wild birds influence infection and exposure to influenza A viruses. *PLoS ONE.* 2013;8:e57614. [PubMed http://dx.doi.org/10.1371/journal.pone.0057614](http://dx.doi.org/10.1371/journal.pone.0057614)
50. Pedersen K, Swafford SR, DeLiberto TJ. Low pathogenicity avian influenza subtypes isolated from wild birds in the United States, 2006–2008. *Avian Dis.* 2010;54:405–10. [PubMed http://dx.doi.org/10.1637/8693-031309-Reg.1](http://dx.doi.org/10.1637/8693-031309-Reg.1)
51. Hildebrand E, Carstensen M, Butler E. Surveillance for highly pathogenic avian influenza in Minnesota’s migratory birds from 2006–2010. Summaries of wildlife research findings. St. Paul MN): Minnesota Department of Natural Resources; 2010. p. 203–11.

52. Buscaglia C. A survey for avian influenza from gulls on the coasts of the District of Pinamar and the Lagoon Salada Grande, General Madariaga, Argentina. *Avian Dis.* 2012;56:1017–20. [PubMed](#)
<http://dx.doi.org/10.1637/10147-040712-Reg.1>
53. Cumming GS, Caron A, Abolnik C, Cattoli G, Bruinzeel LW, Burger CE, et al. The ecology of influenza A viruses in wild birds in southern Africa. *EcoHealth.* 2011;8:4–13. [PubMed](#)
<http://dx.doi.org/10.1007/s10393-011-0684-z>
54. Gaidet N, Caron A, Cappelle J, Cumming GS, Balanca G, Hammoumi S, et al. Understanding the ecological drivers of avian influenza virus infection in wildfowl: a continental-scale study across Africa. *Proc Biol Sci.* 2012;279:1131–41. [PubMed](#) <http://dx.doi.org/10.1098/rspb.2011.1417>
55. Goujgoulova G, Petkova AM, Georgiev G. Avian influenza viruses isolated from mallards in Bulgaria. *Avian Dis.* 2010;54:450–2. [PubMed](#) <http://dx.doi.org/10.1637/8681-030709-ResNote.1>
56. Han Y, Hou G, Jiang W, Han C, Liu S, Chen J, et al. A survey of avian influenza in tree sparrows in China in 2011. *PLoS ONE.* 2012;7:e33092. [PubMed](#)
<http://dx.doi.org/10.1371/journal.pone.0033092>
57. Zhu G, Wang RJ, Xuan FJ, Daszak P, Anthony SJ, Zhang SY, et al. Characterization of recombinant H9N2 influenza viruses isolated from wild ducks in China. *Vet Microbiol.* 2013;166:327–36. [PubMed](#) <http://dx.doi.org/10.1016/j.vetmic.2013.05.013>
58. Montalvo-Corral M, Lopez-Robles G, Hernandez J. Avian influenza survey in migrating waterfowl in Sonora, Mexico. *Transbound Emerg Dis.* 2011;58:63–8. [PubMed](#)
<http://dx.doi.org/10.1111/j.1865-1682.2010.01182.x>
59. Williams RA, Segovia-Hinojosa K, Ghersi BM, Gonzaga V, Peterson AT, Montgomery JM. Avian influenza infections in nonmigrant land birds in Andean Peru. *J Wildl Dis.* 2012;48:910–7. [PubMed](#) <http://dx.doi.org/10.7589/2011-02-052>
60. Siengsan-Lamont J, Robertson I, Blacksell SD, Ellis T, Fenwick S, Saengchoowong S, et al. Virological and molecular epidemiological investigations into the role of wild birds in the epidemiology of influenza A/H5N1 in central Thailand. *Vet Microbiol.* 2011;148:213–8. [PubMed](#)
<http://dx.doi.org/10.1016/j.vetmic.2010.09.028>
61. Boynukara B, Gulhan T, Adizel O, Ilhan Z, Aksakal A, Durmus A, et al. Determination of avian influenza A viruses in some avian species in Van Lake Basin by real-time PCR, isolation and subtyping. *Journal of Animal and Veterinary Advances.* 2011;10:502–10.
<http://dx.doi.org/10.3923/javaa.2011.502.510>

62. El-Zoghby EF, Aly MM, Nasef SA, Hassan MK, Arafa AS, Selim AA, et al. Surveillance on A/H5N1 virus in domestic poultry and wild birds in Egypt. *Virology*. 2013;10:203. [PubMed](#)
<http://dx.doi.org/10.1186/1743-422X-10-203>
63. Soliman A, Saad M, Ellassal E, Amir E, Plathonoff C, Bahgat V, et al. Surveillance of avian influenza viruses in migratory birds in Egypt, 2003–09. *J Wildl Dis*. 2012;48:669–75. [PubMed](#)
<http://dx.doi.org/10.7589/0090-3558-48.3.669>
64. Lewis NS, Javakhishvili Z, Russell CA, Machabishvili A, Lexmond P, Verhagen JH, et al. Avian influenza virus surveillance in wild birds in Georgia: 2009–2011. *PLoS ONE*. 2013;8:e58534. [PubMed](#) <http://dx.doi.org/10.1371/journal.pone.0058534>
65. Odoom JK, Bel-Nono S, Rodgers D, Agbenohevi PG, Dafeamekpor CK, Sowa RM, et al. Troop education and avian influenza surveillance in military barracks in Ghana, 2011. *BMC Public Health*. 2012;12:957. [PubMed](#) <http://dx.doi.org/10.1186/1471-2458-12-957>
66. González-Reiche AS, Morales-Betoulle ME, Alvarez D, Betoulle J-L, Mueller ML, Sosa SM, et al. Influenza A viruses from wild birds in Guatemala belong to the North American lineage. *PLoS ONE*. 2012;7:e32873. [PubMed](#) <http://dx.doi.org/10.1371/journal.pone.0032873>
67. Pawar SD, Kale SD, Rawankar AS, Koratkar SS, Raut CG, Pande SA, et al. Avian influenza surveillance reveals presence of low pathogenic avian influenza viruses in poultry during 2009–2011 in the West Bengal State, India. *Virology*. 2012;9:151. [PubMed](#)
<http://dx.doi.org/10.1186/1743-422X-9-151>
68. Gilbert M, Jambal L, Karesh WB, Fine A, Shiilegdamba E, Dulam P, et al. Highly pathogenic avian influenza virus among wild birds in Mongolia. *PLoS ONE*. 2012;7:e44097. [PubMed](#)
<http://dx.doi.org/10.1371/journal.pone.0044097>
69. Sakoda Y, Sugar S, Batchluun D, Erdene-Ochir T-O, Okamatsu M, Isoda N, et al. Characterization of H5N1 highly pathogenic avian influenza virus strains isolated from migratory waterfowl in Mongolia on the way back from southern Asia to their northern territory. *Virology*. 2010;406:88–94. [PubMed](#) <http://dx.doi.org/10.1016/j.virol.2010.07.007>
70. Columba Teru V, Manu SA, Ahmed GI, Junaidu K, Newman S, Nyager J, et al. Situation-based survey of avian influenza viruses in possible “bridge” species of wild and domestic birds in Nigeria. *Influenza Res Treat*. 2012;2012:8567601. Epub 2012 Sep 2. doi: 10.1155/2012/567601.
71. Siddique N, Naeem K, Abbas MA, Ahmed Z, Malik SA. Sequence and phylogenetic analysis of highly pathogenic avian influenza H5N1 viruses isolated during 2006–2008 outbreaks in Pakistan

- reveals genetic diversity. *Virology*. 2012;9:300–13. [PubMed http://dx.doi.org/10.1186/1743-422X-9-300](http://dx.doi.org/10.1186/1743-422X-9-300)
72. Kulak MV, Ilinykh FA, Zaykovskaya AV, Epanchinzeva AV, Evstaphiev IL, Tovtunec NN, et al. Surveillance and identification of influenza A viruses in wild aquatic birds in the Crimea, Ukraine (2006–2008). *Avian Dis.* 2010;54:1086–90. [PubMed http://dx.doi.org/10.1637/9272-020510-ResNote.1](http://dx.doi.org/10.1637/9272-020510-ResNote.1)
73. Takakuwa H, Yamashiro T, Le MQ, Phuong LS, Ozaki H, Tsunekuni R, et al. The characterization of low pathogenic avian influenza viruses isolated from wild birds in northern Vietnam from 2006 to 2009. *Comp Immunol Microbiol Infect Dis.* 2013;36:581–90. [PubMed http://dx.doi.org/10.1016/j.cimid.2013.06.004](http://dx.doi.org/10.1016/j.cimid.2013.06.004)
74. Tinh TV, Gilbert M, Bunpapong N, Amonsin A, Nguyen DT, Doherty PF Jr, et al. Avian influenza viruses in wild land birds in northern Vietnam. *J Wildl Dis.* 2012;48:195–200. [PubMed http://dx.doi.org/10.7589/0090-3558-48.1.195](http://dx.doi.org/10.7589/0090-3558-48.1.195)
75. Simulundu E, Ishii A, Igarashi M, Mweene AS, Suzuki Y, Hang'ombe BM, et al. Characterization of influenza A viruses isolated from wild waterfowl in Zambia. *J Gen Virol.* 2011;92:1416–27. [PubMed http://dx.doi.org/10.1099/vir.0.030403-0](http://dx.doi.org/10.1099/vir.0.030403-0)
76. Khatun A, Giasuddin M, Islam KM, Khanom S, Samad MA, Islam MR, et al. Surveillance of avian influenza virus type A in semi-scavenging ducks in Bangladesh. *BMC Vet Res.* 2013;9:196. [PubMed http://dx.doi.org/10.1186/1746-6148-9-196](http://dx.doi.org/10.1186/1746-6148-9-196)
77. Theary R, San S, Davun H, Allal L, Lu HG. New outbreaks of H5N1 highly pathogenic avian influenza in domestic poultry and wild birds in Cambodia in 2011. *Avian Dis.* 2012;56:861–4. [PubMed http://dx.doi.org/10.1637/10195-041012-ResNote.1](http://dx.doi.org/10.1637/10195-041012-ResNote.1)
78. Ofula VO, Franklin AB, Root JJ, Sullivan HJ, Gichuki P, Makio A, et al. Detection of avian influenza viruses in wild waterbirds in the Rift Valley of Kenya using fecal sampling. *Vector Borne Zoonotic Dis.* 2013;13:394–400. [PubMed http://dx.doi.org/10.1089/vbz.2011.0926](http://dx.doi.org/10.1089/vbz.2011.0926)