

# *Rickettsia sibirica mongolitimonae* Infection, France, 2010–2014

Emmanouil Angelakis, Herve Richet,  
Didier Raoult

To further characterize human infections caused by *Rickettsia sibirica mongolitimonae*, we tested skin biopsy and swab samples and analyzed clinical, epidemiologic, and diagnostic characteristics of patients with a rickettsiosis. The most common (38%) indigenous species was *R. sibirica mongolitimonae*. Significantly more cases of *R. sibirica mongolitimonae* infection occurred during spring and summer.

Tickborne rickettsioses are zoonoses caused by spotted fever group (SFG) *Rickettsia* spp. (1). The first human infection with *R. sibirica mongolitimonae* was reported in France in 1996 (2). This patient had rope-like lymphangitis from the eschar to the draining lymph node, and *R. sibirica mongolitimonae* infection was thus named lymphangitis-associated rickettsiosis (3,4). Since then, other cases with or without rope-like lymphangitis have been described (5). Several SFG rickettsioses that have been considered nonpathogenic for decades are now associated with human infections, making these diseases useful as a paradigm for understanding emerging and reemerging infections (6). To further characterize human infections caused by *R. sibirica mongolitimonae*, we tested skin biopsy and swab samples and analyzed the clinical, epidemiologic, and diagnostic characteristics of patients with a rickettsiosis.

## The Study

During 2010–2014, we tested skin biopsy (7) and cutaneous swab samples from rickettsiosis inpatients and outpatients throughout France. These samples were received frozen or in transport media; when possible, serum samples were also collected and sent at room temperature. For patients with positive *Rickettsia* results, epidemiologic and clinical data were collected.

We extracted total genomic DNA from samples by using a QIAamp tissue kit (QIAGEN, Hilden, Germany). We screened samples for *Rickettsia* spp. by using a quantitative PCR assay selective for a 109-bp fragment of a hypothetical protein (8). For positive samples, PCR amplification and sequencing selective for the *gltA* and *ompA* genes were performed (8). Samples were cultured in human embryonic lung fibroblasts (9). All serum samples were tested by immunofluorescence assay for SFG rickettsial antigens and typhus

group rickettsiae (10). Student *t* or  $\chi^2$  tests were performed by using Epi Info version 6.0 software (Centers for Disease Control and Prevention, Atlanta, GA, USA). Means were compared by using analysis of variance or the Kruskal-Wallis test, on the basis of results of the Bartlett test for inequality of population variances. Proportions were compared by using the Mantel-Haenszel  $\chi^2$  or Fisher exact tests when the expected value of a cell was  $<0.05$ . *R. sibirica mongolitimonae* seasonality was assessed by using the autocorrelation module of PASW software version 17.02 (<http://www.spss.com/hk/statistics/>).  $p < 0.05$  was considered significant.

We classified patients as definitively having a rickettsiosis if direct evidence of rickettsial infection was found on culture or molecular assays. Of 465 patients examined, 91 (20%) were infected with *Rickettsia* spp., most commonly *R. africae* ( $n = 36$ , 40%), followed by *R. conorii* ( $n = 21$ , 23%), *R. sibirica mongolitimonae* ( $n = 20$ , 22%), and *R. slovaca* ( $n = 14$ , 15%). Two cases of *R. sibirica mongolitimonae* infection in France have been reported (11,12).

For patients infected with *R. sibirica mongolitimonae*, median age  $\pm$  SD (interquartile range) was  $43 \pm 21$  (2–70) years, and most (12, 60%) were male (online Technical Appendix Table 1, <http://wwwnc.cdc.gov/EID/article/22/5/14-1989-Techapp1.pdf>). The most common *Rickettsia* species in France was *R. sibirica mongolitimonae*. Only 1 patient mentioned recent travel to Spain; all others denied recent travel. Five patients mentioned recent outdoor activities, 8 mentioned frequent contact with dogs, and 1 mentioned contact with horses. A tick bite or tick handling was reported by 6 patients. An autocorrelation analysis revealed significant seasonality for *R. sibirica mongolitimonae* cases ( $p < 0.001$ ). Significantly more cases occurred during spring (April–June) (11 cases, 55%;  $p = 0.006$ ), followed by summer (July–September) (8 cases, 40%,  $p = 0.01$ ). One case occurred in October and none in winter.

The symptoms at disease onset included fever for all patients (duration 4–14 days), myalgia ( $n = 11$ , 55%), and headache ( $n = 3$ , 15%). Generalized maculopapular rash and an inoculation eschar developed in all patients. One patient had 3 eschars (buttocks, right hand, breast). A rope-like lymphangitis from the eschar to the draining lymph node was detected in 7 (35%) patients. One patient was admitted to an intensive care unit. For all 5 patients for whom an initial laboratory examination was available, increased liver enzymes (alanine aminotransferase, aspartate aminotransferase) and thrombocytopenia were found; 2 patients had hypoproteinemia. Oral doxycycline (7–14 days) was given to 19 patients; pristinamycin (7 days) was given to 1 patient. All outcomes were successful.

Author affiliation: Aix Marseille Université, Marseille, France

DOI: <http://dx.doi.org/10.3201/eid2205.141989>

**Table.** Epidemiologic and clinical characteristics of the main spotted fever group rickettsioses identified at the Unité de Recherche sur les Maladies Infectieuses et Tropicales Émergentes, Marseilles, France, 2010–2014\*

| Characteristic           | <i>Rickettsia africae</i> | <i>R. conorii</i>                                | <i>R. slovaca</i> | <i>R. sibirica mongolitimonae</i> |
|--------------------------|---------------------------|--------------------------------------------------|-------------------|-----------------------------------|
| No. cases                | 36                        | 21                                               | 14                | 20                                |
| Geographic location      | Zimbabwe and South Africa | Algeria, France, Morocco, Portugal, South Africa | France            | France, Spain                     |
| Median age ± SD (IQR), y | 58 ± 12 (31–80)           | 53 ± 18 (10–80)                                  | 36 ± 23 (6–65)    | 43 ± 21 (2–70)                    |
| Female sex               | 14 (39)                   | 7 (33)                                           | 9 (64)            | 8 (40)                            |
| Recent travel            | 36                        | 17                                               | 0                 | 1                                 |
| Clinical signs           |                           |                                                  |                   |                                   |
| Fever                    | 35 (97)                   | 21 (100)                                         | 5 (36)            | 20 (100)                          |
| Rash                     | 24 (67)                   | 20 (95)                                          | 3 (21)            | 19 (95)                           |
| Enlarged lymph nodes     | 15 (42)                   | 3 (14)                                           | 14 (100)          | 12 (60)                           |
| Lymphadenopathy location |                           |                                                  |                   |                                   |
| Cervical                 | 1 (3)                     | 3 (14)                                           | 14 (100)          | 5 (25)                            |
| Inguinal                 | 14 (39)                   | 0                                                | 0                 | 3 (15)                            |
| Axillary                 | 0                         | 0                                                | 0                 | 4 (20)                            |
| Eschar                   | 36 (100)                  | 18 (86)                                          | 14 (100)          | 20 (100)                          |
| Multiple eschars         | 13 (36)                   | 0                                                | 0                 | 3 (15)                            |
| Eschar location          |                           |                                                  |                   |                                   |
| Scalp                    | 0                         | 2 (10)                                           | 14 (100)          | 0                                 |
| Lower limbs              | 32 (89)                   | 3 (14)                                           | 0                 | 7 (33)                            |
| Upper limbs              | 2 (6)                     | 2 (10)                                           | 0                 | 4 (20)                            |
| Trunk                    | 2 (6)                     | 5 (24)                                           | 0                 | 3 (15)                            |
| Neck                     | 0                         | 0                                                | 0                 | 4 (20)                            |
| Lymphangitis             | 0                         | 0                                                | 0                 | 7 (35)                            |
| Treatment (duration, d)  |                           |                                                  |                   |                                   |
| Doxycycline              | 34 (1–20)                 | 21 (7–21)                                        | 12 (1–7)          | 19 (7–14)                         |
| Amoxicillin              | 2 (7)                     | None                                             | None              | None                              |
| Pristinamycin            | None                      | 1 (7)                                            | None              | 1 (7)                             |
| Azithromycin             | None                      | None                                             | 2 (4)             | None                              |

\*Values are no. (%) patients unless otherwise indicated. IQR, interquartile range.

An eschar swab sample was available for 13 patients (13), and a skin biopsy sample was available for 10; all samples were positive for *R. sibirica mongolitimonae*. An acute-phase serum sample was also available for 13 patients; results of serologic testing were positive for only 2 (15%). A convalescent-phase serum sample was available from 5 patients; results were positive for 4 (80%). A skin biopsy sample was also positive for *R. sibirica mongolitimonae* by culture.

Statistical comparison of the 4 rickettsioses (Table) showed that a recent travel history was more common among patients with *R. africae* infection ( $p < 0.001$ ). *R. slovaca* infection was associated with absence of fever or rash ( $p < 0.001$  for each). Multiple eschars were associated with *R. africae* infection ( $p < 0.001$ ). An eschar on the neck was a characteristic of infection with *R. sibirica mongolitimonae* ( $p = 0.002$ ); on the scalp, *R. slovaca* ( $p < 0.001$ ); on the trunk, *R. conorii* ( $p = 0.05$ ); and on the lower limbs, *R. africae* ( $p < 0.001$ ). For patients with rope-like lymphangitis, the probability of *R. sibirica mongolitimonae* infection was 100% ( $p < 0.001$ ). Cervical lymphadenitis was associated with *R. slovaca* ( $p < 0.001$ ), inguinal lymphadenitis with *R. africae* ( $p < 0.001$ ), and axillary lymphadenitis with *R. sibirica mongolitimonae* infection ( $p = 0.01$ ).

### Conclusions

*R. sibirica mongolitimonae* is considered a rare pathogen; only 30 cases of infection with this organism have been reported in Europe and Africa (online Technical Appendix Table 2), of which 11 patients had lymphangitis, 27 inoculation eschars, and 18 a rash. In agreement with previous authors, we found that the most common signs of *R. sibirica mongolitimonae* infection were fever and rash. The addition of rope-like lymphangitis cases to those in the literature revealed that 17 (35%) of patients with *R. sibirica mongolitimonae* infection had this manifestation. Ramos et al. proposed that the term lymphangitis-associated rickettsiosis may be unwarranted for *R. sibirica mongolitimonae* infection because it is not found in all patients infected with this organism and because other rickettsioses produce lymphangitis (14). However, only *R. sibirica mongolitimonae* infection is associated with rope-like lymphangitis extending from the eschar to the draining lymph node; to our knowledge, only 1 case of mild, local, but not rope-like lymphangitis in a patient with *R. africae* infection has been described (15). In accordance with previous reports from France and Spain (3,14), we found that *R. sibirica mongolitimonae* infection was seasonal and that most cases occurred in the spring and summer.

Our strategy for diagnosing *Rickettsia* spp. infection on the basis of skin biopsy and cutaneous swab samples modified our knowledge of the epidemiology of SFG rickettsioses in France. We provide evidence that *R. sibirica mongolitimonae* infection is a frequent rickettsiosis, probably more frequent than *R. conorii* infection, which for decades has been considered the most common *Rickettsia* species in France.

Dr. Angelakis is a medical microbiologist and researcher at the Unité de Recherche sur les Maladies Infectieuses et Tropicales Émergentes, Unités Mixtes de Recherche 6236, Centre Nationale de la Recherche Scientifique, Institut de Recherche pour le Développement, in Marseille. His research interests are zoonotic pathogens.

## References

1. Merhej V, Angelakis E, Socolovschi C, Raoult D. Genotyping, evolution and epidemiological findings of *Rickettsia* species. *Infect Genet Evol.* 2014;25:122–37. <http://dx.doi.org/10.1016/j.meegid.2014.03.014>
2. Raoult D, Brouqui P, Roux V. A new spotted-fever-group rickettsiosis. *Lancet.* 1996;348:412. [http://dx.doi.org/10.1016/S0140-6736\(05\)65037-4](http://dx.doi.org/10.1016/S0140-6736(05)65037-4)
3. Fournier PE, Gouriet F, Brouqui P, Lucht F, Raoult D. Lymphangitis-associated rickettsiosis, a new rickettsiosis caused by *Rickettsia sibirica mongolitimonae*: seven new cases and review of the literature. *Clin Infect Dis.* 2005;40:1435–44. <http://dx.doi.org/10.1086/429625>
4. Fournier PE, Tissot-Dupont H, Gallais H, Raoult D. *Rickettsia mongolitimonae*: a rare pathogen in France. *Emerg Infect Dis.* 2000;6:290–2. <http://dx.doi.org/10.3201/eid0603.000309>
5. Foissac M, Socolovschi C, Raoult D. Lymphangitis-associated rickettsiosis caused by *Rickettsia sibirica mongolitimonae*. *Ann Dermatol Venereol.* 2013;140:521–7. <http://dx.doi.org/10.1016/j.annder.2013.04.096>
6. Germanakis A, Chochlakis D, Angelakis E, Tselentis Y, Psaroulaki A. *Rickettsia aeschlimanii* infection in a man, Greece. *Emerg Infect Dis.* 2013;19:1176–7. <http://dx.doi.org/10.3201/eid1907.130232>
7. Angelakis E, Richet H, Rolain JM, La Scola B, Raoult D. Comparison of real-time quantitative PCR and culture for the diagnosis of emerging rickettsioses. *PLoS Negl Trop Dis.* 2012;6:e1540. <http://dx.doi.org/10.1371/journal.pntd.0001540>
8. Bechah Y, Socolovschi C, Raoult D. Identification of rickettsial infections by using cutaneous swab specimens and PCR. *Emerg Infect Dis.* 2011;17:83–6. <http://dx.doi.org/10.3201/eid1701.100854>
9. Gouriet F, Fenollar F, Patrice JY, Drancourt M, Raoult D. Use of shell-vial cell culture assay for isolation of bacteria from clinical specimens: 13 years of experience. *J Clin Microbiol.* 2005;43:4993–5002. <http://dx.doi.org/10.1128/JCM.43.10.4993-5002.2005>
10. Fournier PE, Jensenius M, Laferl H, Vene S, Raoult D. Kinetics of antibody responses in *Rickettsia africae* and *Rickettsia conorii* infections. *Clin Diagn Lab Immunol.* 2002;9:324–8.
11. Edouard S, Parola P, Socolovschi C, Davoust B, La Scola B, Raoult D. Clustered cases of *Rickettsia sibirica mongolitimonae* infection, France. *Emerg Infect Dis.* 2013;19:337–8. <http://dx.doi.org/10.3201/eid1902.120863>
12. Solary J, Socolovschi C, Aubry C, Brouqui P, Raoult D, Parola P. Detection of *Rickettsia sibirica mongolitimonae* by using cutaneous swab samples and quantitative PCR. *Emerg Infect Dis.* 2014;20:716–8. <http://dx.doi.org/10.3201/eid2004.130575>
13. Bechah Y, Socolovschi C, Raoult D. Identification of rickettsial infections by using cutaneous swab specimens and PCR. *Emerg Infect Dis.* 2011;17:83–6. <http://dx.doi.org/10.3201/eid1701.100854>
14. Ramos JM, Jado I, Padilla S, Masia M, Anda P, Gutierrez F. Human infection with *Rickettsia sibirica mongolitimonae*, Spain, 2007–2011. *Emerg Infect Dis.* 2013;19:267–9. <http://dx.doi.org/10.3201/eid1902.111706>
15. Althaus F, Greub G, Raoult D, Genton B. African tick-bite fever: a new entity in the differential diagnosis of multiple eschars in travelers. Description of five cases imported from South Africa to Switzerland. *Int J Infect Dis.* 2010;14(Suppl 3):e274–6. <http://dx.doi.org/10.1016/j.ijid.2009.11.021>

Address for correspondence: Didier Raoult, Faculté de Médecine, Université de la Méditerranée, URMITE, UMR CNRS 6236, IRD 198, Centre National de Référence, 27 Blvd Jean Moulin, Marseille 13005, France; email: didier.raoult@gmail.com

## EID Podcast: Louseborne Relapsing Fever in Europe

During June 9–September 30, 2015, five cases of louseborne relapsing fever were identified in Turin, Italy. All 5 cases were in young refugees from Somalia, 2 of whom had lived in Italy since 2011. This report seems to confirm the possibility of local transmission of louse-borne relapsing fever.



Visit our website to listen:  
<http://www2c.cdc.gov/podcasts/player.asp?f=8639774>

# Expanding Distribution of Lethal Amphibian Fungus *Batrachochytrium salamandrivorans* in Europe

## Technical Appendix

**Technical Appendix Table 1.** Field sites where *Bsal* was detected, sampled species, numbers of *Bsal*-positive and total sampled specimens\*

| Site no., location, and amphibian collected                        | No. <i>Bsal</i> -positive/total tested (year)       | Observed prevalence (Bayesian 95% credible intervals) | Remarks                                                                             |                                                  |
|--------------------------------------------------------------------|-----------------------------------------------------|-------------------------------------------------------|-------------------------------------------------------------------------------------|--------------------------------------------------|
| <b>The Netherlands</b>                                             |                                                     |                                                       |                                                                                     |                                                  |
| 1, Bunderbos, deciduous forest<br>Fire salamander                  | 3/3 (2010)                                          | 1.00 (0.42–1.00)                                      | Past mass deaths; 99.9% population decline (1997–2014)                              |                                                  |
|                                                                    | 1/1 (2011)                                          | 1.00                                                  |                                                                                     |                                                  |
|                                                                    | 1/1 (2012)                                          | 1.00                                                  |                                                                                     |                                                  |
|                                                                    | 0/3 (2014)                                          | 0 (0–0.61)                                            |                                                                                     |                                                  |
|                                                                    | 2/14 (2015)                                         | 0.14 (0.04–0.40)                                      |                                                                                     |                                                  |
|                                                                    | 0/1 (2016)                                          | 0                                                     |                                                                                     |                                                  |
|                                                                    | Alpine newt                                         | 1/1 (2013)                                            | 1.00                                                                                | Possibly declining (monitoring started in 2013)† |
|                                                                    | 1/39 (2014)                                         | 0.03 (0.01–0.13)                                      |                                                                                     |                                                  |
|                                                                    | 1/10 (2015)                                         | 0.10 (0.02–0.43)                                      |                                                                                     |                                                  |
|                                                                    | 0/6 (2016)                                          | 0 (0–0.43)                                            |                                                                                     |                                                  |
| 2, Putberg, deciduous forest<br>Smooth newt                        | 0/2 (2014)                                          | 0 (0–0.70)                                            | Possibly declining†                                                                 |                                                  |
|                                                                    | Alpine newt                                         | 0/10 (2014)                                           | 0 (0–0.31)                                                                          | Possibly declining†                              |
|                                                                    | 1/1‡ (2014)                                         | 1.00                                                  |                                                                                     |                                                  |
|                                                                    | 1/1‡ (2015)                                         | 1.00                                                  |                                                                                     |                                                  |
| 3, Meerssen, garden pond<br>Fire salamander                        | 0/1 (2015)                                          | 0                                                     | No evidence of decline§                                                             |                                                  |
|                                                                    | Smooth newt                                         | 4/43 (2015)                                           | 0.09 (0.04–0.21)                                                                    | No evidence of decline§                          |
|                                                                    | Alpine newt                                         | 0/9 (2015)                                            | 0 (0–0.30)                                                                          | No evidence of decline§                          |
| 4, Wormdal, clusters of natural ponds in nature conservation area¶ | Smooth newt                                         | 1/22 (2015)                                           | 0.05 (0.01–0.21)                                                                    | 87% decline(2000–2013)†#                         |
|                                                                    | Alpine newt                                         | 0/12 (2015)                                           | 0 (0–0.26)                                                                          | 96% decline (2000–2013)†#                        |
| 5, Pepinusbeekdal, extensive agriculture<br>Smooth newt            | 1/2‡ (2014)                                         | 0.50 (0.09–0.88)                                      | No evidence of decline†                                                             |                                                  |
|                                                                    | 6, Berg en Dal, garden pond<br>Alpine newt          | 12/12‡ (2015)                                         | 1.00 (0.74–1.00)                                                                    | Yearly mass deaths; species still present§       |
| 7, Vijlenerbosch, deciduous forest<br>Alpine newt                  |                                                     | 0/1 (2013)                                            | 0                                                                                   | No evidence of decline§                          |
|                                                                    |                                                     | 0/30 (2014)                                           | 0 (0–0.11)                                                                          |                                                  |
|                                                                    | 1/18 (2015)                                         | 0.05 (0.02–0.24)                                      |                                                                                     |                                                  |
| Smooth newt                                                        | 0/8 (2014)                                          | 0 (0–0.31)                                            | No evidence of decline§                                                             |                                                  |
|                                                                    | 0/11 (2015)                                         | 0 (0–0.26)                                            |                                                                                     |                                                  |
| Palmate newt                                                       | 0/1 (2014)                                          | 0                                                     | No evidence of decline§                                                             |                                                  |
|                                                                    | 0/9 (2015)                                          | 0 (0–0.30)                                            |                                                                                     |                                                  |
| <b>Belgium</b>                                                     |                                                     |                                                       |                                                                                     |                                                  |
| 8, Eupen, deciduous forest<br>Fire salamander                      | 1/2 (2013)                                          | 0.50 (0.09–0.88)                                      | Deaths, probably fire salamanders severely declining, no monitoring trend available |                                                  |
|                                                                    | 9, Robertville, deciduous forest<br>Fire salamander | 16/30 (2014)                                          | 0.53 (0.36–0.69)                                                                    | Deaths, severe decline, monitoring ongoing       |
| 10, Liège, deciduous forest                                        |                                                     |                                                       |                                                                                     |                                                  |

| Site no., location, and amphibian collected | No. <i>Bsal</i> -positive/total tested (year) | Observed prevalence (Bayesian 95% credible intervals) | Remarks                                                 |
|---------------------------------------------|-----------------------------------------------|-------------------------------------------------------|---------------------------------------------------------|
| Fire salamander                             | 5/5 (2014)                                    | 1.00 (0.55–1.00)                                      | Deaths                                                  |
| 11, Duffel, garden pond                     |                                               |                                                       |                                                         |
| Alpine newt                                 | 2/30‡ (2015)                                  | 0.07 (0.02–0.22)                                      | 2 dead in fyke; no evidence of decline                  |
| Smooth newt                                 | 0/16 (2015)                                   | 0 (0–0.20)                                            | No deaths; no evidence of decline                       |
| Germany                                     |                                               |                                                       |                                                         |
| 12, Weisse Wehe, deciduous forest           |                                               |                                                       |                                                         |
| Fire salamander                             | 4/11‡ (2015)                                  | 0.36 (0.15–0.65)                                      | No evidence of decline†                                 |
| 13, Solchbachtal, mixed forest              |                                               |                                                       |                                                         |
| Fire salamander                             | 0/2 (2014)                                    | 0 (0–0.70)                                            | Decreased newts and salamanders§                        |
|                                             | 1/51 (2015)                                   | 0.02 (0.01–0.10)                                      |                                                         |
| Palmate newt                                | 0/19(2014)                                    | 0 (0–0.18)                                            | Decreased newts and salamanders§                        |
| Alpine newt                                 | 0/5(2014)                                     | 0 (0–0.44)                                            | Decreased newts and salamanders§                        |
| 14, Belgenbachtal, mixed forest             |                                               |                                                       |                                                         |
| Fire salamander                             | 21/22‡ (2015)                                 | 0.96 (0.79–0.99)                                      | Remarkable deaths (16 dead), noted only since Nov 2015† |

\**Bsal*, *Batrachochytrium salamandrivorans*. Data provide an overview of novel information and previously published data. Site numbers correspond to those on map (Figure).

†Population monitored.

‡Includes individual(s) found dead by chance.

§Anecdotal reports.

¶At this site, crested newts and smooth newts decreased with similar percentages over the same period (–96%; –94%, respectively).

#[http://www.ravon.nl/EID\\_Sl\\_Spitzen\\_et\\_al\\_2016](http://www.ravon.nl/EID_Sl_Spitzen_et_al_2016).

**Technical Appendix Table 2.** Field sites studied where *Bsal* was not detected, number of sampled species and specimens\*

| Site no., location, and amphibian collected | Number of specimens tested (year) | Observed prevalence (Bayesian 95% credible intervals) | Remarks                                                                                              |
|---------------------------------------------|-----------------------------------|-------------------------------------------------------|------------------------------------------------------------------------------------------------------|
| Belgium                                     |                                   |                                                       |                                                                                                      |
| 15, Nerenbos, deciduous forest              |                                   |                                                       |                                                                                                      |
| Fire salamander                             | 30 (2015)                         | 0 (0–0.11)                                            | No evidence of decline†                                                                              |
| 16, Heilig Geestgoed, deciduous forest      |                                   |                                                       |                                                                                                      |
| Fire salamander                             | 30 (2015)                         | 0 (0–0.11)                                            | No evidence of decline†                                                                              |
| 17, Kasteel van Horst, deciduous forest     |                                   |                                                       |                                                                                                      |
| Fire salamander                             | 30 (2015)                         | 0 (0–0.11)                                            | No evidence of decline†                                                                              |
| 18, Smetledebos, deciduous forest           |                                   |                                                       |                                                                                                      |
| Fire salamander                             | 30 (2015)                         | 0 (0–0.11)                                            | No evidence of decline†                                                                              |
| 19, Kluisbos, deciduous forest              |                                   |                                                       |                                                                                                      |
| Fire salamander                             | 30 (2015)                         | 0 (0–0.11)                                            | No evidence of decline†                                                                              |
| 20, Hallerbos, deciduous forest             |                                   |                                                       |                                                                                                      |
| Fire salamander                             | 30 (2015)                         | 0 (0–0.11)                                            | No evidence of decline†                                                                              |
| 21, Buggenhoutbos, deciduous forest         |                                   |                                                       |                                                                                                      |
| Fire salamander                             | 30 (2015)                         | 0 (0–0.11)                                            | No evidence of decline†                                                                              |
| 22, Raspaillebos, deciduous forest          |                                   |                                                       |                                                                                                      |
| Fire salamander                             | 30 (2015)                         | 0 (0–0.11)                                            | No evidence of decline†                                                                              |
| 23, Haeyesbos, deciduous forest             |                                   |                                                       |                                                                                                      |
| Fire salamander                             | 30 (2015)                         | 0 (0–0.11)                                            | No evidence of decline†                                                                              |
| 24, t Burreken, deciduous forest            |                                   |                                                       |                                                                                                      |
| Fire salamander                             | 30 (2015)                         | 0 (0–0.11)                                            | No evidence of decline†                                                                              |
| Germany                                     |                                   |                                                       |                                                                                                      |
| 25, Lamersiefen, deciduous forest           |                                   |                                                       |                                                                                                      |
| Fire salamander                             | 17 (2014)                         | 0 (0–0.19)                                            | No evidence of decline†                                                                              |
|                                             | 32 (2015)                         | 0 (0–0.11)                                            |                                                                                                      |
| 26, Fischbach, deciduous forest             |                                   |                                                       |                                                                                                      |
| Fire salamander                             | 36 (2014)                         | 0 (0–0.09)                                            | No evidence of decline; 3 dead-found specimens Tested negative for <i>Bsal</i> via histology (2014)† |
|                                             | 51 (2015)                         | 0 (0–0.07)                                            |                                                                                                      |
| Alpine newt                                 | 1 (2015)                          | 0                                                     | Live-studied specimen by chance; no evidence of decline                                              |

| Site no., location, and amphibian collected                      | Number of specimens tested (year) | Observed prevalence (Bayesian 95% credible intervals) | Remarks                                                 |
|------------------------------------------------------------------|-----------------------------------|-------------------------------------------------------|---------------------------------------------------------|
| Palmate newt                                                     | 1 (2015)                          | 0                                                     | Live-studied specimen by chance; no evidence of decline |
| 27, Kallerbach, deciduous forest<br>Fire salamander              | 24(2015)                          | 0 (0–0.15)                                            | No evidence of decline†                                 |
| 28, Rosbach, deciduous forest<br>Fire salamander                 | 47 (2015)                         | 0 (0–0.07)                                            | No evidence of decline†                                 |
| 29, Zweifallshammer, deciduous forest<br>Fire salamander         | 41 (2015)                         | 0 (0–0.08)                                            | No evidence of decline†                                 |
| 30, Peterbach, mixed forest<br>Palmate newt                      | 12 (2014)                         | 0 (0–0.26)                                            | No evidence of decline                                  |
| Alpine newt                                                      | 4 (2014)                          | 0 (0–0.52)                                            | No evidence of decline                                  |
| 31, Haftenbach, deciduous forest<br>Fire salamander              | 46 (2015)                         | 0 (0–0.08)                                            | No evidence of decline†                                 |
| 32, Sauerbach, deciduous forest<br>Fire salamander               | 22 (2015)                         | 0 (0–0.15)                                            | No evidence of decline†                                 |
| Alpine newt                                                      | 1 (2015)                          | 0,00                                                  | No evidence of decline                                  |
| 33, Härtgessief, deciduous forest<br>Fire salamander             | 15 (2014)                         | 0 (0–0.19)                                            | Strong evidence of decline†                             |
| 34, Kottenforst, deciduous forest<br>Fire salamander             | 51 (2015)                         | (0–0.07)                                              | No evidence of decline                                  |
| 35, Großkampenber, mixed forest<br>Alpine newt                   | 4 (2015)                          | 0 (0–0.52)                                            | No evidence of decline                                  |
| Palmate newt                                                     | 1 (2015)                          | 0                                                     | No evidence of decline                                  |
| 36, Lützkampen -mixed forest<br>Alpine newt                      | 8 (2015)                          | 0 (0–0.31)                                            | No evidence of decline                                  |
| 37, Ferschweiler- mixed forest<br>Alpine newt                    | 2 (2015)                          | 0 (0–0.70)                                            | No evidence of decline                                  |
| Palmate newt                                                     | 8 (2015)                          | 0 (0–0.31)                                            | No evidence of decline                                  |
| 38, Ernzen, mixed forest<br>Fire salamander                      | 4 (2015)                          | 0 (0–0.52)                                            | No evidence of decline†                                 |
| The Netherlands                                                  |                                   |                                                       |                                                         |
| 39, Moerveld surroundings (A), Bunderbos vicinity<br>Alpine newt | 13 (2015)                         | 0 (0–0.22)                                            | No evidence of decline‡                                 |
| 40, Moerveld surroundings (B), Bunderbos vicinity<br>Alpine newt | 34 (2015)                         | 0 (0–0.11)                                            | No evidence of decline‡                                 |
| 41, Snijdersbergweg 21, garden pond<br>Alpine newt               | 60 (2015)                         | 0 (0–0.06)                                            | No evidence of decline‡                                 |
| 42, Mevr van der Meijstraat 12, garden pond<br>Alpine newt       | 19 (2015)                         | 0 (0–0.18)                                            | No evidence of decline‡                                 |
| 43, Mevr van der Meijstraat 20, garden pond<br>Alpine newt       | 17 (2015)                         | 0 (0–0.19)                                            | No evidence of decline‡                                 |
| 44, Snijdersbergweg 20, 2 garden ponds<br>Alpine newt            | 30 (2015)                         | 0 (0–0.11)                                            | No evidence of decline‡                                 |
| 45, Snijdersbergweg 23b, garden pond<br>Alpine newt              | 15 (2015)                         | 0 (0–0.19)                                            | No evidence of decline‡                                 |
| 46, Broekhoven, garden pond<br>Fire salamander                   | 2 (2015)                          | 0 (0–0.70)                                            | No evidence of decline‡                                 |
| 47, Meerssen, deciduous forest<br>Fire salamander                | 57 (2013)                         | 0 (0–0.06)                                            | No deaths; no evidence of decline†                      |
|                                                                  | 43 (2014)                         | 0 (0–0.08)                                            |                                                         |
|                                                                  | 29 (2015)                         | 0 (0–0.11)                                            |                                                         |
|                                                                  | 2 (2016)                          | 0 (0–0.70)                                            |                                                         |
| 48, Carisberg, deciduous forest<br>Alpine newt                   | 8 (2014)                          | 0 (0–0.31)                                            | No information available                                |
| Palmate newt                                                     | 23 (2014)                         | 0 (0–0.14)                                            | No information available                                |
| Smooth newt                                                      | 2 (2014)                          | 0 (0–0.70)                                            | No information available                                |
| Additional far-out sites (Germany)                               |                                   |                                                       |                                                         |
| N.S., Solling, deciduous forest<br>Fire salamander               | 23 (2015)                         | 0 (0–0.14)                                            | No evidence of decline‡                                 |
| N.S., Ilsenburg, deciduous forest<br>Fire salamander             | 8 (2015)                          | 0 (0–0.31)                                            | No evidence of decline‡                                 |
| N.S., Lelm, deciduous forest<br>Alpine newt                      | 57 (2015)                         | 0 (0–0.06)                                            | No evidence of decline‡                                 |
| Palmate newt                                                     | 6 (2015)                          | 0 (0–0.43)                                            | No evidence of decline‡                                 |

| Site no., location, and amphibian collected                    | Number of specimens tested (year) | Observed prevalence (Bayesian 95% credible intervals) | Remarks                 |
|----------------------------------------------------------------|-----------------------------------|-------------------------------------------------------|-------------------------|
| Smooth newt                                                    | 16 (2015)                         | 0 (0–0.20)                                            | No evidence of decline‡ |
| Crested newt                                                   | 29 (2015)                         | 0 (0–0.11)                                            | No evidence of decline‡ |
| N.S., Kleiwiesen, exposed ponds surrounded by deciduous forest |                                   |                                                       |                         |
| Alpine newt                                                    | 27 (2015)                         | 0 (0–0.13)                                            | No evidence of decline‡ |
| Smooth newt                                                    | 117 (2015)                        | 0 (0–0.03)                                            | No evidence of decline‡ |
| Crested newt                                                   | 27 (2015)                         | 0 (0–0.13)                                            | No evidence of decline‡ |
| N.S., Waldecker Schlossgrund, deciduous forest                 |                                   |                                                       |                         |
| Fire salamander                                                | 22 (2015)                         | 0 (0–0.15)                                            | No evidence of decline‡ |
| N.S., Closewitz, exposed ponds surrounded by deciduous forest  |                                   |                                                       |                         |
| Crested newt                                                   | 23 (2015)                         | 0 (0–0.14)                                            | No evidence of decline‡ |
| Additional far-out site (the Netherlands)                      |                                   |                                                       |                         |
| N.S., Veluwe, deciduous forest                                 |                                   |                                                       |                         |
| Italian crested newt                                           | 0 (2015)                          | 0 (0–0.11)                                            | No evidence of decline‡ |

\**Bsal*, *Batrachochytrium salamandrivorans*; N.S., not shown on map (Figure). Data provide an overview of novel information and previously published data. Site numbers correspond to those on map (Figure).

†Population monitored.

‡Anecdotal report.