

3. Jordan MB, Allen CE, Weitzman S, Filipovich AH, McClain KL. How I treat hemophagocytic lymphohistiocytosis. *Blood*. 2011;118:4041–52. <http://dx.doi.org/10.1182/blood-2011-03-278127>
4. Wheat LJ, Freifeld AG, Kleiman MB, Baddley JW, McKinsey DS, Loyd JE, et al. Clinical practice guidelines for the management of patients with histoplasmosis: 2007 update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2007;45:807–25. <http://dx.doi.org/10.1086/521259>
5. Townsend JL, Shanbhag S, Hancock J, Bowman K, Nijhawan AE. Histoplasmosis-induced hemophagocytic syndrome: a case series and review of the literature. *Open Forum Infect Dis*. 2015;2:ofv055. PMID: 26380347
6. Henter JI, Horne A, Arico M, Egeler RM, Filipovich AH, Imashuku S, et al. HLH-2004: diagnostic and therapeutic guidelines for hemophagocytic lymphohistiocytosis. *Pediatr Blood Cancer*. 2007;48:124–31. <http://dx.doi.org/10.1002/pbc.21039>
7. Trottestam H, Horne A, Arico M, Egeler RM, Filipovich AH, Gadner H, et al. Chemoimmunotherapy for hemophagocytic lymphohistiocytosis: long-term results of the HLH-94 treatment protocol. *Blood*. 2011;118:4577–84. <http://dx.doi.org/10.1182/blood-2011-06-356261>
8. Majluf-Cruz AS, Hurtado Monroy R, Souto-Meirino C, del Rio Chiriboga C, Simon J. Hemophagocytic syndrome associated with histoplasmosis in the acquired immunodeficiency syndrome: description of 3 cases and review of the literature [in Spanish]. *Sangre (Barc)*. 1993;38:51–5.
9. Keller FG, Kurtzberg J. Disseminated histoplasmosis: a cause of infection-associated hemophagocytic syndrome. *Am J Pediatr Hematol Oncol*. 1994;16:368–71.
10. Sanchez A, Celaya AK, Victorio A. Histoplasmosis-associated hemophagocytic syndrome: a case report. *AIDS Read*. 2007;17:496–9.

Address for correspondence: Joshua Wolf, St. Jude Children's Research Hospital, Department of Infectious Diseases, 262 Danny Thomas Pl, Memphis, TN 38105, USA; email: joshua.wolf@stjude.org

Novel Avian Influenza A(H5N8) Viruses in Migratory Birds, China, 2013–2014

Li-Chen Zhou,¹ Jing Liu,¹ En-Le Pei,¹ Wen-Jie Xue,¹ Jia-Min Lyu, Yin-Ting Cai, Di Wu, Wei Wu, Yu-Yi Liu, Hui-Yu Jin, Yu-Wei Gao, Zheng-Huan Wang, Tian-Hou Wang

Author affiliations: Laboratory of Wildlife Epidemic Diseases, Shanghai Key Laboratory for Urban Ecological Processes and Eco-Restoration, School of Life Sciences, East China Normal University, Shanghai, China (L.-C. Zhou, J. Liu, J.-M. Lyu, Z.-H. Wang, T.-H. Wang); Shanghai Municipal Agency of Wildlife Conservation, Shanghai (E.-L. Pei, D. Wu, H.-Y. Jin); Chongming Dongtan National Nature Reserve, Shanghai (W.-J. Xue, W. Wu); Jiuduansha Wetland National Nature Reserve, Shanghai, China (Y.-T. Cai) Wildlife Conservation Section, Shanghai Municipal

Forestry Bureau, Shanghai (Y.-Y. Liu); Research Center of Wildlife Disease, Key Laboratory of Jilin Province for Zoonosis Prevention and Control, Military Veterinary Research Institute of Academy of Military Medical Sciences, Changchun, China (Y.-W. Gao)

DOI: <http://dx.doi.org/10.3201/eid2206.151754>

To the Editor: Novel highly pathogenic avian influenza (HPAI) A(H5N8) virus infections were first detected in poultry in eastern China in 2010 (1); the virus caused outbreaks in South Korea and Japan in 2014 (2) and reached Europe and North America by early 2015 (3–6). Phylogenetic analysis indicated that novel HPAI subtype H5N8 viruses might have originated in China and then circulated in East Asia countries and that the global geographic dissemination of this virus was strongly associated with the migration of wild birds (7). However, the role of migratory birds in the initial introduction and spread of novel H5N8 strains in China and other countries in the region is unclear. Shanghai, located at the Yangtze River estuary on the eastern coast of China, is a crucial stopover for migratory birds in East Asia. We report the presence of novel H5N8 strains from migratory birds sampled in Shanghai from October 2013 through December 2014.

A total of 26 novel H5N8 viruses were detected from migratory ducks and curlews captured and swabbed during their wintering period at the coastal wetlands of Shanghai. We collected 19 H5N8 viruses from 16 common teals (*Anas crecca*), 2 falcated ducks (*A. falcata*), and 1 spot-billed duck (*A. poecilorhyncha*) sampled in 2013 and 7 viruses from Eurasian curlews (*Numenius arquata*) sampled in 2014. Common teals were also found to be infected with subtype H5N1, detected by N1 gene fragments in 3 mixed-infection and 2 single-infection samples (online Technical Appendix, <http://wwwnc.cdc.gov/EID/article/22/6/15-1754-Techapp1.pdf>). Sequences from this study were deposited in GenBank (accession nos. KT936635–KT936716).

Homology BLAST (<http://blast.st-va.ncbi.nlm.nih.gov/Blast.cgi>) searches showed that H5 and N8 genes of 18 influenza A(H5N8) viruses in ducks had $\geq 98\%$ similarity to H5N8 isolates W24 and 6D18 detected in poultry in Zhejiang Province (2), adjacent to Shanghai. The H5 gene in A/common teal/Shanghai/1108-1/2013(mixed) (PD1108-1) was 96% related to low pathogenicity avian influenza (LPAI) subtype H5N1 isolated from a European teal sampled in Russia in 2011 (GenBank accession no. KF462362). Of the 7 viruses from curlews, H5 and N8 isolates were closely related to isolates H68 and H297 from wild ducks reported in South Korea in early 2014 (8). Matrix genes of all novel subtype H5N8 viruses were closely related (95%–99%) to isolates from China (S11090, W24), Japan (156), and South Korea (Gochang1, S005) (online Technical Appendix Table).

¹These authors contributed equally to this article.




Figure. Phylogenetic tree of the hemagglutinin (HA) genes of influenza A subtype H5 viruses from wild birds of Shanghai, China, 2013–2014. Boldface indicates viruses from this study; representative isolates are underlined and referred to in abbreviated form in brackets. A total of 109 HA gene sequences ($\geq 1,594$ nt) were used for tree reconstruction. Representative strains and clades are recommended by WHO/OIE/FAO H5N1 Evolution Working Group and were retrieved from Influenza Virus Resource Database (<http://www.ncbi.nlm.nih.gov/genomes/FLU/Database/select.cgi>) and GISAID's EpiFluTM Database (<http://platform.gisaid.org/epi3/frontend>). The phylogenetic tree was constructed by using the maximum likelihood method based on the general time reversible model with bootstrap analysis (100 replicates), by MEGA version 6 (<http://www.megasoftware.net/>). Bootstrap values $\geq 50\%$ are shown. Scale bar indicates nucleotide substitutions per site. LPAI, low pathogenicity avian influenza. HPAI, highly pathogenic avian influenza.

Phylogenetic analysis of HPAI H5 descendants of A/goose/Guangdong/1/1996(H5N1) showed that clade 2.3.4.4 (10) H5N8 viruses fall into 2 distinct groups, closely related to group A (Buan2-like) and group B (Gochang1-like) (8). The hemagglutinin (HA) genes of the 18 subtype H5N8 viruses from ducks in 2013 shared a protease cleavage site motif of REKRRKR/GLF and the sequence cluster with H5N8 viruses from eastern China poultry (W24, 6D18) (2) and Korean group B isolates (Gochang1, H52) (8,9) to form group B. The HA genes from all 7 H5N8 isolates from curlews in 2014 had a protease cleavage site motif of RERRRKR/GLF and the sequence cluster, along with Korean group A (Buan2-like) (8,9), European (3,4), and North American (7) H5N8 lineage viruses to form group A. The HA from PD1108-1 had a cleavage site motif (RE–TR/GLF) characteristic of LPAI HA, and its sequence clustered with the Eurasian LPAI H5 lineage (Figure).

According to the sampling dates, the identification of the 18 group B H5N8 isolates from Shanghai was the earliest detection of HPAI H5N8 virus in wild birds in East Asia, before the first reported outbreak in South Korea in January 2014. Although poultry isolates from China obtained during the same period were phylogenetically clustered with group A (Figure), no group A viruses were detected in wild birds during the 2013–2014 wintering season in China. Notably, 2 of the group A Chinese poultry isolates (Q1 and s13124) have the HA cleavage site motifs of group B. Their topologically basal positions in group A (Figure) implied the connection between the 2 groups. Eurasian curlews are widely distributed in the Northern Hemisphere, including Europe, Siberia, Japan, the Korean Peninsula, and China (<http://ibc.lynxeds.com/species/eurasian-curlew-numenius-arquata>). Populations wintering in Shanghai have overlapped migratory routes and habitat distribution with duck species in East Asia (Shanghai Chongming Dongtan National Nature Reserve, unpub. data), which suggested possible transmission routes through overlapped habitats in their northern territory (breeding areas) or close contacts among these species. These data support the theory that asymptomatic migratory birds may have played a role in geographic dissemination of HPAI subtype H5N8 and facilitation of viral evolution and reassortment. Moreover, that HPAI subtypes H5N1 and H5N8 co-infected and co-circulated in migratory ducks suggests that rapid and active mutation and reassortment of H5 subtypes may take place in these hosts. Therefore, to monitor and then control the epidemics of H5 subtype viruses, it is urgent that more intensive surveillance be carried out in poultry and wild birds and that information be promptly shared among countries.

Acknowledgment

We thank Shanghai Pudong Forestry Station for field sampling assistance.

This work was supported by Shanghai Wildlife Epidemic Disease Monitoring Program (G061255), Shanghai Science and Technology Committee Project (2013QLG001).

References

- Zhao K, Gu M, Zhong L, Duan Z, Zhang Y, Zhu Y, et al. Characterization of three H5N5 and one H5N8 highly pathogenic avian influenza viruses in China. *Vet Microbiol.* 2013; 163:351–7. <http://dx.doi.org/10.1016/j.vetmic.2012.12.025>
- Wu H, Peng X, Xu L, Jin C, Cheng L, Lu X, et al. Novel reassortant influenza A(H5N8) viruses in domestic ducks, eastern China. *Emerg Infect Dis.* 2014;20:1315–8. <http://dx.doi.org/10.3201/eid2008.140339>
- Bouwstra R, Heutink R, Bossers A, Harders F, Koch G, Elbers A. Full-genome sequence of influenza A(H5N8) virus in poultry linked to sequences of strains from Asia, the Netherlands, 2014. *Emerg Infect Dis.* 2015;21:872–4. <http://dx.doi.org/10.3201/eid2105.141839>
- Harder T, Maurer-Stroh S, Pohlmann A, Starick E, Horeth-Bontgen D, Albrecht K, et al. Influenza A(H5N8) virus similar to strain in Korea causing highly pathogenic avian influenza in Germany. *Emerg Infect Dis.* 2015;21:860–3. <http://dx.doi.org/10.3201/eid2105.141897>
- Hanna A, Banks J, Marston DA, Ellis RJ, Brookes SM, Brown IH. Genetic characterization of highly pathogenic avian influenza (H5N8) virus from domestic ducks, England, November 2014. *Emerg Infect Dis.* 2015;21:879–82. <http://dx.doi.org/10.3201/eid2105.141954>
- Ip HS, Torchetti MK, Crespo R, Kohrs P, DeBruyn P, Mansfield KG, et al. Novel Eurasian highly pathogenic avian influenza A H5 viruses in wild birds, Washington, USA, 2014. *Emerg Infect Dis.* 2015;21:886–90. <http://dx.doi.org/10.3201/eid2105.142020>
- Lee DH, Torchetti MK, Winker K, Ip HS, Song CS, Swayne DE. Intercontinental spread of Asian-origin H5N8 to North America through Beringia by migratory birds. *J Virol.* 2015;89:6521–4. <http://dx.doi.org/10.1128/JVI.00728-15>
- Jeong J, Kang HM, Lee EK, Song BM, Kwon YK, Kim HR, et al. Highly pathogenic avian influenza virus (H5N8) in domestic poultry and its relationship with migratory birds in South Korea during 2014. *Vet Microbiol.* 2014;173:249–57. <http://dx.doi.org/10.1016/j.vetmic.2014.08.002>
- Lee YJ, Kang H, Lee E, Song B, Jeong J, Kwon Y, et al. Novel reassortant influenza A(H5N8) viruses, South Korea, 2014. *Emerg Infect Dis.* 2014;20:1087–9. <http://dx.doi.org/10.3201/eid2006.140233>
- World Health Organization. Evolution of the influenza A(H5) haemagglutinin: WHO/OIE/FAO H5 Working Group reports a new clade designated 2.3.4.4. 2015. [cited 2016 Feb 2]. http://www.who.int/influenza/gisrs_laboratory/h5_nomenclature_clade2344/en/

Address for correspondence: Zheng-Huan Wang or Tian-Hou Wang, Laboratory of Wildlife Epidemic Diseases, Shanghai Key Lab for Urban Ecological Processes and Eco-Restoration, School of Life Sciences, East China Normal University, No. 3663, North Zhongshan Rd, Shanghai, China; email: zhwang@bio.ecnu.edu.cn or thwang@bio.ecnu.edu.cn

Novel Avian Influenza A(H5N8) Viruses in Migratory Birds, China, 2013–2014

Technical Appendix

Materials and Methods

Wild Bird Surveillance and Sampling

Wild birds were captured and sampled with the permission and supervision of Shanghai Wild Life Conservation and Management Office. Swabs (1 oropharyngeal and 1 cloacal of each bird) were taken and transported in viral transport medium to the laboratory at 4°C for downstream molecular diagnosis within 6 hours.

Virus Detection

Viral RNA was extracted from viral transport medium by using MagMAX™ Pathogen RNA/DNA Kit (Applied Biosystems, Foster City, CA, USA) on a Magmax-96 Express (Applied Biosystems) according to the manufacturer's instructions. Real time reverse transcription-PCR was performed with matrix gene-specific probed primers to detect the presence of avian influenza viruses according to the protocol of the Chinese Center for Disease Control and Prevention (1). The results were further confirmed by sequencing the PCR products.

Subtype Identification and Gene Sequencing

RNA of viral-positive samples were reverse-transcribed by PrimeScript II Reverse Transcriptase Kit (TaKaRa, Biotechnology [Dalian] Co., Ltd, Dalian, China) according to the kit protocol. PCR amplifications were performed for subtyping of hemagglutinin (HA) and neuraminidase, and gene segments were amplified by using the previously published primers (2–

4) and primers designed in this study (on request). PCR products were gel-purified by using the QIAquick gel extraction kit (QIAGEN, Valencia, CA, USA) and were sequenced with the BigDye terminator kit (Applied Biosystems) on an ABI 3730 (Applied Biosystems). Sequences were assembled and edited with the Lasergene sequence analysis software package (<http://www.dnastar.com/t-dnastar-lasergene.aspx>). Primers for sequencing were similar to those used for amplification.

Phylogenetic Analysis

A phylogenetic tree was constructed of the HA genes of H5 viruses in wild birds of Shanghai, China, 2013–2014 (see text Figure). The viruses of this study are indicated in boldface. Representative isolates are underlined and referred to in abbreviation form in brackets. A total of 109 HA gene sequences ($\geq 1,594$ nt) were used for tree reconstruction. Representative strains and clades are recommended by WHO/OIE/FAO H5N1 Evolution Working Group and were retrieved from Influenza Virus Resource Database (<http://www.ncbi.nlm.nih.gov/genomes/FLU/Database/select.cgi>) and GISAID's EpiFluTM Database (<http://platform.gisaid.org/epi3/frontend>). The phylogenetic tree was constructed by using the maximum likelihood method based on the general time reversible model (best fit model: general time reversible + I + R, computed by jModelTest 2.1.4 (5,6) with bootstrap analysis (100 replicates), by MEGA v. 6 (7). Scale bar indicates nucleotide substitutions per site. Bootstrap values $\geq 50\%$ are shown.

References

1. Chinese Center for Disease Control and Prevention. Chinese National Influenza Center Standard Operation Protocol. Beijing: The Center; 2007. p. 101–25.
2. Hoffmann E, Stech J, Guan Y, Webster RG, Perez DR. Universal primer set for the full-length amplification of all influenza A viruses. Arch Virol. 2001;146:2275–89. [PubMed](https://pubmed.ncbi.nlm.nih.gov/11511111/)
<http://dx.doi.org/10.1007/s007050170002>

3. Tsukamoto K, Ashizawa H, Nakanishi K, Kaji N, Suzuki K, Okamatsu M, et al. Subtyping of avian influenza viruses H1 to H15 on the basis of hemagglutinin genes by PCR assay and molecular determination of pathogenic. *J Clin Microbiol.* 2008;46:3048–55. [PubMed](#)
<http://dx.doi.org/10.1128/JCM.02386-07>
4. Fereidouni SR, Starick E, Grund C, Globig A, Mettenleiter TC, Beer M, et al. Rapid molecular subtyping by reverse transcription polymerase chain reaction of the neuraminidase gene of avian influenza A viruses. *Vet Microbiol.* 2009;135:253–60. [PubMed](#)
<http://dx.doi.org/10.1016/j.vetmic.2008.09.077>
5. Darriba D, Taboada GL, Doallo R, Posada D. jModelTest 2: more models, new heuristics and parallel computing. *Nat Methods.* 2012;9:772. [PubMed](#) <http://dx.doi.org/10.1038/nmeth.2109>
6. Guindon S, Gascuel O. A simple, fast and accurate method to estimate large phylogenies by maximum-likelihood. *Syst Biol.* 2003;52:696–704. [PubMed](#) <http://dx.doi.org/10.1080/10635150390235520>
7. Tamura K, Stecher G, Peterson D, FilipSKI A, Kumar S. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol Biol Evol.* 2013;30:2725–9. [PubMed](#)
<http://dx.doi.org/10.1093/molbev/mst197>

Table. Genetic details of H5 viruses detected in migratory birds of Shanghai, 2013–2014*

Virus name	Collection date	Closest related virus strain/nt identity			HA cleavage site	Pathogenicity
		HA (%)	NA (%)	MP (%)		
A/Common Teal/Shanghai/PD1108-1/2013(mixed)	2013 Nov 8	Novo (96)	6D18 (99)	W24 (96)	RE- - - TR/GLF	Low
A/Common Teal/Shanghai/PD1108-3/2013(H5N8)	2013 Nov 8	w24 (98)	6D18 (99)	S11090 (95)	REKRRKR/GLF	High
A/Common Teal/Shanghai/PD1108-5/2013(mixed)	2013 Nov 8	w24 (99)	6D18 (99)	W24 (99)	REKRRKR/GLF	High
A/Common Teal/Shanghai/PD1108-6/2013(H5N8)	2013 Nov 8	w24 (99)	6D18 (99)	Gochang1 (99)	REKRRKR/GLF	High
A/Common Teal/Shanghai/PD1108-7/2013(mixed)	2013 Nov 8	w24 (99)	6D18 (99)	W24 (99)	REKRRKR/GLF	High
A/Common Teal/Shanghai/PD1108-13/2013(H5N8)	2013 Nov 8	w24 (98)	6D18 (99)	Gochang1 (99)	REKRRKR/GLF	High
A/Common Teal/Shanghai/PD1108-20/2013(H5N8)	2013 Nov 8	w24 (99)	6D18 (99)	Gochang1 (98)	REKRRKR/GLF	High
A/Common Teal/Shanghai/PD1108-24/2013(H5N8)	2013 Nov 8	w24 (99)	6D18 (99)	Gochang1 (99)	REKRRKR/GLF	High
A/Common Teal/Shanghai/PD1108-25/2013(H5N8)	2013 Nov 8	w24 (98)	6D18 (99)	Gochang1 (98)	REKRRKR/GLF	High
A/Common Teal/Shanghai/PD1108-27/2013(H5N8)	2013 Nov 8	w24 (99)	6D18 (99)	Gochang1 (98)	REKRRKR/GLF	High
A/Common Teal/Shanghai/PD1118-1/2013(H5N8)	2013 Nov 18	w24 (99)	6D18 (99)	W24 (98)	REKRRKR/GLF	High
A/Common Teal/Shanghai/PD1121-10/2013(H5N8)	2013 Nov 21	w24 (99)	6D18 (99)	Gochang1 (98)	REKRRKR/GLF	High
A/Common Teal/Shanghai/PD1121-15/2013(H5N8)	2013 Nov 21	w24 (99)	6D18 (99)	W24 (99)	REKRRKR/GLF	High
A/Common Teal/Shanghai/PD1121-18/2013(H5N8)	2013 Nov 21	w24 (99)	6D18 (99)	W24 (99)	REKRRKR/GLF	High
A/Common Teal/Shanghai/PD1121-19/2013(H5N8)	2013 Nov 21	w24 (99)	6D18 (99)	W24 (99)	REKRRKR/GLF	High
A/Falcatad Duck/Shanghai/PD1121-26/2013(H5N8)	2013 Nov 21	w24 (99)	6D18 (99)	W24 (98)	REKRRKR/GLF	High
A/Falcatad Duck/Shanghai/PD1121-27/2013(H5N8)	2013 Nov 21	w24 (99)	6D18 (99)	156 (95)	REKRRKR/GLF	High
A/Spot-billed Duck/Shanghai/PD1202-3/2013(H5N8)	2013 Dec 2	w24 (99)	6D18 (99)	W24 (99)	REKRRKR/GLF	High
A/Common Teal/Shanghai/PD1202-9/2013(H5N8)	2013 Dec 2	w24 (99)	6D18 (99)	W24 (99)	REKRRKR/GLF	High
A/Eurasian Curlew/Shanghai/DT1215-1/2014(H5N8)	2014 Dec 15	H68 (99)	H297 (99)	S005 (99)	RERRRKR/GLF	High
A/Eurasian Curlew/Shanghai/DT1231-1/2014(H5N8)	2014 Dec 31	H68 (99)	H297 (99)	S005 (99)	RERRRKR/GLF	High
A/Eurasian Curlew/Shanghai/DT1231-2/2014(H5N8)	2014 Dec 31	H68 (99)	H297 (99)	S005 (99)	RERRRKR/GLF	High

Virus name	Collection date	Closest related virus strain/nt identity			HA cleavage site	Pathogenicity
		HA (%)	NA (%)	MP (%)		
A/Eurasian Curlew/Shanghai/DT1231-3/2014(H5N8)	2014 Dec 31	H68 (99)	H297 (99)	S005 (99)	RERRRKR/GLF	High
A/Eurasian Curlew/Shanghai/DT1231-4/2014(H5N8)	2014 Dec 31	H68 (99)	H297 (99)	S005 (99)	RERRRKR/GLF	High
A/Eurasian Curlew/Shanghai/DT1231-5/2014(H5N8)	2014 Dec 31	H68 (99)	H297 (99)	S005 (99)	RERRRKR/GLF	High
A/Eurasian Curlew/Shanghai/DT1231-6/2014(H5N8)	2014 Dec 31	H68 (99)	H297 (99)	S005 (99)	RERRRKR/GLF	High
A/Common Teal/Shanghai/PD1108-4/2013(H5N1)	2013 Nov 8	Novo (96)	/	W24 (97)	RE- - - TR/GLF	Low
A/Common Teal/Shanghai/PD1108-8/2013(H5N1)	2013 Nov 8	w24 (95)	/	MHC40-28 (97)	REKRRKR/GLF	High

*HA, hemagglutinin; MP, matrix protein; NA, neuraminidase; species: common teal (*Anas crecca*); falcated duck (*Anas falcata*); spot-billed duck (*Anas poecilorhyncha*); Eurasian curlew (*Numenius arquata*); sampling site: PD, Pudong eastern coastal wetlands; DT, Dongtan Bird Nature Reserve; homologue abbreviations: HA: novo, A/European teal/Novosibirsk/203/2011(H5N1)KF462362; w24, A/duck/Zhejiang/W24/2013(H5N8)KJ476669; H68, A/Baikal teal/Korea/H68/2014(H5N8)KJ509004; NA: 6D18, A/duck/Zhejiang/6D18/2013(H5N8)KJ476674; H297, A/mallard/Korea/H297/2014(H5N8)KJ509134; M: w24, A/duck/Zhejiang/W24/2013(H5N8)KJ476675; S11090, A/duck/Hunan/S11090/2012(H4N6)CY146551; Gochang1, A/breeder duck/Korea/Gochang1/2014(H5N8)KJ413837; 156, A/duck/Hokkaido/156/2006(H11N9)AB428725; S005, A/waterfowl/Korea/S005/2014(H5N8)KJ511815; MHC40-28, A/wild duck/Korea/MHC40-28/2010(H7N7)KC609942; / indicates no data.