

7. Doyle A, Bhalla KS, Jones JM III, Ennis DM. Myocardial involvement in Rocky Mountain spotted fever: a case report and review. *Am J Med Sci*. 2006;332:208–10. <http://dx.doi.org/10.1097/0000441-200610000-00009>
8. Kularatne SA, Rajapakse RP, Wickramasinghe WM, Nanayakkara DM, Budagoda SS, Weerakoon KG, et al. Rickettsioses in the central hills of Sri Lanka: serological evidence of increasing burden of spotted fever group. *Int J Infect Dis*. 2013;17:e988–92. <http://dx.doi.org/10.1016/j.ijid.2013.05.014>
9. Kushawaha A, Brown M, Martin I, Evenhuis W. Hitch-hiker taken for a ride: an unusual case of myocarditis, septic shock and adult respiratory distress syndrome. *BMJ Case Rep*. 2013;2013. pii: bcr2012007155. <http://dx.doi.org/10.1136/bcr-2012-007155>
10. Solary J, Socolovschi C, Aubry C, Brouqui P, Raoult D, Parola P. Detection of *Rickettsia sibirica mongolitimonae* by using cutaneous swab samples and quantitative PCR. *Emerg Infect Dis*. 2014;20:716–8. <http://dx.doi.org/10.3201/eid2004.130575>

Address for correspondence: Pablo Revilla-Marti, Hospital Clinico Universitario Lozano Blesa, Avenida San Juan Bosco 15, Zaragoza 50009, Spain; email: pablorevillamarti@gmail.com

Enteropathogenic *Escherichia coli* O80:H2 in Young Calves with Diarrhea, Belgium

Damien Thiry, Marc Saulmont, Shino Takaki, Klara De Rauw, Jean-Noël Duprez, Atsushi Iguchi, Denis Piérard, Jacques G. Mainil

Author affiliations: University of Liège, Liège, Belgium (D. Thiry, S. Takaki, J.-N. Duprez, J.G. Mainil); Association Régionale de Santé et d'Identification Animales, Ciney, Belgium (M. Saulmont); Universitair Ziekenhuis Brussel, Brussels, Belgium (K. De Rauw, D. Piérard); University of Miyazaki, Miyazaki, Japan (S. Takaki, A. Iguchi)

DOI: <https://doi.org/10.3201/eid2312.170450>

Serogroup O80 was detected in 40% of 104 enteropathogenic *Escherichia coli* isolates from calves with diarrhea from 42 farms in Belgium during 2008–2015. These isolates harbored the *eae-ξ* and *fliC_{H2}* genes, similar to the O80 attaching-effacing Shigatoxigenic *E. coli* isolates found in humans in France. This strain might be emerging.

Enteropathogenic and attaching-effacing Shigatoxigenic *Escherichia coli* (EPEC and AE-STECS) cause bloody diarrhea in humans and young calves. For clarity,

we use the term AE-STECS instead of enterohemorrhagic *E. coli*, similar to a previous publication (1), to refer to STECS isolates from animals that produce attaching-effacing lesions. EPEC and AE-STECS that infect humans are diverse and comprise scores of serotypes (2); in contrast, most calf AE-STECS strains comprise a few serotypes, mostly O5:H-, O26:H11, O111:H-, and O118:H16 (3). The O26:H11 serotype is also the most common among calf EPEC. However, most serotypes that infect calves have not been identified (3). Therefore, during November 2008–June 2015, we conducted a study on 104 EPEC and 153 AE-STECS isolates collected from the feces or the intestinal contents of calves suffering diarrhea (1 isolate/calf) at the Association Régionale de Santé et d'Identification Animales in Ciney, Belgium. Isolates were screened by PCR for genes of the 10 most pathogenic and common calf and human O serogroups: O5, O26, O103, O104, O111, O118, O121, O145, O157, and O165. Confirming published results (3), 80% (122/153) of AE-STECS isolates and only 21% (22/104) of EPEC isolates tested positive for 1 of these (J.G. Mainil, unpub. data) (4). We sought to further characterize this collection of calf EPEC with unidentified O serogroups.

We submitted 9 calf EPECs with unidentified serogroups to the O-typing multiplex PCR platform (5); 6 of 9 EPEC isolates contained the O80 serogroup–encoding gene, and 3 belonged to 3 other O serogroups. We subsequently performed an O80 serogroup–specific PCR (5) of all 31 AE-STECS and 82 EPEC isolates with unidentified serogroups, along with one O80-positive *E. coli* strain and negative controls; 42 EPEC isolates and the O80-positive *E. coli* strain but no AE-STECS isolates or negative controls tested positive.

We further tested the calf EPEC isolates and 3 human Shiga toxin 2–encoding gene (*stx2*)–positive AE-STECS O80 isolates from the STECS National Reference Center (Brussels, Belgium) by PCR for *fliC_{H2}* and *eae-ξ* genes found in human AE-STECS O80 strains. For amplifying *eae-ξ*, we used previously published PCR conditions (6), and for amplifying *fliC_{H2}*, we used primers H2_F (5'-TGATCCGACATCTCCTGATG-3') and H2_R (5'-CCGTCATCACCAATCAACGC-3') and the following thermocycler conditions: initial denaturation at 94°C for 1 min; 30 cycles of denaturation for 30 s at 94°C, annealing for 30 s at 58°C, and elongation for 1 min at 72°C; and final elongation at 72°C for 2 min. All 42 calf EPEC and 3 human AE-STECS isolates tested positive by both PCRS.

Among the 104 calf EPEC isolates, O80:H2 was frequently found (40% were PCR positive) and, thus, could be considered emerging. Indeed, the EPEC O80 isolates were isolated from calves from 42 farms. The yearly EPEC O80:H2 isolation rate varied from 12% in 2009 to 40%–50% during 2010–2013 to as high as 73% for the

Table. Comparison of yearly isolation rates of EPEC O80:H2 and O26 from calves with diarrhea, Belgium, January 2009–June 2015*

Year	Total no.	EPEC	
		O80:H2, %	O26, %
2009	17	24	12
2010	19	47	5
2011	12	50	33
2012	9	33	22
2013	15	47	20
2014	20	25	25
2015	11	73	9
Total	104	41	16

*In 2008, only isolates collected in November and December were studied, and 1 EPEC was identified. EPEC, enteropathogenic *Escherichia coli*.

first 6 months of 2015 (Table). In comparison, the rate for EPEC O26 serotype was 5%–25%. Although prevalence data before 2009 are lacking, EPEC O80 isolates have been found infrequently in animals: 8 in dead poultry (7,8), 1 in a piglet with diarrhea (9), 1 in a healthy cow (9), and 5 in lambs with diarrhea (10). However, even fewer AE-STECS O80 isolates have been found: 2 in healthy cattle (6); 1 in a calf with diarrhea in January 1987 (J.G. Mainil, unpub. data); and 1 in raw cow's milk cheese (9). According to the literature, the O80:H2 serotype might be emerging in France, where human cases of AE-STECS O80:H2 have been reported (9).

Molecular virulotyping results indicate that our calf EPEC O80 isolates appeared to be more closely related to human AE-STECS (because they all harbored *eae*- ξ and *flhC*_{H26}) than to ovine and poultry EPEC O80 (which usually harbor *eae*- β and *flhC*_{H26}) (J.G. Mainil, unpub. data) (8,10). Further studies are needed to characterize these calf EPEC O80:H2 isolates, and the isolation rate of EPEC O80:H2 in calves with diarrhea must be tracked. Additional PCR virulotyping should be performed with our isolates to identify, if present, other EPEC-related virulence genes and extraintestinal *E. coli*-related virulence genes. Some genes could be located on plasmids, like those that were found in AE-STECS O80:H2 patients with bacteremia and internal organ infections (9), although the infected calves from which our isolates were taken did not show evidence of septicemia before or after necropsy. The relationship among calf EPEC O80:H2 isolates (which are all independent isolates, not constituting a single strain until proven otherwise) and between calf and human isolates needs to be further characterized with pulsed-field gel electrophoresis and whole-genome sequencing. The prevalence of O80:H2 EPEC and AE-STECS in healthy cattle at slaughterhouses and farms in Belgium should be examined. Finally, we need to determine whether these calf EPEC O80:H2 isolates are true EPEC, AE-STECS derivatives that have lost *stx* genes, or AE-STECS precursors that could acquire *stx* genes in the future. This work will aid in the detection, prevention, and control of this potentially emerging pathogen.

Acknowledgments

The authors thank Maika Furukawa for her efficient technical support and Melha Mellata for helpful discussions.

The University of Liège provided financial support (Fonds Spéciaux de la Recherche, 2016–2018). S.T. is a trainee veterinary student from the University of Miyazaki, Japan, under the Japan Public-Private Partnership Student Study Abroad Program of the Japan Student Services Organization.

Dr. Thiry is an assistant professor at the Bacteriology Laboratory at the Faculty of Veterinary Medicine of the University of Liège. His areas of research are the identification of enteropathogenic and Shiga toxin-producing *Escherichia coli* isolated from young diarrheic calves and bacteriophages as an alternative treatment for bacterial infections (*Escherichia coli* and *Staphylococcus aureus*) in cattle.

References

- Piérard D, De Greve H, Haesebrouck F, Mainil J. O157:H7 and O104:H4 vero/Shiga toxin-producing *Escherichia coli* outbreaks: respective role of cattle and humans. *Vet Res (Faisalabad)*. 2012;43:13. <http://dx.doi.org/10.1186/1297-9716-43-13>
- Tozzoli R, Scheutz F. Diarrhoeagenic *Escherichia coli* infections in humans. In: Morabito S, editor. *Pathogenic Escherichia coli: molecular and cellular microbiology*. Norwich (UK): Horizon Scientific Press and Caister Academic Press; 2014.
- Mainil JG, Fairbrother JM. Pathogenic *Escherichia coli* in domestic mammals and birds. In: Morabito S, editor. *Pathogenic Escherichia coli: molecular and cellular microbiology*. Norwich (UK): Horizon Scientific Press and Caister Academic Press; 2014.
- Fakih I, Thiry D, Duprez JN, Saulmont M, Iguchi A, Piérard D, et al. Identification of Shiga toxin-producing (STEC) and enteropathogenic (EPEC) *Escherichia coli* in diarrhoeic calves and comparative genomics of O5 bovine and human STECS. *Vet Microbiol*. 2017;202:16–22. <http://dx.doi.org/10.1016/j.vetmic.2016.02.017>
- Iguchi A, Iyoda S, Seto K, Morita-Ishihara T, Scheutz F, Ohnishi M; Pathogenic *E. coli* Working Group in Japan. *Escherichia coli* O-genotyping PCR: a comprehensive and practical platform for molecular O serogrouping. *J Clin Microbiol*. 2015;53:2427–32. <http://dx.doi.org/10.1128/JCM.00321-15>
- Blanco M, Blanco JE, Mora A, Dahbi G, Alonso MP, González EA, et al. Serotypes, virulence genes, and intimin types of Shiga toxin (verotoxin)-producing *Escherichia coli* isolates from cattle in Spain and identification of a new intimin variant gene (*eae*- ξ). *J Clin Microbiol*. 2004;42:645–51. <http://dx.doi.org/10.1128/JCM.42.2.645-651.2004>
- Schouler C, Schaeffer B, Brée A, Mora A, Dahbi G, Biet F, et al. Diagnostic strategy for identifying avian pathogenic *Escherichia coli* based on four patterns of virulence genes. *J Clin Microbiol*. 2012;50:1673–8. <http://dx.doi.org/10.1128/JCM.05057-11>
- Oh JY, Kang MS, An BK, Shin EG, Kim MJ, Kim YJ, et al. Prevalence and characteristics of intimin-producing *Escherichia coli* strains isolated from healthy chickens in Korea. *Poult Sci*. 2012;91:2438–43. <http://dx.doi.org/10.3382/ps.2012-02301>
- Soysal N, Mariani-Kurkdjian P, Smail Y, Liguori S, Gouali M, Loukiadis E, et al. Enterohemorrhagic *Escherichia coli* hybrid pathotype O80:H2 as a new therapeutic challenge. *Emerg Infect Dis*. 2016;22:1604–12. <http://dx.doi.org/10.3201/eid2209.160304>

10. Cid D, Ruiz-Santa-Quiteria JA, Marín I, Sanz R, Orden JA, Amils R, et al. Association between intimin (*eae*) and *espB* gene subtypes in attaching and effacing *Escherichia coli* strains isolated from diarrhoeic lambs and goat kids. *Microbiology*. 2001;147: 2341–53. <http://dx.doi.org/10.1099/00221287-147-8-2341>

Address for correspondence: Jacques G. Mainil, Bacteriology Laboratory, Department of Infectious Diseases, Faculty of Veterinary Medicine, Campus du Sart Tilman B43a, Quartier Vallée 2, Avenue de Cureghem 6, Université de Liège, 4000, Liège, Belgium; email: jg.mainil@uliege.be

Incentives for Bushmeat Consumption and Importation among West African Immigrants, Minnesota, USA

Emily Walz, David Wilson, Jacob C. Stauffer, Danushka Wanduragala, William M. Stauffer, Dominic A. Travis,¹ Jonathan D. Alpern¹

Author affiliations: University of Minnesota, Minneapolis, Minnesota, USA (E. Walz, W.M. Stauffer, D.A. Travis, J.D. Alpern); African Career, Education, and Resources, Inc., Brooklyn Park, Minnesota, USA (D. Wilson); Stillwater High School, Stillwater, Minnesota, USA (J.C. Stauffer); Minnesota Department of Health, St. Paul, Minnesota, USA (D. Wanduragala)

DOI: <https://doi.org/10.3201/eid2312.170563>

The knowledge, attitudes, and practices surrounding bushmeat consumption and importation in the United States are not well described. Focus groups of West African persons living in Minnesota, USA, found that perceived risks are low and unlikely to deter consumers. Incentives for importation and consumption were multifactorial in this community.

Bushmeat hunting and butchery are risk factors for zoonotic disease transmission (1–3). However, less is known about health risks to those who consume products that are already butchered when purchased. Bushmeat in this report refers to meat from wild African animals such as rodents, hooved animals, carnivores, primates, and bats (3).

Thousands of pounds of bushmeat are illegally imported into the United States annually (4), mostly from West

Africa (5). A previous study of bushmeat consumption by African immigrants in the United States described mixed perceptions regarding the risks and benefits of consuming bushmeat (5). Improved understanding of the complex social drivers of these practices is needed to better characterize risk and formulate communication strategies.

To identify the cultural perspectives and knowledge, attitudes, and practices surrounding bushmeat importation and consumption, we held focus groups with members of the Liberian community living in the Minneapolis–St. Paul area of Minnesota, USA. Minneapolis–St. Paul has the largest Liberia-born population in the United States, and ranks fifth in overall African populations in US metropolitan areas (6). Recognizing the history of stigmatization associated with increased risk for Ebola virus among persons from West Africa, we engaged a community-based organization to partner in the planning and execution of this study (7,8). Creating a comfortable environment where participants share personal experiences and insights freely is a key tenet of focus group methodology (9); this partnership was essential in gaining trust and maintaining cultural sensitivity.

Inclusion criteria for participant selection included: 1) minimum age 18 years, 2) self-identification as West African, and 3) willingness to discuss bushmeat in a group setting. The partner organization recruited community members by using a combination of purposeful sampling and social media advertisement and facilitated 3 focus groups (10–12 participants, each for 90 min) in January and February 2016; a designated research team member attended each session. A standard guide for questions was used for each session (online Technical Appendix, <https://wwwnc.cdc.gov/EID/article/23/12/17-0563-Techapp1.pdf>). The University of Minnesota Institutional Review Board approved this study.

Sessions were audio recorded and transcribed; participants were not identified. Nonverbal cues (i.e., gestures, emotions, points of hesitation, nods of agreement) and other participant interactions were added to the transcript by a notetaker. We analyzed the collected data by using a modified grounded theory method with inductive analysis as previously described (10). Two authors (E.W., J.D.A.) analyzed each transcript by using an open and selective coding approach. Subsequently, all transcripts were analyzed together by using axial coding further describing relationships among themes (Table); representative quotes from participants were selected to exemplify a relationship or common theme (9) (Table). We supported validity of findings by using member-checking, triangulation of findings with multiple sources, and peer debriefing (9). Many themes were repeated in all groups; however, this study was limited by inability to confirm that we had reached saturation of perspectives. According to Creswell, it is ideal

¹These senior authors contributed equally to this article.