

39. Saeed AA, Al-Hamdan NA, Fontaine RE. Plague from eating raw camel liver. *Emerg Infect Dis.* 2005;11:1456–7. <https://doi.org/10.3201/eid1109.050081>
40. Brygoo E, Dodin A. Apropos of telluric plague and burrow plague: Madagascar data [in French]. *Bull Soc Pathol Exot.* 1965;1:14–7.
41. Chernin E. Richard Pearson Strong and the Manchurian epidemic of pneumonic plague, 1910–1911. *J Hist Med Allied Sci.* 1989;44:296–319. <https://doi.org/10.1093/jhmas/44.3.296>
42. Kool JL. Risk of person-to-person transmission of pneumonic plague. *Clin Infect Dis.* 2005;40:1166–72. <https://doi.org/10.1086/428617>
43. Didelot X, Whittles LK, Hall I. Model-based analysis of an outbreak of bubonic plague in Cairo in 1801. *J R Soc Interface.* 2017;14:20170160. <https://doi.org/10.1098/rsif.2017.0160>
44. Whittles LK, Didelot X. Epidemiological analysis of the Eyam plague outbreak of 1665–1666. *Proc Biol Sci.* 2016;283:20160618. <https://doi.org/10.1098/rspb.2016.0618>
45. Teh W, Chun W, Pollitzer R. Plague in Manchuria: I. observations made during and after the second Manchurian plague epidemic of 1920–21, II. The rôle of the tarabagan in the epidemiology of plague. *J Hyg (Lond).* 1923;21:307–58. <https://doi.org/10.1017/S0022172400031521>
46. Strong R, Teague O. Studies on pneumonic plague and plague immunisation. II the method of transmission of the infection in pneumonic plague and manner of spread of the disease during the epidemic. *The Philippine Journal of Tropical Medicine.* 1912;7:137–56.

Address for correspondence: Sophie Jullien, Institute of Global Health, University of Barcelona, Carrer Rosselló 132, 5-2, 08036 Barcelona, Spain; email: [sophjullien@gmail.com](mailto:sophjullien@gmail.com)

## etymologia

### *Culex quinquefasciatus* [*'kyō leks 'kwinkwə fa she 'ah tus*]

Sarah Anne J. Guagliardo, Rebecca S. Levine

In 1823, the American entomologist Thomas Say described *Culex* (Latin for “gnat”) *quinquefasciatus*, which he collected along the Mississippi River. Originally written as “C. 5-fasciatus,” the name refers to 5 (“quinque”) black, broad, transverse bands (“fasciatus” or “fasciae”) on the mosquito’s dorsal abdomen. The name remains despite later revelations of more than 5 fasciae, thanks to improved microscopy. Although *quinquefasciatus* is the official scientific name, there are at least 5 synonymous names for this species.

Say described this species as “exceedingly numerous and troublesome.” “Quinx” are among the world’s most abundant peridomestic mosquitoes, earning the nickname “southern house mosquito.” *Cx. quinquefasciatus* is found throughout subtropical and tropical areas worldwide, except for exceedingly dry or cold regions. This mosquito is a principal vector of many pathogens, transmitting the phlebovirus Rift Valley fever virus and the 2 flaviviruses St. Louis encephalitis virus and West Nile virus, in addition to filarial worms and avian malarial parasites.




Figure. Female *Culex quinquefasciatus* mosquito. Image credit: CDC Public Health Image Library, 1976.

#### Sources

- Belkin J. *Quinquefasciatus* or *Fatigans* for the tropical (Southern) house mosquito (Diptera: Culicidae). *Proc Entomol Soc Wash.* 1977;79:45–52.
- Farajollahi A, Fonseca DM, Kramer LD, Marm Kilpatrick A. “Bird biting” mosquitoes and human disease: a review of the role of *Culex pipiens* complex mosquitoes in epidemiology. *Infect Genet Evol.* 2011;11:1577–85. <https://doi.org/10.1016/j.meegid.2011.08.013>
- Harrison BA, Byrd BD, Sither CB, Whitt PB. *The Mosquitoes of the Mid-Atlantic Region: An Identification Guide.* Cullowhee (NC): Western Carolina University; 2016.
- Say T. Descriptions of dipterous insects of the United States. *Journal of the Academy of Natural Sciences.* 1823;3:9–54.
- University of Florida, Department of Entomology and Nematology. Featured creatures. [cited 2021 Mar 3]. [http://entnemdept.ufl.edu/creatures/aquatic/southern\\_house\\_mosquito.htm](http://entnemdept.ufl.edu/creatures/aquatic/southern_house_mosquito.htm)

Author affiliation: Centers for Disease Control and Prevention, Atlanta, Georgia, USA

Address for correspondence: Sarah Anne J. Guagliardo, Centers for Disease Control and Prevention, 1600 Clifton Rd NE, Atlanta, GA 30327-4027, USA; email: [sguagliardo@cdc.gov](mailto:sguagliardo@cdc.gov)

DOI: <https://doi.org/10.3201/eid2708.ET2708>